



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 03:44 AM UTC

PDB ID : 2XAO / pdb_00002xao
Title : Inositol 1,3,4,5,6-pentakisphosphate 2-kinase from *A. thaliana* in complex with IP5
Authors : Gonzalez, B.; Banos-Sanz, J.I.; Villate, M.; Brearley, C.A.; Sanz-Aparicio, J.
Deposited on : 2010-03-31
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	451	2% 67% 20% • 10%
1	B	451	4% 71% 19% • 6%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6668 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called INOSITOL-PENTAKISPHOSPHATE 2-KINASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	404	Total	C	N	O	S	0	0	0
			3224	2052	545	614	13			
1	B	424	Total	C	N	O	S	0	0	0
			3378	2148	572	645	13			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	54	SER	ALA	conflict	UNP Q93YN9
A	90	GLN	LYS	conflict	UNP Q93YN9
A	157	THR	SER	SEE REMARK 999	UNP Q93YN9
A	204	ILE	ASN	conflict	UNP Q93YN9
A	224	ARG	SER	conflict	UNP Q93YN9
A	321	CYS	SER	conflict	UNP Q93YN9
A	325	ILE	LEU	conflict	UNP Q93YN9
A	337	ARG	LYS	SEE REMARK 999	UNP Q93YN9
B	54	SER	ALA	conflict	UNP Q93YN9
B	90	GLN	LYS	conflict	UNP Q93YN9
B	157	THR	SER	SEE REMARK 999	UNP Q93YN9
B	204	ILE	ASN	conflict	UNP Q93YN9
B	224	ARG	SER	conflict	UNP Q93YN9
B	321	CYS	SER	conflict	UNP Q93YN9
B	325	ILE	LEU	conflict	UNP Q93YN9
B	337	ARG	LYS	conflict	UNP Q93YN9

- Molecule 2 is MYO-INOSITOL-(1,3,4,5,6)-PENTAKISPHOSPHATE (CCD ID: 5MY) (formula: C₆H₁₇O₂₁P₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	O	P	0	0
			32	6	21	5		
2	B	1	Total	C	O	P	0	0
			32	6	21	5		


- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		
3	B	1	Total	Zn	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: INOSITOL-PENTAKISPHOSPHATE 2-KINASE



• Molecule 1: INOSITOL-PENTAKISPHOSPHATE 2-KINASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.76Å 112.74Å 144.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	88.74 – 2.90 88.74 – 2.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (88.74-2.90) 100.0 (88.74-2.90)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.31 (at 2.91Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.225 , 0.291 (Not available) , 0.260	Depositor DCC
R_{free} test set	1141 reflections (5.19%)	wwPDB-VP
Wilson B-factor (Å ²)	42.7	Xtriage
Anisotropy	0.092	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 34.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	6668	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, 5MY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.61	1/3280 (0.0%)	0.90	0/4416
1	B	0.62	1/3441 (0.0%)	0.96	5/4640 (0.1%)
All	All	0.62	2/6721 (0.0%)	0.93	5/9056 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	153	SER	C-O	7.71	1.39	1.23
1	A	94	ILE	CA-CB	7.41	1.57	1.54

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	337	ARG	CA-C-N	8.68	130.69	119.84
1	B	337	ARG	C-N-CA	8.68	130.69	119.84
1	B	384	GLU	CA-C-N	6.92	127.30	119.83
1	B	384	GLU	C-N-CA	6.92	127.30	119.83
1	B	177	SER	N-CA-C	5.73	118.37	110.35

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3224	0	3240	57	0
1	B	3378	0	3393	55	0
2	A	32	0	7	1	0
2	B	32	0	7	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
All	All	6668	0	6647	110	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (110) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:101:HIS:CD2	1:A:303:ARG:HH22	1.75	1.03
1:B:40:ARG:HD3	1:B:146:LEU:CD1	2.10	0.81
1:A:126:ARG:HH21	1:A:131:VAL:HA	1.49	0.77
1:A:67:HIS:O	1:A:70:ARG:HD2	1.88	0.73
1:A:15:TYR:HA	1:A:25:LEU:HD12	1.71	0.72
1:A:319:ILE:HD13	1:A:420:TYR:HB2	1.70	0.72
1:B:8:LYS:HD2	1:B:8:LYS:H	1.54	0.71
1:A:101:HIS:HD2	1:A:303:ARG:HH22	1.31	0.71
1:A:101:HIS:CD2	1:A:303:ARG:NH2	2.56	0.70
1:B:40:ARG:HD3	1:B:146:LEU:HD11	1.72	0.70
1:B:277:GLN:H	1:B:397:GLN:HE22	1.38	0.70
1:A:153:SER:HB3	1:A:402:LYS:NZ	2.07	0.69
1:B:348:LEU:HG	1:B:352:GLU:HG2	1.75	0.69
1:B:204:ILE:HG22	1:B:204:ILE:O	1.94	0.67
1:A:126:ARG:NH2	1:A:131:VAL:HA	2.09	0.67
1:B:14:ILE:HD12	1:B:15:TYR:H	1.60	0.66
1:B:222:LYS:NZ	1:B:222:LYS:HB3	2.10	0.66
1:B:101:HIS:HD2	1:B:303:ARG:NH2	1.94	0.65
1:B:226:LEU:HA	1:B:292:SER:OG	1.97	0.64
1:A:422:LEU:O	1:A:426:ILE:HG12	1.96	0.64
1:B:345:LEU:O	1:B:348:LEU:HB2	1.98	0.63
1:B:74:GLU:N	1:B:74:GLU:OE1	2.33	0.62
1:B:16:ARG:NH2	1:B:148:ASP:OD1	2.32	0.62
1:A:295:VAL:HG13	1:A:301:LEU:HD22	1.82	0.61
1:A:129:TRP:HB2	1:B:178:ARG:HG2	1.82	0.60
1:B:101:HIS:HD2	1:B:303:ARG:HH22	1.49	0.60
1:B:215:LEU:O	1:B:219:SER:HB3	2.01	0.60
1:A:47:ASN:OD1	1:B:188:THR:HG21	2.03	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:113:GLU:O	1:B:117:CYS:HB2	2.03	0.58
1:A:275:PHE:HD1	1:A:276:ILE:HD12	1.68	0.58
1:A:101:HIS:HD2	1:A:303:ARG:NH2	1.99	0.58
1:A:325:ILE:HD11	1:A:354:LEU:HA	1.86	0.57
1:A:190:VAL:HB	1:A:195:MET:HE2	1.87	0.55
1:A:97:LEU:HG	1:A:307:ILE:CD1	2.38	0.54
1:A:153:SER:HB3	1:A:402:LYS:HZ1	1.71	0.54
1:B:277:GLN:H	1:B:397:GLN:NE2	2.05	0.54
1:B:39:ILE:HD12	1:B:41:ILE:HG13	1.90	0.53
1:B:52:LYS:HB2	1:B:58:VAL:N	2.23	0.53
1:A:153:SER:HB3	1:A:402:LYS:HZ2	1.73	0.53
1:A:215:LEU:O	1:A:219:SER:HB3	2.09	0.52
2:A:500:5MY:O35	2:A:500:5MY:O34	2.28	0.52
1:A:18:GLU:OE1	1:A:126:ARG:HG2	2.11	0.51
1:B:393:LYS:N	1:B:394:PRO:HD2	2.26	0.51
1:B:39:ILE:CD1	1:B:41:ILE:HG13	2.41	0.50
1:A:168:LYS:HE3	1:A:170:LYS:HD3	1.92	0.50
1:A:94:ILE:HD13	1:A:102:VAL:HB	1.92	0.50
1:B:251:GLY:HA2	1:B:257:THR:HG21	1.94	0.50
1:B:198:LEU:HD12	1:B:430:TYR:HD1	1.75	0.50
1:B:222:LYS:HB3	1:B:222:LYS:HZ3	1.77	0.49
1:A:295:VAL:HG12	1:A:296:TYR:N	2.28	0.49
1:A:319:ILE:HD13	1:A:420:TYR:CB	2.41	0.49
1:A:112:LYS:HG3	1:A:138:THR:O	2.13	0.49
1:A:325:ILE:HG12	1:A:354:LEU:HD23	1.95	0.48
1:B:300:VAL:HG11	1:B:403:VAL:HG21	1.94	0.48
1:B:345:LEU:HA	1:B:348:LEU:HD22	1.95	0.48
1:A:199:LEU:HD21	1:A:425:LYS:HE3	1.95	0.48
1:B:187:LYS:HB3	1:B:195:MET:HE1	1.96	0.48
1:B:172:GLY:HA3	1:B:218:PHE:CD2	2.49	0.47
1:B:25:LEU:HB2	1:B:39:ILE:CG2	2.45	0.47
1:A:197:GLN:HB3	1:A:209:GLU:HG3	1.97	0.47
1:A:72:ASN:HB3	1:A:74:GLU:OE2	2.15	0.47
1:B:341:ALA:HA	1:B:344:SER:HB2	1.97	0.47
1:B:39:ILE:HD11	1:B:143:ALA:HB1	1.97	0.47
1:A:119:ASP:HB2	1:A:136:VAL:CG2	2.45	0.46
1:B:226:LEU:HA	1:B:292:SER:HG	1.80	0.46
1:B:40:ARG:CD	1:B:146:LEU:CD1	2.88	0.46
1:B:184:ASN:O	1:B:187:LYS:HG2	2.15	0.45
1:A:200:LYS:HB3	1:A:206:ILE:HG12	1.98	0.45
1:A:170:LYS:HA	1:A:367:LYS:O	2.17	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:337:ARG:HA	1:B:338:PRO:HD3	1.65	0.45
1:B:308:GLN:NE2	1:B:367:LYS:HA	2.32	0.44
1:B:325:ILE:HD12	1:B:354:LEU:HD23	2.00	0.44
1:A:46:ARG:NH1	1:B:182:LYS:HG2	2.32	0.44
1:B:341:ALA:O	1:B:345:LEU:N	2.49	0.44
1:B:351:ASP:HA	1:B:354:LEU:HD12	1.99	0.44
1:B:186:LEU:HD12	1:B:186:LEU:HA	1.90	0.44
1:A:216:ASP:OD1	1:A:224:ARG:NE	2.45	0.44
1:A:186:LEU:HD12	1:A:186:LEU:HA	1.83	0.44
1:A:79:PRO:HG2	1:A:83:VAL:HG21	1.99	0.43
1:A:300:VAL:HG11	1:A:403:VAL:HG11	2.00	0.43
1:A:163:ILE:HG13	1:A:275:PHE:CD1	2.54	0.43
1:B:165:VAL:HG22	1:B:242:VAL:HG22	2.01	0.43
1:B:204:ILE:O	1:B:204:ILE:CG2	2.65	0.43
1:A:172:GLY:HA3	1:A:218:PHE:CD2	2.53	0.43
1:B:97:LEU:HD13	1:B:307:ILE:HG13	2.01	0.43
1:A:166:GLU:HB3	1:A:241:ARG:HB2	2.00	0.42
1:A:38:VAL:HG13	1:A:148:ASP:HB2	2.00	0.42
1:A:93:ILE:HG23	1:A:97:LEU:HD12	2.00	0.42
1:B:16:ARG:HB3	1:B:24:VAL:O	2.20	0.42
1:B:198:LEU:HD12	1:B:430:TYR:CD1	2.52	0.42
1:A:73:ASN:O	1:A:76:ILE:HG13	2.19	0.42
1:A:148:ASP:OD1	1:A:148:ASP:C	2.62	0.42
1:A:212:TYR:CE2	1:A:214:PRO:HG3	2.54	0.42
1:A:322:TYR:HE2	1:A:417:GLU:HG3	1.84	0.42
1:B:212:TYR:CE2	1:B:214:PRO:HG3	2.55	0.42
1:B:74:GLU:O	1:B:87:ARG:NH1	2.51	0.41
1:A:18:GLU:OE2	1:A:126:ARG:NH1	2.53	0.41
1:A:185:MET:O	1:A:188:THR:HB	2.20	0.41
1:A:342:GLU:N	1:A:342:GLU:OE1	2.54	0.41
1:A:251:GLY:HA2	1:A:257:THR:HG21	2.03	0.41
1:A:295:VAL:HG23	1:A:373:ILE:HD11	2.03	0.41
1:A:322:TYR:CE2	1:A:417:GLU:HG3	2.56	0.41
1:B:25:LEU:HB2	1:B:39:ILE:HG22	2.01	0.41
1:A:97:LEU:HD22	1:A:405:PHE:CE1	2.56	0.41
1:B:36:GLY:HA2	1:B:151:LEU:HD11	2.03	0.41
1:B:101:HIS:CD2	1:B:303:ARG:HH22	2.34	0.41
1:B:313:LEU:O	1:B:314:ASP:HB3	2.21	0.41
1:A:287:PHE:O	1:A:290:LEU:HB3	2.21	0.41
1:A:90:GLN:HA	1:A:94:ILE:HG13	2.03	0.40
1:B:165:VAL:C	1:B:372:MET:HE3	2.46	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	392/451 (87%)	374 (95%)	16 (4%)	2 (0%)	24	54
1	B	418/451 (93%)	396 (95%)	21 (5%)	1 (0%)	43	72
All	All	810/902 (90%)	770 (95%)	37 (5%)	3 (0%)	30	59

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	49	LYS
1	A	70	ARG
1	A	177	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	362/399 (91%)	330 (91%)	32 (9%)	9	29
1	B	378/399 (95%)	328 (87%)	50 (13%)	4	13
All	All	740/798 (93%)	658 (89%)	82 (11%)	6	20

All (82) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	MET
1	A	5	LEU
1	A	25	LEU
1	A	47	ASN
1	A	72	ASN
1	A	94	ILE
1	A	96	LEU
1	A	136	VAL
1	A	164	SER
1	A	174	LEU
1	A	176	THR
1	A	186	LEU
1	A	188	THR
1	A	198	LEU
1	A	206	ILE
1	A	207	SER
1	A	223	GLU
1	A	234	SER
1	A	238	ASN
1	A	244	LEU
1	A	247	SER
1	A	276	ILE
1	A	295	VAL
1	A	301	LEU
1	A	310	LEU
1	A	343	LEU
1	A	348	LEU
1	A	350	LEU
1	A	351	ASP
1	A	372	MET
1	A	391	SER
1	A	435	LYS
1	B	4	ILE
1	B	5	LEU
1	B	8	LYS
1	B	16	ARG
1	B	30	SER
1	B	39	ILE
1	B	52	LYS
1	B	75	LEU
1	B	83	VAL
1	B	97	LEU
1	B	103	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	117	CYS
1	B	120	LYS
1	B	124	LYS
1	B	125	GLN
1	B	145	ILE
1	B	151	LEU
1	B	163	ILE
1	B	176	THR
1	B	186	LEU
1	B	189	SER
1	B	196	HIS
1	B	198	LEU
1	B	207	SER
1	B	208	GLU
1	B	222	LYS
1	B	234	SER
1	B	237	GLN
1	B	253	SER
1	B	276	ILE
1	B	285	GLU
1	B	290	LEU
1	B	292	SER
1	B	300	VAL
1	B	301	LEU
1	B	334	LYS
1	B	344	SER
1	B	348	LEU
1	B	352	GLU
1	B	355	LYS
1	B	371	ILE
1	B	372	MET
1	B	373	ILE
1	B	377	SER
1	B	379	ASN
1	B	398	THR
1	B	403	VAL
1	B	418	SER
1	B	421	LYS
1	B	437	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	22	ASN
1	A	47	ASN
1	A	91	ASN
1	A	101	HIS
1	A	135	ASN
1	A	149	HIS
1	A	238	ASN
1	A	239	ASN
1	A	327	ASN
1	B	22	ASN
1	B	42	GLN
1	B	147	ASN
1	B	149	HIS
1	B	282	HIS
1	B	397	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	5MY	A	500	-	32,32,32	0.68	0	53,53,53	1.15	5 (9%)
2	5MY	B	500	-	32,32,32	0.80	0	53,53,53	1.08	2 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	5MY	A	500	-	-	5/25/49/49	0/1/1/1
2	5MY	B	500	-	-	2/25/49/49	0/1/1/1

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	500	5MY	O15-C5-C6	-2.88	102.64	108.76
2	A	500	5MY	O15-C5-C6	-2.67	103.09	108.76
2	B	500	5MY	O24-P4-O44	2.65	117.74	107.80
2	A	500	5MY	O21-P1-O31	2.41	116.85	107.80
2	A	500	5MY	O24-P4-O44	2.37	116.67	107.80
2	A	500	5MY	O45-P5-O25	2.35	116.60	107.80
2	A	500	5MY	O43-P3-O23	2.12	115.73	107.80

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	500	5MY	C2-C3-O13-P3
2	B	500	5MY	C1-O11-P1-O41
2	B	500	5MY	C4-O14-P4-O34
2	A	500	5MY	C3-O13-P3-O43
2	A	500	5MY	C4-C3-O13-P3
2	A	500	5MY	C1-O11-P1-O21
2	A	500	5MY	C4-O14-P4-O34


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	500	5MY	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	404/451 (89%)	0.07	10 (2%) 58 48	24, 38, 57, 69	0
1	B	424/451 (94%)	0.20	20 (4%) 36 28	24, 38, 61, 81	0
All	All	828/902 (91%)	0.13	30 (3%) 46 38	24, 38, 58, 81	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	47	ASN	5.0
1	B	254	GLY	4.9
1	A	160	GLY	4.8
1	B	381	TRP	3.8
1	B	50	ALA	3.3
1	B	256	SER	3.0
1	A	59	SER	2.9
1	B	48	ASP	2.9
1	A	3	MET	2.9
1	A	103	ASP	2.9
1	B	153	SER	2.8
1	B	253	SER	2.8
1	A	436	ALA	2.8
1	B	45	ARG	2.6
1	B	51	ILE	2.6
1	B	338	PRO	2.6
1	A	47	ASN	2.6
1	B	52	LYS	2.5
1	B	341	ALA	2.5
1	A	343	LEU	2.3
1	B	103	ASP	2.3
1	A	335	GLU	2.2
1	B	336	GLY	2.2
1	B	3	MET	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	377	SER	2.1
1	A	255	GLU	2.1
1	B	152	PHE	2.0
1	A	48	ASP	2.0
1	B	280	ASP	2.0
1	B	124	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	5MY	A	500	32/32	0.98	0.06	13,21,26,28	0
2	5MY	B	500	32/32	0.98	0.06	17,21,26,27	0
3	ZN	B	700	1/1	0.99	0.04	45,45,45,45	0
3	ZN	A	700	1/1	1.00	0.03	40,40,40,40	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 5MY A 500:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 5MY B 500:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.