



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2026 – 10:29 PM UTC

PDB ID : 4XA5 / pdb_00004xa5
Title : Crystal structure of the pre-catalytic ternary complex of DNA polymerase lambda with a templating A and an incoming 8-oxo-dGTP
Authors : Burak, M.J.; Guja, K.E.; Garcia-Diaz, M.
Deposited on : 2014-12-12
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	325	14% 83% 12% • •
2	D	4	100%
3	P	6	50% 50%
4	T	11	64% 36%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
8	GOL	A	608	-	-	X	-

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 3118 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase lambda.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	314	Total	As	C	N	O	S	0	1	0
			2388	1	1511	425	440	11			

- Molecule 2 is a DNA chain called DOWNSTREAM PRIMER DNA (5'-D(P*GP*CP*CP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	4	Total	C	N	O	P	0	0	0
			83	38	16	25	4			


- Molecule 3 is a DNA chain called PRIMER DNA (5'-D(*CP*AP*GP*TP*AP*(DOC))-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	P	6	Total	C	N	O	P	0	0	0
			118	58	23	32	5			

- Molecule 4 is a DNA chain called TEMPLATE DNA (5'-D(*CP*GP*GP*CP*AP*GP*TP*AP*CP*TP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	T	11	Total	C	N	O	P	0	0	0
			224	107	43	64	10			

- Molecule 5 is 8-OXO-2'-DEOXYGUANOSINE-5'-TRIPHOSPHATE (CCD ID: 8DG) (formula: C₁₀H₁₆N₅O₁₄P₃).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			32	10	5	14	3		

- Molecule 6 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	2	Total Mg 2 2	0	0

- Molecule 7 is ACETATE ION (CCD ID: ACT) (formula: $\text{C}_2\text{H}_3\text{O}_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total C O 4 2 2	0	0
7	A	1	Total C O 4 2 2	0	0
7	A	1	Total C O 4 2 2	0	0

- Molecule 8 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total C O 6 3 3	0	0
8	A	1	Total C O 6 3 3	0	0


- Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	178	Total O 178 178	0	0
9	D	6	Total O 6 6	0	0
9	P	24	Total O 24 24	0	0
9	T	39	Total O 39 39	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA polymerase lambda




- Molecule 2: DOWNSTREAM PRIMER DNA (5'-D(P*GP*CP*CP*G)-3')



- Molecule 3: PRIMER DNA (5'-D(*CP*AP*GP*TP*AP*(DOC))-3')



- Molecule 4: TEMPLATE DNA (5'-D(*CP*GP*GP*CP*AP*GP*TP*AP*CP*TP*G)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	55.97Å 62.34Å 141.61Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.79 – 1.90 46.79 – 1.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (46.79-1.90) 100.0 (46.79-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.23 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.199 , 0.227 0.210 , 0.237	Depositor DCC
R_{free} test set	1971 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	32.2	Xtriage
Anisotropy	0.167	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 52.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3118	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CAS, 8DG, GOL, MG, DOC, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.20	2/2431 (0.1%)	1.12	4/3294 (0.1%)
2	D	0.30	0/92	0.52	0/138
3	P	0.38	0/112	0.70	0/171
4	T	0.49	0/251	1.03	2/386 (0.5%)
All	All	1.11	2/2886 (0.1%)	1.08	6/3989 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	574	ASP	CA-C	-6.54	1.46	1.53
1	A	481	GLY	N-CA	5.11	1.51	1.44

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	T	2	DG	C2'-C3'-O3'	-6.14	102.29	111.50
1	A	275	ARG	N-CA-C	-5.99	104.83	111.36
1	A	332	VAL	O-C-N	5.15	123.72	120.42
1	A	515	SER	N-CA-C	5.10	116.53	111.07
1	A	442	GLY	N-CA-C	5.07	122.11	115.47
4	T	1	DC	C5'-C4'-O4'	5.06	116.99	109.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2388	0	2291	52	0
2	D	83	0	45	5	0
3	P	118	0	69	1	0
4	T	224	0	125	3	0
5	A	32	0	12	0	0
6	A	2	0	0	0	0
7	A	12	0	9	1	0
8	A	12	0	16	5	0
9	A	178	0	0	6	1
9	D	6	0	0	2	0
9	P	24	0	0	0	0
9	T	39	0	0	2	1
All	All	3118	0	2567	58	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:568:ARG:NH2	1:A:575:TRP:CZ2	2.29	0.99
1:A:568:ARG:HH22	1:A:575:TRP:HZ2	1.14	0.92
1:A:404:GLN:OE1	9:A:701:HOH:O	1.88	0.92
1:A:568:ARG:NH2	1:A:575:TRP:HZ2	1.68	0.90
1:A:308:ARG:NH1	2:D:1:DG:OP3	2.16	0.79
1:A:494:VAL:HG12	8:A:608:GOL:H32	1.68	0.74
1:A:429:ASP:HB3	1:A:506:PHE:CZ	2.25	0.72
1:A:429:ASP:OD1	9:A:746:HOH:O	2.09	0.70
1:A:568:ARG:NH2	1:A:575:TRP:CE2	2.58	0.69
1:A:568:ARG:CZ	1:A:575:TRP:HZ2	2.07	0.68
2:D:1:DG:OP2	9:D:104:HOH:O	2.12	0.66
1:A:431:LEU:C	1:A:431:LEU:HD23	2.22	0.65
1:A:494:VAL:HG12	8:A:608:GOL:C3	2.26	0.64
1:A:568:ARG:NH2	1:A:575:TRP:NE1	2.45	0.64
1:A:568:ARG:HH22	1:A:575:TRP:NE1	1.98	0.62
1:A:568:ARG:NH2	1:A:575:TRP:HE1	1.99	0.61
1:A:568:ARG:NH1	1:A:575:TRP:HZ2	1.99	0.59
1:A:316:ILE:O	1:A:319:SER:O	2.22	0.58
1:A:254:LEU:HD12	1:A:286:LEU:HB3	1.86	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:471:GLN:HG2	8:A:608:GOL:H2	1.86	0.56
1:A:269:VAL:O	1:A:346[A]:THR:HG23	2.05	0.56
1:A:575:TRP:CD1	1:A:575:TRP:N	2.73	0.53
1:A:427:ASP:OD2	9:A:746:HOH:O	2.18	0.53
1:A:431:LEU:HD23	1:A:432:ILE:N	2.24	0.52
2:D:3:DC:H2'	2:D:4:DG:C8	2.44	0.52
1:A:568:ARG:NH1	1:A:575:TRP:CZ2	2.74	0.51
1:A:429:ASP:HB3	1:A:506:PHE:CE2	2.45	0.51
1:A:516:MET:HG2	1:A:527:LEU:CD2	2.41	0.51
1:A:521:LYS:NZ	4:T:4:DC:OP1	2.38	0.50
4:T:8:DA:OP2	9:T:132:HOH:O	2.19	0.50
1:A:269:VAL:O	1:A:346[B]:THR:HG22	2.12	0.50
2:D:2:DC:H2'	2:D:3:DC:C6	2.47	0.49
1:A:275:ARG:NH1	2:D:1:DG:OP2	2.46	0.49
1:A:271:GLY:CA	1:A:346[A]:THR:HG21	2.43	0.49
1:A:466:GLU:CB	9:A:831:HOH:O	2.62	0.48
1:A:267:TYR:CZ	1:A:275:ARG:HD2	2.49	0.47
4:T:8:DA:OP2	9:T:139:HOH:O	2.20	0.47
1:A:291:LYS:HG2	1:A:292:PRO:O	2.15	0.47
1:A:431:LEU:C	1:A:431:LEU:CD2	2.89	0.46
1:A:253:ASN:O	1:A:253:ASN:ND2	2.36	0.46
1:A:308:ARG:NH1	9:D:104:HOH:O	2.50	0.45
1:A:434:HIS:CE1	1:A:436:ASP:HB3	2.52	0.44
1:A:254:LEU:HA	1:A:254:LEU:HD23	1.70	0.44
1:A:254:LEU:CD1	1:A:286:LEU:HB3	2.48	0.43
1:A:471:GLN:CG	8:A:608:GOL:H2	2.47	0.43
1:A:271:GLY:N	1:A:346[A]:THR:HG21	2.34	0.43
1:A:427:ASP:C	1:A:427:ASP:OD1	2.61	0.43
1:A:568:ARG:HH21	1:A:575:TRP:HE1	1.64	0.43
1:A:253:ASN:O	1:A:254:LEU:HG	2.17	0.43
1:A:534:THR:HG22	1:A:535:ALA:O	2.19	0.43
1:A:493:VAL:O	8:A:608:GOL:O2	2.29	0.42
1:A:534:THR:HG22	1:A:535:ALA:N	2.34	0.42
1:A:472:LYS:HE3	9:A:769:HOH:O	2.19	0.41
1:A:534:THR:HB	1:A:549:ARG:H	1.85	0.41
1:A:442:GLY:N	7:A:605:ACT:H3	2.35	0.41
3:P:4:DT:H2''	3:P:5:DA:O5'	2.20	0.41
1:A:557:LYS:HG3	9:A:827:HOH:O	2.21	0.40
1:A:557:LYS:HG2	1:A:567:TYR:CD1	2.57	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:A:703:HOH:O	9:T:102:HOH:O[4_555]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	310/325 (95%)	296 (96%)	14 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	236/272 (87%)	231 (98%)	5 (2%)	47	44

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	253	ASN
1	A	408	SER
1	A	411	LEU
1	A	431	LEU
1	A	506	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	350	GLN
1	A	355	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	DOC	P	6	3,4	16,19,20	1.06	1 (6%)	20,26,29	1.26	3 (15%)
1	CAS	A	300	1	5,8,9	1.02	0	1,9,11	0.75	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	DOC	P	6	3,4	-	0/7/18/19	0/2/2/2
1	CAS	A	300	1	-	0/0/7/9	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	P	6	DOC	C6-C5	2.83	1.41	1.35

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	P	6	DOC	O4'-C1'-N1	-2.94	102.64	107.86

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	P	6	DOC	C3'-C2'-C1'	2.64	105.92	102.87
3	P	6	DOC	C2'-C1'-N1	2.54	117.21	112.40

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
8	GOL	A	608	-	5,5,5	0.80	0	5,5,5	0.93	0
7	ACT	A	606	-	3,3,3	0.75	0	3,3,3	0.83	0
8	GOL	A	607	-	5,5,5	0.28	0	5,5,5	0.79	0
7	ACT	A	605	-	3,3,3	0.85	0	3,3,3	0.53	0
7	ACT	A	604	-	3,3,3	0.76	0	3,3,3	0.65	0
5	8DG	A	601	6	33,34,34	2.14	8 (24%)	44,54,54	2.42	12 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	GOL	A	607	-	-	3/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	GOL	A	608	-	-	4/4/4/4	-
5	8DG	A	601	6	-	5/22/34/34	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	601	8DG	C5-C4	7.31	1.47	1.37
5	A	601	8DG	PB-O3B	4.47	1.64	1.59
5	A	601	8DG	PA-O3A	3.92	1.63	1.59
5	A	601	8DG	C6-N1	-3.47	1.32	1.38
5	A	601	8DG	C8-N9	-2.83	1.35	1.40
5	A	601	8DG	O8-C8	2.81	1.28	1.23
5	A	601	8DG	C5-N7	-2.18	1.34	1.37
5	A	601	8DG	C4-N9	-2.06	1.35	1.39

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	601	8DG	N7-C8-N9	9.27	116.92	106.61
5	A	601	8DG	N9-C4-N3	6.09	133.74	126.13
5	A	601	8DG	C5-N7-C8	-4.80	102.86	109.47
5	A	601	8DG	C2-N3-C4	3.88	118.98	112.30
5	A	601	8DG	O8-C8-N9	-3.76	120.88	126.00
5	A	601	8DG	O6-C6-C5	-3.42	119.02	127.26
5	A	601	8DG	O6-C6-N1	2.39	124.61	120.11
5	A	601	8DG	O3A-PB-O1B	-2.34	103.65	110.70
5	A	601	8DG	O2B-PB-O1B	2.34	123.34	112.44
5	A	601	8DG	O8-C8-N7	-2.28	122.19	126.47
5	A	601	8DG	C2'-C3'-C4'	-2.12	98.51	102.80
5	A	601	8DG	C4'-O4'-C1'	-2.03	104.69	109.51

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	601	8DG	O4'-C1'-N9-C4
5	A	601	8DG	C2'-C1'-N9-C4
8	A	607	GOL	C1-C2-C3-O3
8	A	608	GOL	O1-C1-C2-C3
8	A	608	GOL	O1-C1-C2-O2
8	A	608	GOL	C1-C2-C3-O3

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
8	A	607	GOL	O2-C2-C3-O3
8	A	608	GOL	O2-C2-C3-O3
8	A	607	GOL	O1-C1-C2-O2
5	A	601	8DG	O4'-C1'-N9-C8
5	A	601	8DG	PB-O3A-PA-O2A
5	A	601	8DG	C2'-C1'-N9-C8

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	A	608	GOL	5	0
7	A	605	ACT	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	313/325 (96%)	0.83	45 (14%) 6 6	15, 39, 80, 97	1 (0%)
2	D	4/4 (100%)	0.23	0 100 100	38, 39, 39, 42	0
3	P	5/6 (83%)	-0.71	0 100 100	24, 24, 29, 32	0
4	T	11/11 (100%)	0.18	0 100 100	28, 37, 60, 66	0
All	All	333/346 (96%)	0.78	45 (13%) 7 7	15, 39, 78, 97	1 (0%)

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	254	LEU	6.7
1	A	575	TRP	6.3
1	A	406	PHE	6.3
1	A	536	VAL	6.3
1	A	252	HIS	4.7
1	A	256	ILE	4.6
1	A	253	ASN	4.4
1	A	255	HIS	4.3
1	A	443	ILE	4.1
1	A	286	LEU	3.9
1	A	467	ASN	3.8
1	A	547	PRO	3.7
1	A	289	PHE	3.6
1	A	313	ILE	3.5
1	A	293	VAL	3.5
1	A	295	SER	3.4
1	A	314	ILE	3.3
1	A	409	GLY	3.2
1	A	294	THR	3.1
1	A	316	ILE	3.0
1	A	411	LEU	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	535	ALA	2.9
1	A	440	HIS	2.8
1	A	302	ILE	2.8
1	A	574	ASP	2.7
1	A	453	GLN	2.7
1	A	518	ALA	2.7
1	A	297	GLN	2.6
1	A	534	THR	2.6
1	A	299	ALA	2.6
1	A	468	GLY	2.5
1	A	521	LYS	2.5
1	A	511	HIS	2.5
1	A	287	LYS	2.5
1	A	317	LEU	2.4
1	A	565	LEU	2.3
1	A	506	PHE	2.3
1	A	434	HIS	2.3
1	A	548	GLY	2.2
1	A	290	HIS	2.2
1	A	522	THR	2.0
1	A	304	GLY	2.0
1	A	436	ASP	2.0
1	A	437	GLY	2.0
1	A	524	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	CAS	A	300	9/10	0.93	0.13	77,83,96,98	0
3	DOC	P	6	18/19	0.98	0.07	26,29,32,33	0

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
8	GOL	A	608	6/6	0.73	0.20	45,51,55,60	0
8	GOL	A	607	6/6	0.82	0.16	41,59,65,72	0
7	ACT	A	605	4/4	0.83	0.17	46,51,54,58	4
7	ACT	A	606	4/4	0.83	0.23	51,57,58,63	0
7	ACT	A	604	4/4	0.91	0.11	53,54,54,55	0
5	8DG	A	601	32/32	0.96	0.06	23,28,33,34	0
6	MG	A	602	1/1	0.97	0.05	32,32,32,32	0
6	MG	A	603	1/1	0.99	0.04	25,25,25,25	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.