



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 29, 2026 – 03:58 PM UTC

PDB ID : 6XAY / pdb\_00006xay  
Title : Structure of a fragment of human fibronectin containing the 10th, 11th and 12th type III domains  
Authors : Mou, T.C.; Nepomuceno, P.A.; Sprang, S.R.; Briknarova, K.  
Deposited on : 2020-06-05  
Resolution : 2.48 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

|                                |   |                                                                  |
|--------------------------------|---|------------------------------------------------------------------|
| MolProbity                     | : | <b>FAILED</b>                                                    |
| Mogul                          | : | 2022.3.0, CSD as543be (2022)                                     |
| Xtriage (Phenix)               | : | 2.0                                                              |
| EDS                            | : | 3.0                                                              |
| Percentile statistics          | : | 20250101.v01 (using entries in the PDB archive January 1st 2025) |
| CCP4                           | : | 9.0.010 (Gargrove)                                               |
| Density-Fitness                | : | 1.0.12                                                           |
| Ideal geometry (proteins)      | : | Engh & Huber (2001)                                              |
| Ideal geometry (DNA, RNA)      | : | Parkinson et al. (1996)                                          |
| Validation Pipeline (wwPDB-VP) | : | 2.49                                                             |


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric        | Whole archive<br>(#Entries) | Similar resolution<br>(#Entries, resolution range(Å)) |
|---------------|-----------------------------|-------------------------------------------------------|
| $R_{free}$    | 180053                      | 7589 (2.50-2.46)                                      |
| RSRZ outliers | 180081                      | 7593 (2.50-2.46)                                      |

MolProbity failed to run properly - the sequence quality summary graphics cannot be shown.

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 8315 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fibronectin.

| Mol | Chain | Residues | Atoms |      |     |     |   | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|---------|-------|
| 1   | A     | 272      | Total | C    | N   | O   | S | 0       | 0       | 0     |
|     |       |          | 2042  | 1286 | 336 | 416 | 4 |         |         |       |
| 1   | B     | 269      | Total | C    | N   | O   | S | 0       | 0       | 0     |
|     |       |          | 2022  | 1274 | 332 | 412 | 4 |         |         |       |
| 1   | C     | 271      | Total | C    | N   | O   | S | 0       | 0       | 0     |
|     |       |          | 2036  | 1283 | 335 | 414 | 4 |         |         |       |
| 1   | D     | 272      | Total | C    | N   | O   | S | 0       | 0       | 0     |
|     |       |          | 2042  | 1286 | 336 | 416 | 4 |         |         |       |

There are 20 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment             | Reference  |
|-------|---------|----------|--------|---------------------|------------|
| A     | 1443    | GLY      | -      | expression tag      | UNP P02751 |
| A     | 1444    | SER      | -      | expression tag      | UNP P02751 |
| A     | 1445    | HIS      | -      | expression tag      | UNP P02751 |
| A     | 1446    | MET      | -      | expression tag      | UNP P02751 |
| A     | 1643    | GLY      | VAL    | engineered mutation | UNP P02751 |
| B     | 1443    | GLY      | -      | expression tag      | UNP P02751 |
| B     | 1444    | SER      | -      | expression tag      | UNP P02751 |
| B     | 1445    | HIS      | -      | expression tag      | UNP P02751 |
| B     | 1446    | MET      | -      | expression tag      | UNP P02751 |
| B     | 1643    | GLY      | VAL    | engineered mutation | UNP P02751 |
| C     | 1443    | GLY      | -      | expression tag      | UNP P02751 |
| C     | 1444    | SER      | -      | expression tag      | UNP P02751 |
| C     | 1445    | HIS      | -      | expression tag      | UNP P02751 |
| C     | 1446    | MET      | -      | expression tag      | UNP P02751 |
| C     | 1643    | GLY      | VAL    | engineered mutation | UNP P02751 |
| D     | 1443    | GLY      | -      | expression tag      | UNP P02751 |
| D     | 1444    | SER      | -      | expression tag      | UNP P02751 |
| D     | 1445    | HIS      | -      | expression tag      | UNP P02751 |
| D     | 1446    | MET      | -      | expression tag      | UNP P02751 |
| D     | 1643    | GLY      | VAL    | engineered mutation | UNP P02751 |

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula:  $C_3H_8O_3$ ).



| Mol | Chain | Residues | Atoms |   |   | ZeroOcc | AltConf |
|-----|-------|----------|-------|---|---|---------|---------|
| 2   | A     | 1        | Total | C | O | 0       | 0       |
|     |       |          | 6     | 3 | 3 |         |         |
| 2   | A     | 1        | Total | C | O | 0       | 0       |
|     |       |          | 6     | 3 | 3 |         |         |
| 2   | B     | 1        | Total | C | O | 0       | 0       |
|     |       |          | 6     | 3 | 3 |         |         |
| 2   | B     | 1        | Total | C | O | 0       | 0       |
|     |       |          | 6     | 3 | 3 |         |         |
| 2   | B     | 1        | Total | C | O | 0       | 0       |
|     |       |          | 6     | 3 | 3 |         |         |
| 2   | D     | 1        | Total | C | O | 0       | 0       |
|     |       |          | 6     | 3 | 3 |         |         |

- Molecule 3 is water.

| Mol | Chain | Residues | Atoms |    | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|---------|---------|
| 3   | A     | 44       | Total | O  | 0       | 0       |
|     |       |          | 44    | 44 |         |         |
| 3   | B     | 20       | Total | O  | 0       | 0       |
|     |       |          | 20    | 20 |         |         |
| 3   | C     | 27       | Total | O  | 0       | 0       |
|     |       |          | 27    | 27 |         |         |
| 3   | D     | 46       | Total | O  | 0       | 0       |
|     |       |          | 46    | 46 |         |         |

MolProbity failed to run properly - this section is therefore empty.

### 3 Data and refinement statistics

| Property                                                                | Value                                                       | Source           |
|-------------------------------------------------------------------------|-------------------------------------------------------------|------------------|
| Space group                                                             | C 2 2 2                                                     | Depositor        |
| Cell constants<br>a, b, c, $\alpha$ , $\beta$ , $\gamma$                | 125.08Å 440.37Å 60.47Å<br>90.00° 90.00° 90.00°              | Depositor        |
| Resolution (Å)                                                          | 37.40 – 2.48<br>37.40 – 2.48                                | Depositor<br>EDS |
| % Data completeness<br>(in resolution range)                            | 92.1 (37.40-2.48)<br>92.2 (37.40-2.48)                      | Depositor<br>EDS |
| $R_{merge}$                                                             | (Not available)                                             | Depositor        |
| $R_{sym}$                                                               | (Not available)                                             | Depositor        |
| $\langle I/\sigma(I) \rangle$ <sup>1</sup>                              | 2.84 (at 2.48Å)                                             | Xtriage          |
| Refinement program                                                      | PHENIX 1.16                                                 | Depositor        |
| R, $R_{free}$                                                           | 0.246 , 0.296<br>0.246 , 0.296                              | Depositor<br>DCC |
| $R_{free}$ test set                                                     | 2000 reflections (3.33%)                                    | wwPDB-VP         |
| Wilson B-factor (Å <sup>2</sup> )                                       | 41.8                                                        | Xtriage          |
| Anisotropy                                                              | 0.624                                                       | Xtriage          |
| Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> ) | 0.30 , 59.3                                                 | EDS              |
| L-test for twinning <sup>2</sup>                                        | $\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$ | Xtriage          |
| Estimated twinning fraction                                             | No twinning to report.                                      | Xtriage          |
| $F_o, F_c$ correlation                                                  | 0.93                                                        | EDS              |
| Total number of atoms                                                   | 8315                                                        | wwPDB-VP         |
| Average B, all atoms (Å <sup>2</sup> )                                  | 72.0                                                        | wwPDB-VP         |

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 39.23 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.2757e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 4 Model quality [i](#)

### 4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

### 4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

### 4.3 Torsion angles [i](#)

#### 4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

### 4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 4.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 4.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res  | Link | Bond lengths |      |          | Bond angles |      |          |
|-----|------|-------|------|------|--------------|------|----------|-------------|------|----------|
|     |      |       |      |      | Counts       | RMSZ | # Z  > 2 | Counts      | RMSZ | # Z  > 2 |
| 2   | GOL  | B     | 1803 | -    | 5,5,5        | 0.97 | 0        | 5,5,5       | 1.13 | 1 (20%)  |
| 2   | GOL  | B     | 1802 | -    | 5,5,5        | 1.00 | 0        | 5,5,5       | 1.03 | 0        |
| 2   | GOL  | B     | 1801 | -    | 5,5,5        | 0.90 | 0        | 5,5,5       | 1.12 | 1 (20%)  |
| 2   | GOL  | D     | 1801 | -    | 5,5,5        | 0.85 | 0        | 5,5,5       | 1.12 | 0        |
| 2   | GOL  | A     | 1801 | -    | 5,5,5        | 0.99 | 0        | 5,5,5       | 1.16 | 0        |
| 2   | GOL  | A     | 1802 | -    | 5,5,5        | 1.05 | 1 (20%)  | 5,5,5       | 1.11 | 0        |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res  | Link | Chirals | Torsions | Rings |
|-----|------|-------|------|------|---------|----------|-------|
| 2   | GOL  | B     | 1803 | -    | -       | 2/4/4/4  | -     |
| 2   | GOL  | B     | 1802 | -    | -       | 0/4/4/4  | -     |
| 2   | GOL  | B     | 1801 | -    | -       | 0/4/4/4  | -     |
| 2   | GOL  | D     | 1801 | -    | -       | 0/4/4/4  | -     |
| 2   | GOL  | A     | 1801 | -    | -       | 2/4/4/4  | -     |
| 2   | GOL  | A     | 1802 | -    | -       | 0/4/4/4  | -     |

All (1) bond length outliers are listed below:

| Mol | Chain | Res  | Type | Atoms | Z     | Observed(Å) | Ideal(Å) |
|-----|-------|------|------|-------|-------|-------------|----------|
| 2   | A     | 1802 | GOL  | O2-C2 | -2.06 | 1.37        | 1.43     |

All (2) bond angle outliers are listed below:

| Mol | Chain | Res  | Type | Atoms    | Z     | Observed(°) | Ideal(°) |
|-----|-------|------|------|----------|-------|-------------|----------|
| 2   | B     | 1803 | GOL  | C3-C2-C1 | -2.01 | 104.43      | 111.80   |
| 2   | B     | 1801 | GOL  | C3-C2-C1 | -2.00 | 104.46      | 111.80   |

There are no chirality outliers.

All (4) torsion outliers are listed below:

| Mol | Chain | Res  | Type | Atoms       |
|-----|-------|------|------|-------------|
| 2   | B     | 1803 | GOL  | C1-C2-C3-O3 |

*Continued on next page...*

*Continued from previous page...*

| Mol | Chain | Res  | Type | Atoms       |
|-----|-------|------|------|-------------|
| 2   | B     | 1803 | GOL  | O2-C2-C3-O3 |
| 2   | A     | 1801 | GOL  | O1-C1-C2-C3 |
| 2   | A     | 1801 | GOL  | O1-C1-C2-O2 |

There are no ring outliers.

No monomer is involved in short contacts.

#### 4.7 Other polymers [i](#)

There are no such residues in this entry.

#### 4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 5 Fit of model and data [i](#)

### 5.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

| Mol | Chain | Analysed        | <RSRZ> | #RSRZ>2        | OWAB(Å <sup>2</sup> ) | Q<0.9 |
|-----|-------|-----------------|--------|----------------|-----------------------|-------|
| 1   | A     | 272/280 (97%)   | 0.24   | 12 (4%) 39 35  | 38, 55, 102, 145      | 0     |
| 1   | B     | 269/280 (96%)   | 0.87   | 31 (11%) 9 8   | 48, 81, 113, 156      | 0     |
| 1   | C     | 271/280 (96%)   | 1.04   | 54 (19%) 3 2   | 28, 91, 139, 185      | 0     |
| 1   | D     | 272/280 (97%)   | 0.17   | 11 (4%) 42 38  | 29, 53, 102, 146      | 0     |
| All | All   | 1084/1120 (96%) | 0.58   | 108 (9%) 12 10 | 28, 67, 122, 185      | 0     |

The worst 5 of 108 RSRZ outliers are listed below:

| Mol | Chain | Res  | Type | RSRZ |
|-----|-------|------|------|------|
| 1   | A     | 1674 | PRO  | 5.7  |
| 1   | C     | 1522 | THR  | 5.1  |
| 1   | A     | 1581 | GLY  | 4.7  |
| 1   | C     | 1560 | VAL  | 4.5  |
| 1   | C     | 1470 | ALA  | 4.2  |

### 5.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

| Mol | Type | Chain | Res  | Atoms | RSCC | RSR  | B-factors( $\text{\AA}^2$ ) | Q<0.9 |
|-----|------|-------|------|-------|------|------|-----------------------------|-------|
| 2   | GOL  | B     | 1803 | 6/6   | 0.82 | 0.25 | 83,102,126,131              | 0     |
| 2   | GOL  | B     | 1802 | 6/6   | 0.84 | 0.23 | 85,95,124,142               | 0     |
| 2   | GOL  | A     | 1801 | 6/6   | 0.85 | 0.24 | 44,60,76,84                 | 0     |
| 2   | GOL  | B     | 1801 | 6/6   | 0.89 | 0.16 | 65,69,84,91                 | 6     |
| 2   | GOL  | A     | 1802 | 6/6   | 0.92 | 0.14 | 43,50,62,69                 | 6     |
| 2   | GOL  | D     | 1801 | 6/6   | 0.95 | 0.21 | 43,43,69,84                 | 0     |

## 5.5 Other polymers [i](#)

There are no such residues in this entry.