



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 02:50 PM UTC

PDB ID : 1XBI / pdb_00001xbi
Title : High resolution structure of Methanocaldococcus jannaschii L7AE
Authors : Brown II, B.A.; Suryadi, J.; Lieberman, D.V.; Tran, E.J.; Maxwell, E.S.
Deposited on : 2004-08-30
Resolution : 1.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1756 (1.46-1.46)
Clashscore	190562	1795 (1.46-1.46)
Ramachandran outliers	187476	1776 (1.46-1.46)
Sidechain outliers	187428	1776 (1.46-1.46)
RSRZ outliers	180081	1756 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	120	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	EPE	A	150	-	X	-	-

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 1146 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 50S ribosomal protein L7Ae.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	118	Total	C	N	O	S	0	13	0
			958	617	155	184	2			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	cloning artifact	UNP P54066
A	-2	SER	-	cloning artifact	UNP P54066
A	-1	HIS	-	cloning artifact	UNP P54066

- Molecule 2 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: C₈H₁₈N₂O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			15	8	2	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	173	Total 173	O 173	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 50S ribosomal protein L7Ae



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	43.66Å 48.87Å 51.91Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	27.63 – 1.45 27.63 – 1.45	Depositor EDS
% Data completeness (in resolution range)	95.6 (27.63-1.45) 95.6 (27.63-1.45)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.38 (at 1.40Å)	Xtriage
Refinement program	REFMAC 5.2.0001	Depositor
R, R_{free}	0.132 , 0.193 (Not available) , 0.198	Depositor DCC
R_{free} test set	1087 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	11.6	Xtriage
Anisotropy	0.160	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 43.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	1146	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EPE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	3.16	111/1026 (10.8%)	2.49	82/1381 (5.9%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

All (111) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1	MET	N-CA	-26.16	1.12	1.46
1	A	106	LYS	CE-NZ	13.06	1.88	1.49
1	A	68	LYS	CD-CE	12.29	1.89	1.52
1	A	8	LYS	CA-C	-11.20	1.39	1.52
1	A	88	GLU	CD-OE2	9.74	1.43	1.25
1	A	8	LYS	CD-CE	9.27	1.80	1.52
1	A	22	ALA	N-CA	9.23	1.58	1.46
1	A	101	ASP	CA-C	-9.21	1.42	1.52
1	A	12	GLU	N-CA	-9.12	1.35	1.46
1	A	14	GLN	CA-C	-8.88	1.40	1.52
1	A	50	GLU	CA-C	-8.71	1.40	1.52
1	A	106	LYS	CA-C	8.70	1.64	1.52
1	A	33	GLU	CD-OE2	8.67	1.41	1.25
1	A	9	VAL	N-CA	-8.43	1.39	1.46
1	A	40[A]	ARG	CB-CG	-8.36	1.27	1.52
1	A	40[B]	ARG	CB-CG	-8.36	1.27	1.52
1	A	86	GLY	CA-C	-8.20	1.40	1.51
1	A	11	GLU	CB-CG	-8.10	1.28	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	35	THR	C-O	7.97	1.33	1.24
1	A	40[A]	ARG	NE-CZ	-7.80	1.24	1.33
1	A	40[B]	ARG	NE-CZ	-7.80	1.24	1.33
1	A	83	LYS	CE-NZ	7.79	1.72	1.49
1	A	-1	HIS	CA-CB	7.77	1.69	1.53
1	A	29	LYS	CD-CE	-7.77	1.29	1.52
1	A	88	GLU	C-O	-7.67	1.13	1.24
1	A	-1	HIS	N-CA	7.62	1.60	1.46
1	A	117	GLN	CD-OE1	7.54	1.37	1.23
1	A	22	ALA	CA-C	-7.37	1.42	1.52
1	A	8	LYS	CG-CD	7.22	1.74	1.52
1	A	39[A]	GLU	C-O	7.20	1.32	1.24
1	A	39[B]	GLU	C-O	7.20	1.32	1.24
1	A	21	VAL	CA-CB	-7.03	1.46	1.54
1	A	115	LEU	CA-CB	7.02	1.64	1.53
1	A	100	GLY	CA-C	-6.99	1.42	1.51
1	A	99	GLU	C-O	6.88	1.33	1.23
1	A	39[A]	GLU	CA-CB	6.87	1.64	1.53
1	A	39[B]	GLU	CA-CB	6.87	1.64	1.53
1	A	89	VAL	CA-CB	-6.87	1.44	1.54
1	A	16	GLU	C-O	-6.80	1.16	1.24
1	A	3	VAL	CA-CB	6.77	1.62	1.54
1	A	80	ASP	CG-OD1	-6.66	1.12	1.25
1	A	23	LYS	CA-C	-6.62	1.43	1.52
1	A	110	GLU	CD-OE2	6.58	1.37	1.25
1	A	116	LYS	N-CA	-6.43	1.37	1.46
1	A	100	GLY	C-O	6.36	1.31	1.24
1	A	80	ASP	N-CA	-6.22	1.38	1.46
1	A	60	HIS	CD2-NE2	-6.21	1.31	1.37
1	A	59	ALA	C-O	6.17	1.31	1.24
1	A	73	ALA	CA-CB	6.12	1.61	1.53
1	A	9	VAL	C-N	-6.05	1.26	1.33
1	A	31	ALA	CA-C	-6.01	1.45	1.52
1	A	116	LYS	CB-CG	-6.00	1.34	1.52
1	A	30	GLY	N-CA	-6.00	1.39	1.45
1	A	55[A]	GLU	CA-C	-5.97	1.44	1.52
1	A	55[B]	GLU	CA-C	-5.97	1.44	1.52
1	A	76	ALA	N-CA	-5.91	1.39	1.46
1	A	79	GLN	N-CA	5.90	1.53	1.46
1	A	1	MET	CA-C	-5.86	1.45	1.52
1	A	73	ALA	CA-C	-5.85	1.45	1.53
1	A	117	GLN	CG-CD	-5.85	1.37	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	78	LYS	C-O	-5.77	1.16	1.24
1	A	117	GLN	CD-NE2	5.76	1.45	1.33
1	A	114	VAL	CA-CB	5.73	1.61	1.54
1	A	59	ALA	CA-C	-5.70	1.44	1.52
1	A	111	LYS	N-CA	-5.69	1.39	1.46
1	A	105	LEU	N-CA	5.68	1.53	1.46
1	A	15	LYS	CD-CE	5.67	1.69	1.52
1	A	73	ALA	C-O	5.67	1.29	1.24
1	A	13	ILE	CG1-CD1	-5.64	1.29	1.51
1	A	80	ASP	C-O	-5.63	1.17	1.24
1	A	117	GLN	N-CA	5.62	1.56	1.46
1	A	103[A]	GLU	C-O	-5.62	1.17	1.24
1	A	103[B]	GLU	C-O	-5.62	1.17	1.24
1	A	66	GLU	CG-CD	5.61	1.66	1.52
1	A	101	ASP	CG-OD2	-5.60	1.14	1.25
1	A	16	GLU	N-CA	-5.57	1.39	1.46
1	A	18	LEU	C-O	-5.55	1.17	1.24
1	A	83	LYS	N-CA	5.55	1.53	1.46
1	A	37	ALA	CA-C	-5.55	1.45	1.52
1	A	116	LYS	CE-NZ	5.54	1.66	1.49
1	A	36	LYS	C-N	5.54	1.41	1.34
1	A	28	LYS	CE-NZ	5.54	1.66	1.49
1	A	31	ALA	N-CA	5.54	1.53	1.46
1	A	110	GLU	CA-CB	5.54	1.62	1.53
1	A	38	VAL	CA-CB	-5.46	1.47	1.54
1	A	17	LEU	CA-C	-5.43	1.46	1.52
1	A	38	VAL	C-O	-5.41	1.17	1.24
1	A	1	MET	CB-CG	5.41	1.68	1.52
1	A	116	LYS	CG-CD	5.39	1.68	1.52
1	A	11	GLU	CA-CB	5.36	1.61	1.53
1	A	27	ILE	CA-C	-5.33	1.46	1.52
1	A	9	VAL	CA-C	5.32	1.57	1.53
1	A	14	GLN	CD-OE1	5.26	1.33	1.23
1	A	-1	HIS	CB-CG	5.25	1.57	1.50
1	A	21	VAL	N-CA	-5.25	1.40	1.46
1	A	16	GLU	CD-OE1	5.25	1.35	1.25
1	A	85	ALA	C-O	5.23	1.31	1.24
1	A	33	GLU	CG-CD	5.21	1.65	1.52
1	A	32[A]	ASN	CG-ND2	5.18	1.44	1.33
1	A	32[B]	ASN	CG-ND2	5.18	1.44	1.33
1	A	35	THR	CA-C	-5.16	1.46	1.52
1	A	11	GLU	CD-OE1	5.15	1.35	1.25

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	13	ILE	N-CA	5.12	1.53	1.46
1	A	25[A]	GLN	CA-C	-5.10	1.45	1.52
1	A	25[B]	GLN	CA-C	-5.10	1.45	1.52
1	A	2	ALA	N-CA	5.07	1.52	1.45
1	A	57[A]	VAL	CA-CB	-5.07	1.48	1.54
1	A	57[B]	VAL	CA-CB	-5.07	1.48	1.54
1	A	112	VAL	N-CA	-5.06	1.40	1.46
1	A	5	VAL	C-N	-5.03	1.25	1.33
1	A	108	LEU	CA-C	5.02	1.59	1.52

All (82) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	68	LYS	CG-CD-CE	11.24	137.15	111.30
1	A	40[A]	ARG	NE-CZ-NH2	-10.42	109.82	119.20
1	A	40[B]	ARG	NE-CZ-NH2	-10.42	109.82	119.20
1	A	-1	HIS	CA-C-N	9.09	138.90	121.54
1	A	-1	HIS	C-N-CA	9.09	138.90	121.54
1	A	100	GLY	O-C-N	-8.23	112.88	122.34
1	A	30	GLY	CA-C-O	-8.12	116.63	122.23
1	A	36	LYS	O-C-N	-7.89	113.94	122.07
1	A	8	LYS	CD-CE-NZ	-7.76	87.08	111.90
1	A	80	ASP	OD1-CG-OD2	7.76	141.51	122.90
1	A	111	LYS	O-C-N	-7.73	114.05	122.09
1	A	19	ASP	O-C-N	-7.72	112.78	122.27
1	A	-1	HIS	CB-CA-C	7.57	124.49	110.10
1	A	101	ASP	CA-C-N	7.43	130.10	120.44
1	A	101	ASP	C-N-CA	7.43	130.10	120.44
1	A	23	LYS	O-C-N	-7.26	113.27	122.34
1	A	2	ALA	CA-C-O	7.17	129.76	121.87
1	A	24	ALA	CA-C-O	-7.16	113.96	121.55
1	A	-1	HIS	CA-CB-CG	7.11	120.91	113.80
1	A	15	LYS	CD-CE-NZ	-7.08	89.26	111.90
1	A	99	GLU	OE1-CD-OE2	-7.05	105.98	122.90
1	A	53[A]	LYS	O-C-N	-7.04	113.48	121.79
1	A	53[B]	LYS	O-C-N	-7.04	113.48	121.79
1	A	20	ALA	O-C-N	-7.04	114.82	122.07
1	A	99	GLU	CG-CD-OE2	7.00	134.49	118.40
1	A	2	ALA	O-C-N	-6.88	114.70	122.75
1	A	115	LEU	CB-CA-C	-6.54	99.94	110.79
1	A	83	LYS	O-C-N	-6.53	114.70	122.15
1	A	35	THR	O-C-N	-6.37	115.37	122.12

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	76	ALA	N-CA-C	6.34	118.19	111.28
1	A	5	VAL	CA-C-O	-6.31	113.66	121.04
1	A	79	GLN	CB-CA-C	6.25	120.70	110.88
1	A	2	ALA	CA-C-N	6.25	129.34	120.53
1	A	2	ALA	C-N-CA	6.25	129.34	120.53
1	A	3	VAL	N-CA-CB	6.21	118.24	110.47
1	A	36	LYS	CD-CE-NZ	-6.17	92.16	111.90
1	A	78	LYS	N-CA-C	6.10	118.89	111.82
1	A	114	VAL	CA-C-O	6.08	127.27	120.95
1	A	32[A]	ASN	CA-C-N	6.07	128.41	120.28
1	A	32[A]	ASN	C-N-CA	6.07	128.41	120.28
1	A	32[B]	ASN	CA-C-N	6.07	128.41	120.28
1	A	32[B]	ASN	C-N-CA	6.07	128.41	120.28
1	A	104[A]	GLU	CA-C-O	-6.06	114.45	120.82
1	A	104[B]	GLU	CA-C-O	-6.06	114.45	120.82
1	A	111	LYS	CD-CE-NZ	-5.89	93.06	111.90
1	A	20	ALA	CA-C-N	5.89	128.19	120.60
1	A	20	ALA	C-N-CA	5.89	128.19	120.60
1	A	114	VAL	CB-CA-C	-5.85	104.24	112.14
1	A	73	ALA	N-CA-C	5.84	115.94	108.24
1	A	40[A]	ARG	CD-NE-CZ	5.81	132.54	124.40
1	A	40[B]	ARG	CD-NE-CZ	5.81	132.54	124.40
1	A	114	VAL	O-C-N	-5.80	116.24	121.87
1	A	33	GLU	O-C-N	-5.79	115.98	122.12
1	A	111	LYS	N-CA-CB	5.75	118.52	109.94
1	A	21	VAL	N-CA-CB	5.72	116.86	110.51
1	A	113	ASN	N-CA-C	-5.55	105.13	111.07
1	A	80	ASP	CA-CB-CG	5.54	118.14	112.60
1	A	1	MET	CB-CA-C	5.54	121.45	110.42
1	A	53[A]	LYS	CA-C-O	5.50	124.73	119.72
1	A	53[B]	LYS	CA-C-O	5.50	124.73	119.72
1	A	19	ASP	CA-C-N	5.44	127.51	120.44
1	A	19	ASP	C-N-CA	5.44	127.51	120.44
1	A	102	ALA	O-C-N	-5.42	116.48	122.07
1	A	108	LEU	CA-C-O	-5.38	114.27	120.24
1	A	32[A]	ASN	O-C-N	-5.34	116.33	122.09
1	A	32[B]	ASN	O-C-N	-5.34	116.33	122.09
1	A	83	LYS	CG-CD-CE	5.31	123.51	111.30
1	A	14	GLN	CG-CD-OE1	-5.31	110.19	120.80
1	A	14	GLN	O-C-N	-5.29	116.52	122.12
1	A	68	LYS	CD-CE-NZ	5.24	128.68	111.90
1	A	3	VAL	CA-CB-CG2	-5.22	101.52	110.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	101	ASP	O-C-N	-5.22	116.35	123.24
1	A	16	GLU	CG-CD-OE1	-5.21	106.42	118.40
1	A	68	LYS	O-C-N	-5.20	116.09	122.34
1	A	-1	HIS	N-CA-CB	-5.15	101.75	110.50
1	A	88	GLU	CB-CA-C	-5.15	102.02	110.72
1	A	80	ASP	CA-C-O	5.08	125.93	120.55
1	A	100	GLY	CA-C-N	5.07	129.41	121.76
1	A	100	GLY	C-N-CA	5.07	129.41	121.76
1	A	36	LYS	CA-C-O	5.05	126.12	120.82
1	A	110	GLU	N-CA-CB	-5.01	102.75	110.16
1	A	84	ALA	O-C-N	-5.01	116.81	122.12

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	106	LYS	Mainchain
1	A	16	GLU	Sidechain
1	A	88	GLU	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	958	0	1012	38	2
2	A	15	0	17	0	0
3	A	173	0	0	15	4
All	All	1146	0	1029	38	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:103[A]:GLU:CD	1:A:103[A]:GLU:CG	1.77	1.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:8:LYS:CD	1:A:8:LYS:CE	1.80	1.52
1:A:68:LYS:CD	1:A:68:LYS:CE	1.89	1.51
1:A:83:LYS:CE	1:A:83:LYS:NZ	1.72	1.47
1:A:106:LYS:CE	1:A:106:LYS:NZ	1.88	1.36
1:A:104[B]:GLU:OE2	3:A:255:HOH:O	1.69	1.09
1:A:70[B]:ILE:HD11	3:A:305:HOH:O	1.64	0.97
1:A:8:LYS:CD	1:A:8:LYS:NZ	2.34	0.90
1:A:70[B]:ILE:HD13	3:A:224:HOH:O	1.71	0.89
1:A:54:PRO:O	1:A:57[B]:VAL:HG12	1.74	0.88
1:A:70[B]:ILE:CD1	3:A:305:HOH:O	2.23	0.84
1:A:70[B]:ILE:CD1	3:A:224:HOH:O	2.25	0.83
1:A:32[B]:ASN:ND2	3:A:274:HOH:O	2.14	0.78
1:A:83:LYS:NZ	1:A:83:LYS:CD	2.48	0.77
1:A:52:VAL:HG11	1:A:57[B]:VAL:HG13	1.72	0.70
1:A:103[A]:GLU:CD	1:A:103[A]:GLU:CB	2.63	0.69
1:A:52:VAL:CG1	1:A:57[B]:VAL:CG1	2.75	0.65
1:A:52:VAL:CG1	1:A:57[B]:VAL:HG13	2.29	0.63
1:A:29:LYS:NZ	3:A:278:HOH:O	2.26	0.61
1:A:104[B]:GLU:CD	3:A:333:HOH:O	2.46	0.59
1:A:103[B]:GLU:OE2	3:A:263:HOH:O	2.16	0.57
1:A:2:ALA:HB1	1:A:55[A]:GLU:OE2	2.04	0.57
1:A:41:GLY:HA2	3:A:305:HOH:O	2.05	0.56
1:A:-1:HIS:HB3	1:A:1:MET:SD	2.46	0.56
1:A:8:LYS:NZ	1:A:8:LYS:HD3	2.19	0.55
1:A:104[B]:GLU:CD	3:A:255:HOH:O	2.34	0.54
1:A:8:LYS:CD	1:A:8:LYS:HZ2	2.17	0.54
1:A:1:MET:HE3	1:A:1:MET:HA	1.91	0.52
1:A:6[B]:LYS:NZ	1:A:66:GLU:OE1	2.44	0.50
1:A:1:MET:SD	1:A:1:MET:N	2.85	0.48
1:A:14:GLN:NE2	1:A:76:ALA:H	2.11	0.48
1:A:103[B]:GLU:CD	3:A:263:HOH:O	2.59	0.45
1:A:52:VAL:HG12	1:A:57[B]:VAL:CG1	2.46	0.44
1:A:70[B]:ILE:HD11	3:A:224:HOH:O	2.05	0.44
1:A:48:ILE:O	1:A:74:TYR:HA	2.19	0.43
1:A:56[B]:GLU:HG2	3:A:249:HOH:O	2.18	0.42
1:A:32[A]:ASN:HD21	1:A:36:LYS:HE3	1.86	0.40
1:A:14:GLN:NE2	3:A:220:HOH:O	2.53	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:261:HOH:O	3:A:318:HOH:O[4_466]	1.49	0.71
1:A:55[A]:GLU:OE2	3:A:215:HOH:O[2_575]	1.56	0.64
1:A:39[B]:GLU:OE1	3:A:280:HOH:O[2_565]	1.64	0.56
3:A:229:HOH:O	3:A:297:HOH:O[4_566]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	129/120 (108%)	128 (99%)	0	1 (1%)	16 3

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1	MET

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	108/96 (112%)	104 (96%)	4 (4%)	30 4

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	11	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	68	LYS
1	A	117	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	14	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EPE	A	150	-	15,15,15	2.46	7 (46%)	19,20,20	3.63	10 (52%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EPE	A	150	-	-	5/9/19/19	0/1/1/1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	150	EPE	C9-N1	5.46	1.59	1.47
2	A	150	EPE	C3-C2	3.63	1.64	1.51
2	A	150	EPE	O2S-S	2.94	1.53	1.45
2	A	150	EPE	C3-N4	2.88	1.54	1.46
2	A	150	EPE	O3S-S	2.81	1.57	1.47
2	A	150	EPE	C9-C10	-2.56	1.45	1.52
2	A	150	EPE	C6-N1	2.18	1.52	1.46

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	150	EPE	O1S-S-C10	-7.09	96.01	106.73
2	A	150	EPE	C3-C2-N1	-6.23	98.08	110.65
2	A	150	EPE	O2S-S-C10	5.82	115.53	106.73
2	A	150	EPE	C9-N1-C2	-4.84	98.34	111.24
2	A	150	EPE	C5-N4-C3	4.55	118.64	108.84
2	A	150	EPE	C7-N4-C3	-4.38	99.57	111.24
2	A	150	EPE	C7-N4-C5	4.29	122.68	111.24
2	A	150	EPE	O3S-S-C10	-4.12	97.95	106.00
2	A	150	EPE	O3S-S-O1S	3.30	119.66	111.40
2	A	150	EPE	C2-C3-N4	-2.50	105.60	110.65

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	150	EPE	C8-C7-N4-C5
2	A	150	EPE	C10-C9-N1-C2
2	A	150	EPE	C10-C9-N1-C6
2	A	150	EPE	C9-C10-S-O1S
2	A	150	EPE	C9-C10-S-O3S

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	118/120 (98%)	-0.49	2 (1%) 69 72	5, 12, 28, 62	13 (11%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	-1	HIS	3.8
1	A	6[A]	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	EPE	A	150	15/15	0.94	0.09	18,23,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.