



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 02:40 PM UTC

PDB ID : 2XCN / pdb_00002xcn
Title : Crystal structure of HCV NS3 protease with a boronate inhibitor
Authors : Li, X.; Zhang, Y.-K.; Liu, Y.; Ding, C.Z.; Li, Q.; Zhou, Y.; Plattner, J.J.; Baker, S.J.; Qian, X.; Fan, D.; Liao, L.; Ni, Z.-J.; White, G.V.; Mordaunt, J.E.; Lazarides, L.X.; Slater, M.J.; Jarvest, R.L.; Thommes, P.; Ellis, M.; Edge, C.M.; Hubbard, J.A.; Nassau, P.; McDowell, B.; Skarzynski, T.J.; Rowland, P.; Somers, D.O.; Kazmierski, W.M.; Grimes, R.M.; Wright, L.L.; Smith, G.K.; Zou, W.; Wright, J.; Pennicott, L.E.
Deposited on : 2010-04-23
Resolution : 3.02 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.02 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3131 (3.04-3.00)
Clashscore	190562	3444 (3.04-3.00)
Ramachandran outliers	187476	3319 (3.04-3.00)
Sidechain outliers	187428	3322 (3.04-3.00)
RSRZ outliers	180081	3130 (3.04-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	198	
1	B	198	
2	C	23	

Continued on next page...

Ideal geometry (proteins) : Engh & Huber (2001)
 Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
 Validation Pipeline (wwPDB-VP) : 2.49

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	23	 A horizontal bar chart showing the quality of chain D. The bar is divided into three segments: a green segment representing 52%, a yellow segment representing 17%, and a grey segment representing 30%. The percentages are labeled below the corresponding segments.

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 2870 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NS3 PROTEASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	180	Total	C	N	O	S	0	0	0
			1335	833	243	250	9			
1	B	153	Total	C	N	O	S	0	0	0
			1136	710	206	212	8			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-9	ALA	-	expression tag	UNP Q91RS4
A	-8	SER	-	expression tag	UNP Q91RS4
A	-7	MET	-	expression tag	UNP Q91RS4
A	-6	THR	-	expression tag	UNP Q91RS4
A	-5	GLY	-	expression tag	UNP Q91RS4
A	-4	GLY	-	expression tag	UNP Q91RS4
A	-3	GLN	-	expression tag	UNP Q91RS4
A	-2	GLN	-	expression tag	UNP Q91RS4
A	-1	MET	-	expression tag	UNP Q91RS4
A	0	GLY	-	expression tag	UNP Q91RS4
A	181	GLY	-	expression tag	UNP Q91RS4
A	182	SER	-	expression tag	UNP Q91RS4
A	183	HIS	-	expression tag	UNP Q91RS4
A	184	HIS	-	expression tag	UNP Q91RS4
A	185	HIS	-	expression tag	UNP Q91RS4
A	186	HIS	-	expression tag	UNP Q91RS4
A	187	HIS	-	expression tag	UNP Q91RS4
A	188	HIS	-	expression tag	UNP Q91RS4
A	40	THR	ALA	conflict	UNP Q91RS4
A	91	SER	ALA	conflict	UNP Q91RS4
B	-9	ALA	-	expression tag	UNP Q91RS4
B	-8	SER	-	expression tag	UNP Q91RS4
B	-7	MET	-	expression tag	UNP Q91RS4
B	-6	THR	-	expression tag	UNP Q91RS4
B	-5	GLY	-	expression tag	UNP Q91RS4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-4	GLY	-	expression tag	UNP Q91RS4
B	-3	GLN	-	expression tag	UNP Q91RS4
B	-2	GLN	-	expression tag	UNP Q91RS4
B	-1	MET	-	expression tag	UNP Q91RS4
B	0	GLY	-	expression tag	UNP Q91RS4
B	181	GLY	-	expression tag	UNP Q91RS4
B	182	SER	-	expression tag	UNP Q91RS4
B	183	HIS	-	expression tag	UNP Q91RS4
B	184	HIS	-	expression tag	UNP Q91RS4
B	185	HIS	-	expression tag	UNP Q91RS4
B	186	HIS	-	expression tag	UNP Q91RS4
B	187	HIS	-	expression tag	UNP Q91RS4
B	188	HIS	-	expression tag	UNP Q91RS4
B	40	THR	ALA	conflict	UNP Q91RS4
B	91	SER	ALA	conflict	UNP Q91RS4


- Molecule 2 is a protein called NS4A.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	21	Total	C	N	O	0	0	0
			149	99	27	23			
2	D	16	Total	C	N	O	0	0	0
			108	70	20	18			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	19	LYS	-	expression tag	UNP C9WU77
C	20	LYS	-	expression tag	UNP C9WU77
C	40	LYS	-	expression tag	UNP C9WU77
C	41	LYS	-	expression tag	UNP C9WU77
D	19	LYS	-	expression tag	UNP C9WU77
D	20	LYS	-	expression tag	UNP C9WU77
D	40	LYS	-	expression tag	UNP C9WU77
D	41	LYS	-	expression tag	UNP C9WU77

- Molecule 3 is N-[(CYCLOPENTYLOXY)CARBONYL]-3-METHYL-L-VALYL-(4R)-N-[(1R)-3-HYDROXY-1-[HYDROXY(OXIDO)BORANYL]PROPYL]-4-(ISOQUINOLIN-1-YLOXY)-L-PROLINAMIDE (CCD ID: C8D) (formula: C₂₉H₄₀BN₄O₈).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	B	C	N	O	0	0
			42	1	29	4	8		
3	B	1	Total	B	C	N	O	0	0
			42	1	29	4	8		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		
4	C	1	Total	Mg	0	0
			1	1		

- Molecule 5 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Zn	0	0
			1	1		
5	B	1	Total	Zn	0	0
			1	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	24	Total	O	0	0
			24	24		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	21	Total 21	O 21	0	0
6	C	5	Total 5	O 5	0	0
6	D	4	Total 4	O 4	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: NS3 PROTEASE



• Molecule 1: NS3 PROTEASE



• Molecule 2: NS4A



• Molecule 2: NS4A



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	231.37Å 231.37Å 75.67Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	70.71 – 3.02 70.71 – 3.02	Depositor EDS
% Data completeness (in resolution range)	100.0 (70.71-3.02) 99.4 (70.71-3.02)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.80 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.184 , 0.239 0.175 , 0.232	Depositor DCC
R_{free} test set	763 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	69.6	Xtriage
Anisotropy	0.381	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 55.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2870	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: C8D, ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.70	0/1361	0.97	1/1856 (0.1%)
1	B	0.67	0/1160	1.06	6/1583 (0.4%)
2	C	0.89	0/150	1.18	1/201 (0.5%)
2	D	0.86	0/108	1.27	0/145
All	All	0.70	0/2779	1.03	8/3785 (0.2%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	130	ARG	N-CA-C	8.53	115.67	108.07
1	B	159	CYS	N-CA-C	7.97	121.46	108.55
1	B	66	SER	CA-C-N	-7.52	112.11	120.45
1	B	66	SER	C-N-CA	-7.52	112.11	120.45
1	B	163	VAL	N-CA-C	6.54	117.30	107.75
1	A	159	CYS	N-CA-C	5.81	117.62	108.96
1	B	74	MET	N-CA-C	-5.41	106.81	113.41
2	C	24	VAL	N-CA-C	5.26	115.69	108.12

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1335	0	1351	28	0
1	B	1136	0	1139	20	0
2	C	149	0	180	1	0
2	D	108	0	125	2	0
3	A	42	0	40	4	0
3	B	42	0	40	1	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	24	0	0	1	0
6	B	21	0	0	5	0
6	C	5	0	0	0	0
6	D	4	0	0	0	0
All	All	2870	0	2875	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:108:THR:HG22	1:A:110:HIS:H	1.35	0.92
1:B:62:ARG:HG3	1:B:62:ARG:HH11	1.38	0.87
1:B:159:CYS:HB3	1:B:164:ALA:HA	1.66	0.76
1:B:155:ARG:NH1	1:B:168:ASP:OD2	2.20	0.75
1:A:37:SER:OG	1:A:42:THR:HB	1.92	0.69
1:A:108:THR:HG22	1:A:110:HIS:N	2.07	0.67
1:B:145:CYS:HB2	1:B:146:PRO:HD2	1.77	0.64
1:A:40:THR:HB	1:A:41:GLN:HG3	1.83	0.60
1:A:155:ARG:HD3	3:A:1181:C8D:C26	2.31	0.60
1:B:62:ARG:HG3	1:B:62:ARG:NH1	2.13	0.60
1:A:159:CYS:HB2	1:A:163:VAL:O	2.01	0.59
1:A:139:SER:CB	3:A:1181:C8D:H192	2.37	0.55
1:A:11:ARG:HG2	6:A:2004:HOH:O	2.06	0.55
1:B:134:TYR:C	1:B:135:LEU:HD23	2.32	0.54
1:B:130:ARG:HB2	1:B:131:PRO:HD2	1.89	0.54
1:A:11:ARG:NH2	1:A:25:ASP:OD1	2.42	0.52
1:A:80:GLN:HG3	1:A:174:ASN:HD22	1.75	0.52
1:A:96:PRO:HB3	1:A:172:VAL:HG11	1.92	0.52
1:A:105:TYR:CD1	1:B:147:ALA:HB2	2.46	0.51
1:A:1:ALA:HA	1:A:2:PRO:C	2.36	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:153:LEU:O	1:B:169:PHE:HB2	2.11	0.51
1:A:115:PRO:HG3	6:B:2021:HOH:O	2.10	0.50
1:B:103:ASP:O	1:B:146:PRO:HD3	2.11	0.49
1:A:105:TYR:CE1	1:B:147:ALA:HB2	2.48	0.48
1:B:73:GLN:HG3	6:B:2010:HOH:O	2.13	0.48
1:B:132:ILE:HG12	1:B:164:ALA:HB2	1.96	0.47
1:A:63:THR:HG22	1:A:71:VAL:C	2.39	0.47
1:A:127:LEU:HD12	1:B:99:CYS:HA	1.95	0.47
2:D:34:LYS:HA	2:D:35:PRO:HD3	1.74	0.46
1:B:135:LEU:HD23	1:B:135:LEU:N	2.31	0.46
1:A:14:LEU:O	1:A:18:ILE:HD12	2.16	0.46
1:A:118:ARG:NH2	1:A:121:ASP:O	2.47	0.46
1:A:103:ASP:O	1:A:146:PRO:HD3	2.16	0.46
1:A:59:ALA:O	1:A:62:ARG:HD2	2.16	0.45
1:A:108:THR:CG2	1:A:110:HIS:H	2.19	0.45
1:A:172:VAL:O	1:A:175:LEU:HB2	2.17	0.45
3:A:1181:C8D:H5	3:A:1181:C8D:H132	1.68	0.45
3:B:1181:C8D:H14	6:B:2020:HOH:O	2.18	0.43
1:A:6:TYR:HB2	2:D:34:LYS:HD2	2.00	0.43
1:A:25:ASP:OD1	1:A:25:ASP:C	2.62	0.42
1:A:159:CYS:HB2	1:A:163:VAL:C	2.44	0.42
1:B:126:LEU:N	1:B:126:LEU:HD23	2.35	0.42
1:B:155:ARG:NH2	6:B:2019:HOH:O	2.52	0.42
1:B:115:PRO:HD2	6:B:2017:HOH:O	2.20	0.41
1:B:109:ARG:HD2	1:B:137:GLY:HA3	2.03	0.41
1:B:130:ARG:CB	1:B:131:PRO:HD2	2.51	0.41
1:A:139:SER:HB3	3:A:1181:C8D:H192	2.03	0.41
1:A:33:VAL:HB	2:C:29:ILE:HB	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	178/198 (90%)	172 (97%)	6 (3%)	0	100	100
1	B	151/198 (76%)	142 (94%)	9 (6%)	0	100	100
2	C	19/23 (83%)	19 (100%)	0	0	100	100
2	D	14/23 (61%)	14 (100%)	0	0	100	100
All	All	362/442 (82%)	347 (96%)	15 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	146/159 (92%)	125 (86%)	21 (14%)	3	14
1	B	125/159 (79%)	105 (84%)	20 (16%)	2	12
2	C	17/19 (90%)	13 (76%)	4 (24%)	1	4
2	D	12/19 (63%)	10 (83%)	2 (17%)	2	11
All	All	300/356 (84%)	253 (84%)	47 (16%)	2	12

All (47) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	LEU
1	A	21	LEU
1	A	28	GLN
1	A	37	SER
1	A	40	THR
1	A	41	GLN
1	A	42	THR
1	A	63	THR
1	A	66	SER
1	A	72	ILE
1	A	83	VAL
1	A	92	ARG
1	A	95	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	108	THR
1	A	123	ARG
1	A	128	SER
1	A	132	ILE
1	A	139	SER
1	A	160	THR
1	A	172	VAL
1	A	179	MET
1	B	29	VAL
1	B	40	THR
1	B	41	GLN
1	B	61	THR
1	B	62	ARG
1	B	78	VAL
1	B	83	VAL
1	B	95	THR
1	B	106	LEU
1	B	122	SER
1	B	123	ARG
1	B	126	LEU
1	B	127	LEU
1	B	128	SER
1	B	132	ILE
1	B	135	LEU
1	B	139	SER
1	B	143	LEU
1	B	161	ARG
1	B	172	VAL
2	C	22	SER
2	C	28	ARG
2	C	34	LYS
2	C	37	ILE
2	D	22	SER
2	D	24	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	49	ASN
1	A	174	ASN
1	B	41	GLN
1	B	49	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	174	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	C8D	B	1181	1	40,45,45	1.25	6 (15%)	52,64,64	1.78	10 (19%)
3	C8D	A	1181	1	40,45,45	1.25	3 (7%)	52,64,64	1.69	10 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	C8D	B	1181	1	-	2/36/60/60	0/4/4/4
3	C8D	A	1181	1	-	8/36/60/60	0/4/4/4

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1181	C8D	C21-N4	3.83	1.37	1.31
3	A	1181	C8D	C23-C22	3.26	1.41	1.36
3	B	1181	C8D	C21-N4	3.23	1.36	1.31
3	B	1181	C8D	C23-C22	2.92	1.40	1.36
3	A	1181	C8D	C5-C12	2.77	1.57	1.53
3	B	1181	C8D	C27-C28	2.64	1.42	1.36
3	B	1181	C8D	C26-C25	2.32	1.41	1.36
3	B	1181	C8D	C5-C12	2.08	1.56	1.53
3	B	1181	C8D	C27-C26	2.08	1.42	1.38

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1181	C8D	C29-C21-N4	-6.17	117.67	124.34
3	A	1181	C8D	C7-O2-C6	4.78	123.34	116.52
3	A	1181	C8D	C29-C21-N4	-4.33	119.66	124.34
3	B	1181	C8D	C15-C14-C13	-4.31	100.09	103.69
3	A	1181	C8D	C1-C2-C5	3.59	117.06	109.71
3	B	1181	C8D	O2-C6-N1	3.48	114.70	110.26
3	A	1181	C8D	C5-C12-N2	-3.25	114.58	118.46
3	B	1181	C8D	C7-O2-C6	3.14	121.01	116.52
3	A	1181	C8D	C2-C5-C12	3.05	117.29	113.23
3	A	1181	C8D	O2-C6-N1	3.00	114.08	110.26
3	B	1181	C8D	C15-C16-C17	-2.99	105.63	111.36
3	B	1181	C8D	C12-C5-N1	-2.68	104.16	107.40
3	B	1181	C8D	O2-C6-O1	-2.61	120.70	124.55
3	A	1181	C8D	C15-C14-C13	-2.51	101.59	103.69
3	B	1181	C8D	C2-C5-N1	2.46	115.77	111.94
3	B	1181	C8D	C2-C5-C12	-2.46	109.95	113.23
3	A	1181	C8D	C15-C16-C17	-2.40	106.77	111.36
3	B	1181	C8D	C22-N4-C21	2.07	122.53	116.28
3	A	1181	C8D	C28-C29-C21	-2.05	120.82	123.45
3	A	1181	C8D	O3-C12-C5	2.03	123.06	119.91

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1181	C8D	C11-C7-O2-C6
3	A	1181	C8D	C1-C2-C5-N1
3	A	1181	C8D	C4-C2-C5-N1

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	A	1181	C8D	C1-C2-C5-C12
3	A	1181	C8D	C3-C2-C5-C12
3	A	1181	C8D	C4-C2-C5-C12
3	B	1181	C8D	C11-C7-O2-C6
3	B	1181	C8D	C18-C19-C20-O5
3	A	1181	C8D	C3-C2-C5-N1
3	A	1181	C8D	N3-C18-C19-C20


There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1181	C8D	1	0
3	A	1181	C8D	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	180/198 (90%)	-0.66	0 100 100	36, 53, 71, 80	0
1	B	153/198 (77%)	-0.32	0 100 100	40, 63, 88, 102	0
2	C	21/23 (91%)	-0.63	0 100 100	38, 48, 63, 79	0
2	D	16/23 (69%)	-0.68	0 100 100	39, 48, 69, 74	0
All	All	370/442 (83%)	-0.52	0 100 100	36, 58, 81, 102	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	C8D	B	1181	42/42	0.94	0.12	56,70,78,80	0
3	C8D	A	1181	42/42	0.95	0.09	49,64,69,71	0
4	MG	A	1182	1/1	0.96	0.11	56,56,56,56	0
4	MG	C	1041	1/1	0.98	0.04	52,52,52,52	0

Continued on next page...

Continued from previous page...



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	ZN	A	1183	1/1	0.99	0.02	54,54,54,54	0
5	ZN	B	1182	1/1	1.00	0.02	54,54,54,54	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around C8D A 1181:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.