



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 10, 2026 – 06:32 AM UTC

PDB ID : 1XDM / pdb_00001xdm
Title : Structure of human aldolase B associated with hereditary fructose intolerance (A149P), at 291K
Authors : Malay, A.D.; Allen, K.N.; Tolan, D.R.
Deposited on : 2004-09-07
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	365	
1	B	365	
1	C	365	
1	D	365	
1	W	365	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	X	365	% 24% 50% 8% • 17%
1	Y	365	% 30% 46% 7% • 17%
1	Z	365	% 17% 54% 9% • 19%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 18550 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fructose-bisphosphate aldolase B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	303	Total	C	N	O	S	0	0	0
			2328	1465	409	442	12			
1	B	304	Total	C	N	O	S	0	0	0
			2343	1475	412	444	12			
1	C	299	Total	C	N	O	S	0	0	0
			2305	1454	404	435	12			
1	D	296	Total	C	N	O	S	0	0	0
			2278	1433	400	433	12			
1	W	297	Total	C	N	O	S	0	0	0
			2286	1441	403	430	12			
1	X	302	Total	C	N	O	S	0	0	0
			2321	1463	407	438	13			
1	Y	304	Total	C	N	O	S	0	0	0
			2343	1475	412	444	12			
1	Z	295	Total	C	N	O	S	0	0	0
			2271	1428	399	432	12			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	cloning artifact	UNP P05062
A	0	SER	-	cloning artifact	UNP P05062
A	149	PRO	ALA	engineered mutation	UNP P05062
B	-1	GLY	-	cloning artifact	UNP P05062
B	0	SER	-	cloning artifact	UNP P05062
B	149	PRO	ALA	engineered mutation	UNP P05062
C	-1	GLY	-	cloning artifact	UNP P05062
C	0	SER	-	cloning artifact	UNP P05062
C	149	PRO	ALA	engineered mutation	UNP P05062
D	-1	GLY	-	cloning artifact	UNP P05062
D	0	SER	-	cloning artifact	UNP P05062
D	149	PRO	ALA	engineered mutation	UNP P05062
W	-1	GLY	-	cloning artifact	UNP P05062

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
W	0	SER	-	cloning artifact	UNP P05062
W	149	PRO	ALA	engineered mutation	UNP P05062
X	-1	GLY	-	cloning artifact	UNP P05062
X	0	SER	-	cloning artifact	UNP P05062
X	149	PRO	ALA	engineered mutation	UNP P05062
Y	-1	GLY	-	cloning artifact	UNP P05062
Y	0	SER	-	cloning artifact	UNP P05062
Y	149	PRO	ALA	engineered mutation	UNP P05062
Z	-1	GLY	-	cloning artifact	UNP P05062
Z	0	SER	-	cloning artifact	UNP P05062
Z	149	PRO	ALA	engineered mutation	UNP P05062

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	W	1	Total	O	S	0	0
			5	4	1		
2	X	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	Y	1	Total	O	S	0	0
			5	4	1		
2	Z	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total	O	0	0
			4	4		
3	B	5	Total	O	0	0
			5	5		
3	C	3	Total	O	0	0
			3	3		
3	D	4	Total	O	0	0
			4	4		
3	W	5	Total	O	0	0
			5	5		
3	X	3	Total	O	0	0
			3	3		
3	Y	6	Total	O	0	0
			6	6		
3	Z	5	Total	O	0	0
			5	5		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Fructose-bisphosphate aldolase B









• Molecule 1: Fructose-bisphosphate aldolase B




• Molecule 1: Fructose-bisphosphate aldolase B







• Molecule 1: Fructose-bisphosphate aldolase B



• Molecule 1: Fructose-bisphosphate aldolase B



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	153.42Å 153.47Å 185.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	79.40 – 3.00 79.40 – 3.00	Depositor EDS
% Data completeness (in resolution range)	89.8 (79.40-3.00) 94.9 (79.40-3.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.79 (at 3.01Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.271 , 0.319 0.264 , 0.310	Depositor DCC
R_{free} test set	7921 reflections (9.48%)	wwPDB-VP
Wilson B-factor (Å ²)	57.9	Xtriage
Anisotropy	0.555	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 79.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.41$, $\langle L^2 \rangle = 0.24$	Xtriage
Estimated twinning fraction	0.448 for k,h,-l	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	18550	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.48	0/2363	1.04	11/3187 (0.3%)
1	B	0.54	0/2378	1.11	24/3207 (0.7%)
1	C	0.50	0/2339	1.05	14/3154 (0.4%)
1	D	0.49	0/2312	1.09	15/3117 (0.5%)
1	W	0.48	0/2321	1.08	15/3130 (0.5%)
1	X	0.56	1/2355 (0.0%)	1.08	17/3176 (0.5%)
1	Y	0.54	0/2378	1.10	23/3207 (0.7%)
1	Z	0.47	0/2304	1.05	11/3105 (0.4%)
All	All	0.51	1/18750 (0.0%)	1.08	130/25283 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	X	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	X	191	ILE	CA-C	8.85	1.62	1.52

The worst 5 of 130 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	X	191	ILE	N-CA-C	12.95	136.86	108.88
1	Y	231	ASN	N-CA-C	-9.61	96.65	110.59
1	W	260	VAL	CA-C-N	9.31	129.07	119.76
1	W	260	VAL	C-N-CA	9.31	129.07	119.76

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	260	VAL	CA-C-N	9.19	128.95	119.76

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	X	213	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2328	0	2341	273	0
1	B	2343	0	2363	205	0
1	C	2305	0	2332	246	0
1	D	2278	0	2291	188	0
1	W	2286	0	2308	205	0
1	X	2321	0	2347	259	0
1	Y	2343	0	2363	183	0
1	Z	2271	0	2284	286	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	C	10	0	0	0	0
2	W	5	0	0	0	0
2	X	5	0	0	0	0
2	Y	5	0	0	0	0
2	Z	5	0	0	0	0
3	A	4	0	0	0	0
3	B	5	0	0	1	0
3	C	3	0	0	0	0
3	D	4	0	0	1	0
3	W	5	0	0	2	0
3	X	3	0	0	0	0
3	Y	6	0	0	0	0
3	Z	5	0	0	0	0
All	All	18550	0	18629	1835	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 49.

The worst 5 of 1835 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:228:LEU:HG	1:X:230:PRO:HD3	1.23	1.16
1:Z:339:LYS:HD3	1:Z:341:GLN:HE21	1.15	1.11
1:C:228:LEU:HG	1:C:230:PRO:HD3	1.37	1.07
1:Y:94:ARG:HD3	1:Y:140:ASP:O	1.58	1.04
1:B:189:GLU:O	1:B:190:VAL:HG23	1.58	1.01

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	293/365 (80%)	231 (79%)	46 (16%)	16 (6%)	1	8
1	B	294/365 (80%)	225 (76%)	60 (20%)	9 (3%)	3	19
1	C	289/365 (79%)	213 (74%)	61 (21%)	15 (5%)	1	9
1	D	286/365 (78%)	227 (79%)	44 (15%)	15 (5%)	1	9
1	W	287/365 (79%)	231 (80%)	40 (14%)	16 (6%)	1	8
1	X	292/365 (80%)	211 (72%)	66 (23%)	15 (5%)	1	10
1	Y	294/365 (80%)	229 (78%)	56 (19%)	9 (3%)	3	19
1	Z	285/365 (78%)	227 (80%)	42 (15%)	16 (6%)	1	8
All	All	2320/2920 (80%)	1794 (77%)	415 (18%)	111 (5%)	2	11

5 of 111 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	24	ALA

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	294	PRO
1	B	24	ALA
1	B	36	VAL
1	C	89	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	246/292 (84%)	220 (89%)	26 (11%)	6	27
1	B	249/292 (85%)	234 (94%)	15 (6%)	17	50
1	C	245/292 (84%)	228 (93%)	17 (7%)	14	45
1	D	241/292 (82%)	223 (92%)	18 (8%)	12	41
1	W	242/292 (83%)	222 (92%)	20 (8%)	10	37
1	X	246/292 (84%)	226 (92%)	20 (8%)	11	38
1	Y	249/292 (85%)	232 (93%)	17 (7%)	14	45
1	Z	240/292 (82%)	216 (90%)	24 (10%)	7	29
All	All	1958/2336 (84%)	1801 (92%)	157 (8%)	11	38

5 of 157 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	X	295	TRP
1	Z	179	GLN
1	Y	173	TYR
1	Y	295	TRP
1	Z	257	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 114 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	336	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	Z	231	ASN
1	W	336	GLN
1	Z	219	HIS
1	Z	25	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	X	364	-	4,4,4	0.43	0	6,6,6	0.17	0
2	SO4	Y	364	-	4,4,4	0.49	0	6,6,6	0.08	0
2	SO4	Z	364	-	4,4,4	0.39	0	6,6,6	0.14	0
2	SO4	A	364	-	4,4,4	0.41	0	6,6,6	0.08	0
2	SO4	B	364	-	4,4,4	0.43	0	6,6,6	0.17	0
2	SO4	W	364	-	4,4,4	0.42	0	6,6,6	0.10	0
2	SO4	C	365	-	4,4,4	0.41	0	6,6,6	0.10	0
2	SO4	C	364	-	4,4,4	0.41	0	6,6,6	0.18	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	303/365 (83%)	0.30	8 (2%) 57 34	21, 72, 92, 92	0
1	B	304/365 (83%)	-0.14	2 (0%) 84 66	11, 47, 80, 92	0
1	C	299/365 (81%)	0.09	3 (1%) 79 59	10, 60, 89, 92	0
1	D	296/365 (81%)	-0.01	3 (1%) 79 59	15, 60, 91, 92	0
1	W	297/365 (81%)	-0.11	1 (0%) 90 79	15, 59, 92, 92	0
1	X	302/365 (82%)	0.04	3 (0%) 79 59	13, 61, 91, 92	0
1	Y	304/365 (83%)	-0.10	3 (0%) 79 59	10, 47, 86, 92	0
1	Z	295/365 (80%)	0.28	5 (1%) 69 45	18, 73, 92, 92	0
All	All	2400/2920 (82%)	0.04	28 (1%) 76 55	10, 59, 92, 92	0

The worst 5 of 28 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Z	314	GLY	3.6
1	X	6	ALA	3.4
1	C	345	THR	3.1
1	Y	345	THR	3.0
1	Z	315	GLY	2.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SO4	W	364	5/5	0.82	0.12	92,92,92,92	0
2	SO4	C	364	5/5	0.87	0.17	92,92,92,92	0
2	SO4	C	365	5/5	0.88	0.11	92,92,92,92	0
2	SO4	X	364	5/5	0.88	0.17	90,90,90,90	0
2	SO4	Y	364	5/5	0.89	0.14	92,92,92,92	0
2	SO4	B	364	5/5	0.92	0.12	92,92,92,92	0
2	SO4	A	364	5/5	0.92	0.08	88,88,88,88	0
2	SO4	Z	364	5/5	0.93	0.08	91,91,91,91	0

6.5 Other polymers [i](#)

There are no such residues in this entry.