



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 04:06 PM UTC

PDB ID : 4XD5 / pdb_00004xd5
Title : Phosphotriesterase variant R2
Authors : Campbell, E.; Kaltenbach, M.; Tokuriki, N.; Jackson, C.J.
Deposited on : 2014-12-19
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	333	7% 86% 12% ..
1	G	333	2% 86% 11% ..

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5809 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Phosphotriesterase variant PTE-R2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	331	Total	C	N	O	S	0	25	0
			2706	1706	495	498	7			
1	G	329	Total	C	N	O	S	0	25	0
			2681	1692	482	499	8			

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		
2	G	2	Total	Zn	0	0
			2	2		

- Molecule 3 is CACODYLATE ION (CCD ID: CAC) (formula: C₂H₆AsO₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	As	C	O	0	0
			5	1	2	2		
3	G	1	Total	As	C	O	0	0
			5	1	2	2		

- Molecule 4 is (4S)-2-METHYL-2,4-PENTANEDIOL (CCD ID: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			8	6	2		
4	A	1	Total	C	O	0	0
			8	6	2		
4	G	1	Total	C	O	0	0
			8	6	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	181	Total	O	0	0
			181	181		
5	G	203	Total	O	0	0
			203	203		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Phosphotriesterase variant PTE-R2



• Molecule 1: Phosphotriesterase variant PTE-R2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	85.70Å 85.87Å 88.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.19 – 1.85 44.19 – 1.85	Depositor EDS
% Data completeness (in resolution range)	90.5 (44.19-1.85) 87.4 (44.19-1.85)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.88 (at 1.86Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.161 , 0.204 0.165 , 0.208	Depositor DCC
R_{free} test set	2565 reflections (3.61%)	wwPDB-VP
Wilson B-factor (Å ²)	16.9	Xtriage
Anisotropy	0.559	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 54.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.017 for -h,l,k 0.017 for -l,-k,-h 0.018 for k,h,-l 0.007 for k,l,h 0.007 for l,h,k	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5809	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: KCX, MPD, ZN, CAC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.46	0/2774	0.74	1/3760 (0.0%)
1	G	0.45	0/2735	0.73	2/3708 (0.1%)
All	All	0.45	0/5509	0.73	3/7468 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	255	ILE	CA-C-N	-5.27	114.60	120.45
1	G	255	ILE	C-N-CA	-5.27	114.60	120.45
1	A	255	ILE	CB-CA-C	-5.22	108.74	113.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2706	0	2744	24	0
1	G	2681	0	2712	25	0
2	A	2	0	0	0	0
2	G	2	0	0	0	0
3	A	5	0	0	0	0
3	G	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	16	0	28	4	0
4	G	8	0	14	1	0
5	A	181	0	0	1	0
5	G	203	0	0	2	1
All	All	5809	0	5498	48	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:343:GLN:HE22	4:A:2404:MPD:HM2	1.58	0.67
1:G:131:TRP:CG	1:G:132:LEU:H	2.16	0.63
1:A:69:TRP:HE1	4:A:2405:MPD:H11	1.63	0.62
1:G:37[A]:ILE:HG22	1:G:360:PRO:HD3	1.80	0.62
1:A:259:ALA:HB1	1:A:262:LEU:HD13	1.82	0.60
1:A:87:LEU:HD12	1:A:116[B]:VAL:HG12	1.83	0.59
1:A:131:TRP:CG	1:A:132:LEU:H	2.22	0.58
4:A:2404:MPD:HO2	4:A:2404:MPD:HO4	1.52	0.57
1:G:286:ALA:O	1:G:290[B]:GLN:HG2	2.05	0.56
1:A:344:GLU:CD	1:A:344:GLU:H	2.15	0.54
1:G:285:LYS:NZ	1:G:289:ASP:OD2	2.34	0.54
1:G:302:TRP:CH2	1:G:321:ASN:HB3	2.42	0.53
1:A:254[A]:ARG:NH2	1:A:271:LEU:O	2.43	0.52
1:A:175:LYS:HA	1:A:211[A]:GLN:HE21	1.75	0.51
1:A:259:ALA:HB2	1:A:272:LEU:HB3	1.92	0.51
1:G:139:ARG:HA	1:G:179:PHE:CE1	2.45	0.50
1:A:67[B]:ARG:HG2	1:G:159:GLU:HG2	1.94	0.50
1:A:57:HIS:HB2	1:A:303:LEU:HB3	1.93	0.50
1:G:57:HIS:HB2	1:G:303:LEU:HB3	1.95	0.49
1:A:46:ILE:HG23	1:A:355:ALA:HB1	1.96	0.48
1:A:89[A]:ARG:HG3	5:A:2523:HOH:O	2.14	0.48
1:A:162:GLY:O	1:A:164[A]:ARG:NH1	2.48	0.46
1:G:131:TRP:CG	1:G:132:LEU:N	2.84	0.46
1:G:300:ASN:OD1	1:G:328:ILE:HG12	2.16	0.46
1:A:148[B]:GLN:HG3	1:A:149:PHE:N	2.29	0.45
1:A:302:TRP:CH2	1:A:321:ASN:HB3	2.51	0.45
1:G:173:THR:O	1:G:175[A]:LYS:N	2.50	0.45
1:G:57:HIS:O	1:G:303:LEU:HA	2.17	0.45
1:A:333:ILE:HB	1:A:334:PRO:HD3	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:172:THR:HG22	1:G:180:GLN:HB3	1.98	0.45
1:G:89[B]:ARG:NH2	5:G:606:HOH:O	2.48	0.44
1:G:173:THR:O	1:G:175[B]:LYS:N	2.50	0.44
1:G:45[A]:THR:OG1	1:G:48[A]:GLU:HG3	2.18	0.44
1:A:339:LYS:HA	1:A:339:LYS:HD3	1.82	0.43
1:G:333:ILE:HG23	1:G:346:LEU:HD13	2.01	0.43
1:G:353:ASN:HB2	1:G:354:PRO:HD3	2.01	0.43
1:A:131:TRP:CG	1:A:132:LEU:N	2.87	0.43
1:A:353:ASN:HB2	1:A:354:PRO:HD3	2.00	0.43
1:A:51:PHE:C	1:A:51:PHE:CD1	2.97	0.42
4:A:2405:MPD:H4	1:G:145:GLU:HG2	2.01	0.42
1:A:57:HIS:O	1:A:303:LEU:HA	2.20	0.42
1:G:136[B]:LEU:HG	1:G:140:LEU:HD22	2.02	0.42
1:A:308:SER:HB3	1:G:136[B]:LEU:HD23	2.01	0.42
1:G:169:KCX:OQ1	1:G:201:HIS:HB2	2.19	0.41
1:G:51:PHE:CD2	4:G:404:MPD:H31	2.55	0.41
1:G:134:PRO:HA	1:G:135[A]:PRO:HD3	1.82	0.41
1:G:152[B]:ARG:NH2	5:G:618:HOH:O	2.46	0.40
1:A:51:PHE:C	1:A:51:PHE:HD1	2.28	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:G:541:HOH:O	5:G:666:HOH:O[4_456]	2.12	0.08

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	353/333 (106%)	336 (95%)	14 (4%)	3 (1%)	16 6

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	G	350/333 (105%)	341 (97%)	8 (2%)	1 (0%)	36	25
All	All	703/666 (106%)	677 (96%)	22 (3%)	4 (1%)	21	10

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	174	GLY
1	A	176	VAL
1	A	260	ILE
1	A	267	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	287/266 (108%)	280 (98%)	7 (2%)	43	28
1	G	284/266 (107%)	276 (97%)	8 (3%)	38	23
All	All	571/532 (107%)	556 (97%)	15 (3%)	42	25

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	152	ARG
1	A	263	GLU
1	A	264	ASP
1	A	271	LEU
1	A	299	SER
1	A	303	LEU
1	A	363	ARG
1	G	46[A]	ILE
1	G	46[B]	ILE
1	G	140	LEU
1	G	152[A]	ARG
1	G	152[B]	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	G	211	GLN
1	G	299	SER
1	G	303	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	KCX	G	169	2,1	10,11,12	1.20	1 (10%)	6,12,14	1.33	1 (16%)
1	KCX	A	169	2,1	10,11,12	1.10	1 (10%)	6,12,14	1.44	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	G	169	2,1	-	1/9/10/12	-
1	KCX	A	169	2,1	-	0/9/10/12	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	169	KCX	CE-NZ	2.37	1.51	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	169	KCX	OQ1-CX	2.36	1.26	1.21

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	169	KCX	OQ1-CX-NZ	-3.31	119.89	124.92
1	G	169	KCX	OQ1-CX-NZ	-2.94	120.46	124.92

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	G	169	KCX	C-CA-CB-CG

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	G	169	KCX	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 4 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	MPD	A	2404	-	7,7,7	0.30	0	9,10,10	0.22	0
3	CAC	G	401	2	2,4,4	2.84	2 (100%)	4,6,6	1.39	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	MPD	A	2405	-	7,7,7	0.30	0	9,10,10	0.42	0
4	MPD	G	404	-	7,7,7	0.36	0	9,10,10	0.36	0
3	CAC	A	2403	2	2,4,4	2.74	2 (100%)	4,6,6	1.63	1 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MPD	A	2404	-	-	4/5/5/5	-
4	MPD	G	404	-	-	5/5/5/5	-
4	MPD	A	2405	-	-	1/5/5/5	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	G	401	CAC	AS-C2	2.98	1.97	1.90
3	A	2403	CAC	AS-C1	2.76	1.96	1.90
3	A	2403	CAC	AS-C2	2.73	1.96	1.90
3	G	401	CAC	AS-C1	2.68	1.96	1.90

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	2403	CAC	O2-AS-C2	2.52	111.94	105.84

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	2404	MPD	C2-C3-C4-C5
4	G	404	MPD	C2-C3-C4-C5
4	G	404	MPD	C1-C2-C3-C4
4	G	404	MPD	CM-C2-C3-C4
4	A	2405	MPD	C2-C3-C4-C5
4	G	404	MPD	O2-C2-C3-C4
4	A	2404	MPD	C2-C3-C4-O4
4	G	404	MPD	C2-C3-C4-O4
4	A	2404	MPD	C1-C2-C3-C4
4	A	2404	MPD	CM-C2-C3-C4

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	2404	MPD	2	0
4	A	2405	MPD	2	0
4	G	404	MPD	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	330/333 (99%)	-0.03	22 (6%) 24 26	6, 16, 71, 145	25 (7%)
1	G	328/333 (98%)	-0.13	8 (2%) 59 63	6, 19, 38, 64	24 (7%)
All	All	658/666 (98%)	-0.08	30 (4%) 37 40	6, 18, 47, 145	49 (7%)

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	271	LEU	7.0
1	A	270	ALA	5.9
1	A	274	ILE	5.9
1	A	272	LEU	5.8
1	A	268	ALA	5.6
1	A	262	LEU	5.2
1	A	266	ALA	4.9
1	A	260	ILE	4.8
1	A	259	ALA	4.5
1	A	261	GLY	4.3
1	A	267	SER	4.1
1	G	362	LEU	4.0
1	A	273	GLY	3.9
1	A	264	ASP	3.9
1	A	362	LEU	3.8
1	A	265	ASN	3.5
1	A	269	SER	3.5
1	A	364	ALA	3.3
1	G	34	GLY	3.3
1	A	263	GLU	3.2
1	A	35	ASP	2.9
1	G	35	ASP	2.9
1	A	275	ARG	2.7
1	A	363	ARG	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	34	GLY	2.6
1	G	204	ALA	2.4
1	G	173	THR	2.4
1	G	203	ALA	2.1
1	G	262	LEU	2.1
1	G	205	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	KCX	A	169	12/13	0.95	0.07	8,10,16,17	0
1	KCX	G	169	12/13	0.96	0.06	8,15,17,19	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MPD	A	2404	8/8	0.77	0.18	37,47,62,66	0
4	MPD	A	2405	8/8	0.79	0.14	42,48,62,72	0
4	MPD	G	404	8/8	0.79	0.18	44,54,58,60	0
3	CAC	A	2403	5/5	0.96	0.13	9,18,25,26	5
3	CAC	G	401	5/5	0.98	0.08	17,19,26,26	5
2	ZN	G	403	1/1	0.99	0.02	17,17,17,17	1
2	ZN	A	2402	1/1	0.99	0.03	14,14,14,14	1
2	ZN	A	2401	1/1	1.00	0.01	13,13,13,13	0
2	ZN	G	402	1/1	1.00	0.01	14,14,14,14	0

6.5 Other polymers [i](#)

There are no such residues in this entry.