



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 11:38 PM UTC

PDB ID : 4XDL / pdb_00004xdl
Title : Crystal structure of human two pore domain potassium ion channel TREK2 (K2P10.1) in complex with a brominated fluoxetine derivative.
Authors : Mackenzie, A.; Pike, A.C.W.; Dong, Y.Y.; Mukhopadhyay, S.; Ruda, G.F.; Brennan, P.E.; Arrowsmith, C.H.; Edwards, A.M.; Bountra, C.; Burgess-Brown, N.A.; Carpenter, E.P.; Structural Genomics Consortium (SGC)
Deposited on : 2014-12-19
Resolution : 3.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1085 (3.54-3.46)
Clashscore	190562	1140 (3.54-3.46)
Ramachandran outliers	187476	1113 (3.54-3.46)
Sidechain outliers	187428	1114 (3.54-3.46)
RSRZ outliers	180081	1084 (3.54-3.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	282	8% 84% 9% 7%
1	B	282	9% 84% 7% 9%
1	C	282	4% 78% 5% 17%
1	D	282	8% 85% 6% 9%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PC1	D	403	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7566 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Potassium channel subfamily K member 10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	261	Total	C	N	O	S	0	0	0
			1926	1283	298	341	4			
1	B	257	Total	C	N	O	S	0	0	0
			1914	1283	292	335	4			
1	C	234	Total	C	N	O	S	0	0	0
			1699	1136	261	299	3			
1	D	258	Total	C	N	O	S	0	0	0
			1935	1290	304	337	4			

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	66	MET	-	initiating methionine	UNP P57789
A	341	ALA	-	expression tag	UNP P57789
A	342	GLU	-	expression tag	UNP P57789
A	343	ASN	-	expression tag	UNP P57789
A	344	LEU	-	expression tag	UNP P57789
A	345	TYR	-	expression tag	UNP P57789
A	346	PHE	-	expression tag	UNP P57789
A	347	GLN	-	expression tag	UNP P57789
B	66	MET	-	initiating methionine	UNP P57789
B	341	ALA	-	expression tag	UNP P57789
B	342	GLU	-	expression tag	UNP P57789
B	343	ASN	-	expression tag	UNP P57789
B	344	LEU	-	expression tag	UNP P57789
B	345	TYR	-	expression tag	UNP P57789
B	346	PHE	-	expression tag	UNP P57789
B	347	GLN	-	expression tag	UNP P57789
C	66	MET	-	initiating methionine	UNP P57789
C	341	ALA	-	expression tag	UNP P57789
C	342	GLU	-	expression tag	UNP P57789
C	343	ASN	-	expression tag	UNP P57789
C	344	LEU	-	expression tag	UNP P57789

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	345	TYR	-	expression tag	UNP P57789
C	346	PHE	-	expression tag	UNP P57789
C	347	GLN	-	expression tag	UNP P57789
D	66	MET	-	initiating methionine	UNP P57789
D	341	ALA	-	expression tag	UNP P57789
D	342	GLU	-	expression tag	UNP P57789
D	343	ASN	-	expression tag	UNP P57789
D	344	LEU	-	expression tag	UNP P57789
D	345	TYR	-	expression tag	UNP P57789
D	346	PHE	-	expression tag	UNP P57789
D	347	GLN	-	expression tag	UNP P57789

- Molecule 2 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total K 2 2	0	0
2	B	1	Total K 1 1	0	0
2	C	3	Total K 3 3	0	0

- Molecule 3 is 1,2-DIACYL-SN-GLYCERO-3-PHOSPHOCHOLINE (CCD ID: PC1) (formula: $C_{44}H_{88}NO_8P$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C 11 11	0	0
3	A	1	Total C 8 8	0	0
3	D	1	Total C 9 9	0	0
3	D	1	Total C 8 8	0	0

- Molecule 4 is CADMIUM ION (CCD ID: CD) (formula: Cd).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Cd 1 1	0	0
4	D	1	Total Cd 1 1	0	0

- Molecule 5 is 3-[2-bromanyl-4-(trifluoromethyl)phenoxy]-N-methyl-3-phenyl-propan-1-amine (CCD ID: 40D) (formula: C₁₇H₁₇BrF₃NO).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Br C F O 12 1 7 3 1	0	0
5	B	1	Total Br C F O 12 1 7 3 1	0	0
5	C	1	Total Br C F O 12 1 7 3 1	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	D	1	Total	Br	C	F	O	0	0
			12	1	7	3	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Potassium channel subfamily K member 10




- Molecule 1: Potassium channel subfamily K member 10




- Molecule 1: Potassium channel subfamily K member 10



- Molecule 1: Potassium channel subfamily K member 10





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	101.71Å 109.84Å 166.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.00 – 3.50 35.00 – 3.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (35.00-3.50) 99.8 (35.00-3.50)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.92 (at 3.49Å)	Xtriage
Refinement program	BUSTER 2.10.1	Depositor
R, R_{free}	0.235 , 0.263 0.281 , 0.307	Depositor DCC
R_{free} test set	1217 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	145.7	Xtriage
Anisotropy	0.208	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 122.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	7566	wwPDB-VP
Average B, all atoms (Å ²)	157.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: K, 40D, CD, PC1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.82	1/1977 (0.1%)	1.47	7/2710 (0.3%)
1	B	0.81	0/1966	1.47	10/2696 (0.4%)
1	C	0.81	1/1744 (0.1%)	1.48	5/2396 (0.2%)
1	D	0.81	0/1986	1.46	8/2718 (0.3%)
All	All	0.82	2/7673 (0.0%)	1.47	30/10520 (0.3%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	255	ILE	CA-CB	5.15	1.56	1.54
1	C	255	ILE	CA-CB	5.07	1.56	1.54

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	286	ASP	CA-CB-CG	5.57	118.17	112.60
1	D	215	PHE	CA-CB-CG	5.51	119.31	113.80
1	B	250	ILE	CA-C-N	5.46	127.44	120.56
1	B	250	ILE	C-N-CA	5.46	127.44	120.56
1	B	169	VAL	CA-C-N	5.45	127.92	120.46
1	B	169	VAL	C-N-CA	5.45	127.92	120.46
1	C	286	ASP	CA-CB-CG	5.39	118.00	112.60
1	D	321	SER	CA-C-N	5.36	127.46	120.28
1	D	321	SER	C-N-CA	5.36	127.46	120.28
1	A	321	SER	CA-C-N	5.36	127.46	120.28
1	A	321	SER	C-N-CA	5.36	127.46	120.28
1	A	286	ASP	CA-CB-CG	5.33	117.93	112.60
1	B	321	SER	CA-C-N	5.29	127.37	120.28
1	B	321	SER	C-N-CA	5.29	127.37	120.28
1	D	286	ASP	CA-CB-CG	5.28	117.88	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	250	ILE	CA-C-N	5.27	127.20	120.56
1	C	250	ILE	C-N-CA	5.27	127.20	120.56
1	A	250	ILE	CA-C-N	5.26	127.18	120.56
1	A	250	ILE	C-N-CA	5.26	127.18	120.56
1	D	250	ILE	CA-C-N	5.24	127.16	120.56
1	D	250	ILE	C-N-CA	5.24	127.16	120.56
1	B	254	THR	CA-C-N	5.24	125.46	120.43
1	B	254	THR	C-N-CA	5.24	125.46	120.43
1	B	177	ASN	N-CA-C	-5.23	105.66	111.36
1	C	254	THR	CA-C-N	5.09	125.32	120.43
1	C	254	THR	C-N-CA	5.09	125.32	120.43
1	A	169	VAL	CA-C-N	5.02	128.40	120.47
1	A	169	VAL	C-N-CA	5.02	128.40	120.47
1	D	254	THR	CA-C-N	5.01	125.24	120.43
1	D	254	THR	C-N-CA	5.01	125.24	120.43

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1926	0	1840	14	0
1	B	1914	0	1856	11	0
1	C	1699	0	1594	7	0
1	D	1935	0	1894	8	0
2	A	2	0	0	0	0
2	B	1	0	0	0	0
2	C	3	0	0	0	0
3	A	19	0	33	0	0
3	D	17	0	26	0	0
4	A	1	0	0	0	0
4	D	1	0	0	0	0
5	A	12	0	3	0	0
5	B	12	0	3	0	0
5	C	12	0	3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	D	12	0	3	0	0
All	All	7566	0	7255	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:170:ILE:HD12	1:A:196:GLY:HA3	1.79	0.64
1:B:170:ILE:HD12	1:B:196:GLY:HA3	1.81	0.63
1:D:170:ILE:CG2	1:D:200:PHE:HB2	2.32	0.59
1:D:170:ILE:HD12	1:D:196:GLY:HA3	1.86	0.57
1:C:170:ILE:HD12	1:C:196:GLY:HA3	1.87	0.56
1:A:144:VAL:HG11	1:B:107:LYS:HA	1.90	0.54
1:C:170:ILE:CG2	1:C:200:PHE:HB2	2.35	0.54
1:A:190:ILE:HG23	1:B:276:VAL:HG11	1.89	0.54
1:A:74:TRP:HD1	1:B:210:GLN:HG3	1.73	0.54
1:A:170:ILE:CG2	1:A:200:PHE:HB2	2.38	0.54
1:A:116:GLU:HG3	1:D:116:GLU:HG3	1.93	0.51
1:B:170:ILE:CG2	1:B:200:PHE:HB2	2.43	0.49
1:D:170:ILE:HG22	1:D:200:PHE:HB2	1.94	0.48
1:B:307:PHE:O	1:B:311:VAL:HG23	2.15	0.47
1:C:307:PHE:O	1:C:311:VAL:HG23	2.15	0.47
1:A:170:ILE:HG22	1:A:200:PHE:HB2	1.97	0.47
1:A:253:VAL:HA	1:A:275:VAL:HG11	1.97	0.47
1:C:144:VAL:HG11	1:D:107:LYS:HA	1.97	0.45
1:C:170:ILE:HG22	1:C:200:PHE:HB2	1.98	0.45
1:D:307:PHE:O	1:D:311:VAL:HG23	2.17	0.44
1:A:107:LYS:HA	1:B:144:VAL:HG11	1.99	0.44
1:D:253:VAL:HA	1:D:275:VAL:HG11	1.99	0.43
1:A:99:GLU:OE2	1:B:185:GLY:HA3	2.18	0.43
1:A:208:GLY:HA2	1:A:318:ALA:HB2	2.01	0.42
1:A:276:VAL:HG21	1:B:190:ILE:HG23	2.01	0.42
1:C:253:VAL:HA	1:C:275:VAL:HG11	2.00	0.42
1:A:300:TRP:C	1:A:303:PRO:HD2	2.45	0.42
1:C:107:LYS:HA	1:D:144:VAL:HG11	2.00	0.42
1:B:253:VAL:HA	1:B:275:VAL:HG11	2.03	0.41
1:B:275:VAL:HG13	1:B:308:TRP:HZ2	1.86	0.41
1:A:275:VAL:HG13	1:A:308:TRP:HZ2	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	259/282 (92%)	249 (96%)	9 (4%)	1 (0%)	30	62
1	B	255/282 (90%)	246 (96%)	9 (4%)	0	100	100
1	C	228/282 (81%)	220 (96%)	7 (3%)	1 (0%)	30	62
1	D	256/282 (91%)	247 (96%)	8 (3%)	1 (0%)	30	62
All	All	998/1128 (88%)	962 (96%)	33 (3%)	3 (0%)	36	67

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	283	GLY
1	A	283	GLY
1	D	283	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/233 (79%)	183 (99%)	2 (1%)	65	74
1	B	188/233 (81%)	187 (100%)	1 (0%)	81	80
1	C	159/233 (68%)	158 (99%)	1 (1%)	78	79
1	D	192/233 (82%)	191 (100%)	1 (0%)	81	80

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	724/932 (78%)	719 (99%)	5 (1%)	76	78

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	156	HIS
1	A	204	LEU
1	B	156	HIS
1	C	156	HIS
1	D	156	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	106	GLN
1	B	127	GLN
1	B	135	HIS
1	C	127	GLN
1	C	135	HIS
1	D	100	GLN
1	D	149	ASN
1	D	152	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 16 ligands modelled in this entry, 8 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PC1	A	603	-	10,10,53	0.29	0	9,9,61	0.13	0
3	PC1	D	402	-	8,8,53	0.22	0	7,7,61	0.13	0
5	40D	C	604	-	12,12,24	0.27	0	18,18,33	0.66	0
5	40D	D	404	-	12,12,24	0.34	0	18,18,33	0.64	0
5	40D	B	402	-	12,12,24	0.33	0	18,18,33	0.62	0
3	PC1	A	606	-	7,7,53	0.20	0	6,6,61	0.12	0
5	40D	A	605	-	12,12,24	0.39	0	18,18,33	0.63	0
3	PC1	D	403	-	7,7,53	0.15	0	6,6,61	0.12	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PC1	A	603	-	-	0/8/8/57	-
3	PC1	D	402	-	-	0/6/6/57	-
5	40D	C	604	-	-	0/6/6/18	0/1/1/2
5	40D	D	404	-	-	0/6/6/18	0/1/1/2
5	40D	B	402	-	-	0/6/6/18	0/1/1/2
3	PC1	A	606	-	-	0/5/5/57	-
5	40D	A	605	-	-	0/6/6/18	0/1/1/2
3	PC1	D	403	-	-	0/5/5/57	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	261/282 (92%)	0.46	22 (8%) 17 11	98, 146, 214, 235	0
1	B	257/282 (91%)	0.45	25 (9%) 13 9	97, 148, 198, 225	0
1	C	234/282 (82%)	0.57	11 (4%) 36 20	116, 159, 244, 261	0
1	D	258/282 (91%)	0.71	23 (8%) 15 10	109, 154, 209, 235	0
All	All	1010/1128 (89%)	0.55	81 (8%) 18 12	97, 152, 218, 261	0

All (81) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	285	GLY	7.6
1	D	286	ASP	6.6
1	D	187	ILE	6.6
1	B	90	THR	6.2
1	A	195	PHE	6.0
1	D	280	THR	5.3
1	D	279	LEU	5.1
1	D	127	GLN	4.8
1	A	285	GLY	4.6
1	D	278	THR	4.6
1	B	231	VAL	4.6
1	B	180	PRO	4.5
1	A	191	LEU	4.0
1	A	78	VAL	3.9
1	D	284	PHE	3.9
1	B	181	SER	3.8
1	B	79	ALA	3.7
1	A	99	GLU	3.7
1	B	179	ALA	3.6
1	D	243	LEU	3.5
1	A	255	ILE	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	79	ALA	3.4
1	B	82	VAL	3.4
1	A	286	ASP	3.4
1	B	157	TRP	3.4
1	D	281	THR	3.3
1	C	147	ILE	3.3
1	B	222	VAL	3.3
1	B	182	THR	3.2
1	C	172	THR	3.2
1	D	175	TYR	3.2
1	C	213	THR	3.1
1	D	184	GLY	3.1
1	B	138	ASP	3.1
1	A	95	PHE	3.1
1	D	283	GLY	3.0
1	A	188	PHE	3.0
1	A	239	ILE	2.9
1	D	239	ILE	2.9
1	A	222	VAL	2.8
1	C	250	ILE	2.8
1	C	138	ASP	2.7
1	C	216	GLY	2.7
1	B	185	GLY	2.7
1	C	176	GLY	2.7
1	B	142	ALA	2.6
1	B	320	LEU	2.6
1	A	302	LYS	2.6
1	C	137	LEU	2.6
1	B	83	VAL	2.5
1	A	75	LYS	2.5
1	D	92	GLY	2.5
1	A	243	LEU	2.5
1	B	176	GLY	2.5
1	A	81	PHE	2.5
1	D	308	TRP	2.5
1	B	232	SER	2.5
1	B	80	ILE	2.4
1	A	202	PHE	2.3
1	D	130	GLU	2.3
1	B	226	PHE	2.3
1	D	313	LEU	2.3
1	D	173	ILE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	282	VAL	2.2
1	C	152	ASN	2.2
1	B	78	VAL	2.2
1	D	274	PHE	2.2
1	C	300	TRP	2.1
1	A	176	GLY	2.1
1	D	303	PRO	2.1
1	B	316	PHE	2.1
1	B	86	VAL	2.1
1	A	97	ALA	2.1
1	A	142	ALA	2.0
1	B	134	GLN	2.0
1	C	93	LEU	2.0
1	A	262	TYR	2.0
1	A	74	TRP	2.0
1	B	204	LEU	2.0
1	B	313	LEU	2.0
1	D	304	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PC1	D	403	8/54	0.64	0.45	117,119,126,127	0
5	40D	B	402	12/23	0.75	0.23	202,205,209,209	1
5	40D	A	605	12/23	0.76	0.18	169,172,174,176	1
5	40D	C	604	12/23	0.79	0.21	226,228,231,234	1
3	PC1	A	606	8/54	0.80	0.48	148,156,160,164	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	PC1	D	402	9/54	0.80	0.35	125,131,145,150	0
3	PC1	A	603	11/54	0.85	0.25	114,118,131,135	0
2	K	C	603	1/1	0.90	0.14	141,141,141,141	0
5	40D	D	404	12/23	0.90	0.16	189,192,194,198	1
2	K	A	601	1/1	0.95	0.09	100,100,100,100	0
2	K	C	601	1/1	0.97	0.12	113,113,113,113	0
2	K	B	401	1/1	0.98	0.03	104,104,104,104	0
4	CD	D	401	1/1	0.98	0.09	136,136,136,136	0
2	K	C	602	1/1	0.98	0.11	128,128,128,128	0
4	CD	A	604	1/1	0.99	0.10	125,125,125,125	0
2	K	A	602	1/1	0.99	0.05	122,122,122,122	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.