



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 30, 2026 – 02:11 AM UTC

PDB ID : 2XE4 / pdb_00002xe4
Title : Structure of Oligopeptidase B from Leishmania major
Authors : McLuskey, K.; Paterson, N.G.; Bland, N.D.; Mottram, J.C.; Isaacs, N.W.
Deposited on : 2010-05-11
Resolution : 1.65 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.66-1.66)
Clashscore	190562	2662 (1.66-1.66)
Ramachandran outliers	187476	2621 (1.66-1.66)
Sidechain outliers	187428	2621 (1.66-1.66)
RSRZ outliers	180081	2564 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	751	 76% 19%
2	B	4	 25% 50% 25%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PGR	A	1747	-	X	-	-
4	PGR	A	1767	-	X	-	-
4	PGR	A	1770	-	X	-	-
4	PGR	A	1782	-	-	X	-
4	PGR	A	1783	-	X	-	-
4	PGR	A	1785	-	X	-	-
4	PGR	A	1787	-	X	-	-
4	PGR	A	1788	-	X	-	-
5	PGO	A	1743	X	-	-	-

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 7215 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called OLIGOPEPTIDASE B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	721	6125	3890	1046	1148	41	0	51	0

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP Q4QHU7
A	-18	GLY	-	expression tag	UNP Q4QHU7
A	-17	SER	-	expression tag	UNP Q4QHU7
A	-16	SER	-	expression tag	UNP Q4QHU7
A	-15	HIS	-	expression tag	UNP Q4QHU7
A	-14	HIS	-	expression tag	UNP Q4QHU7
A	-13	HIS	-	expression tag	UNP Q4QHU7
A	-12	HIS	-	expression tag	UNP Q4QHU7
A	-11	HIS	-	expression tag	UNP Q4QHU7
A	-10	HIS	-	expression tag	UNP Q4QHU7
A	-9	SER	-	expression tag	UNP Q4QHU7
A	-8	SER	-	expression tag	UNP Q4QHU7
A	-7	GLY	-	expression tag	UNP Q4QHU7
A	-6	LEU	-	expression tag	UNP Q4QHU7
A	-5	VAL	-	expression tag	UNP Q4QHU7
A	-4	PRO	-	expression tag	UNP Q4QHU7
A	-3	ARG	-	expression tag	UNP Q4QHU7
A	-2	GLY	-	expression tag	UNP Q4QHU7
A	-1	SER	-	expression tag	UNP Q4QHU7
A	0	HIS	-	expression tag	UNP Q4QHU7
A	25	LEU	PHE	engineered mutation	UNP Q4QHU7

- Molecule 2 is a protein called ANTIPAIN.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	4	43	27	10	6	0	0	0

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	A	1	6	3	3	0	0
3	A	1	6	3	3	0	0
3	A	1	6	3	3	0	0
3	A	1	6	3	3	0	0
3	A	1	6	3	3	0	0
3	A	1	6	3	3	0	0
3	A	1	6	3	3	0	0
3	A	1	6	3	3	0	0
3	A	1	6	3	3	0	0
3	A	1	6	3	3	0	0

- Molecule 4 is R-1,2-PROPANEDIOL (CCD ID: PGR) (formula: C₃H₈O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			5	3	2		
4	A	1	Total	C	O	0	0
			5	3	2		
4	A	1	Total	C	O	0	0
			5	3	2		
4	A	1	Total	C	O	0	0
			5	3	2		
4	A	1	Total	C	O	0	0
			5	3	2		
4	A	1	Total	C	O	0	0
			5	3	2		
4	A	1	Total	C	O	0	0
			5	3	2		
4	A	1	Total	C	O	0	0
			5	3	2		
4	A	1	Total	C	O	0	0
			5	3	2		
4	A	1	Total	C	O	0	0
			5	3	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			5	3	2		
4	A	1	Total	C	O	0	0
			5	3	2		
4	A	1	Total	C	O	0	0
			5	3	2		
4	A	1	Total	C	O	0	0
			5	3	2		
4	A	1	Total	C	O	0	0
			5	3	2		
4	A	1	Total	C	O	0	0
			5	3	2		
4	A	1	Total	C	O	0	0
			5	3	2		
4	A	1	Total	C	O	0	0
			5	3	2		
4	A	1	Total	C	O	0	0
			5	3	2		
4	A	1	Total	C	O	0	0
			5	3	2		
4	A	1	Total	C	O	0	0
			5	3	2		
4	A	1	Total	C	O	0	0
			5	3	2		
4	A	1	Total	C	O	0	0
			5	3	2		
4	A	1	Total	C	O	0	0
			5	3	2		
4	A	1	Total	C	O	0	0
			5	3	2		
4	A	1	Total	C	O	0	0
			5	3	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 5 3 2	0	0
4	A	1	Total C O 5 3 2	0	0
4	A	1	Total C O 5 3 2	0	0
4	A	1	Total C O 5 3 2	0	0
4	A	1	Total C O 5 3 2	0	0
4	A	1	Total C O 5 3 2	0	0
4	A	1	Total C O 5 3 2	0	0
4	A	1	Total C O 5 3 2	0	0
4	A	1	Total C O 5 3 2	0	0
4	A	1	Total C O 5 3 2	0	0
4	A	1	Total C O 5 3 2	0	0
4	A	1	Total C O 5 3 2	0	0
4	A	1	Total C O 5 3 2	0	0
4	A	1	Total C O 5 3 2	0	0
4	A	1	Total C O 5 3 2	0	0

- Molecule 5 is S-1,2-PROPANEDIOL (CCD ID: PGO) (formula: C₃H₈O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			5	3	2		

- Molecule 6 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	P	0	0
			5	4	1		
6	A	1	Total	O	P	0	0
			5	4	1		
6	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 7 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	4	Total Cl 4 4	0	0

- Molecule 8 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	2	Total Na 2 2	0	0

- Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	721	Total O 721 721	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	95.48Å 142.78Å 208.92Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	117.85 – 1.65 117.85 – 1.65	Depositor EDS
% Data completeness (in resolution range)	98.0 (117.85-1.65) 98.0 (117.85-1.65)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.67 (at 1.65Å)	Xtrriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.140 , 0.178 0.146 , 0.182	Depositor DCC
R_{free} test set	8321 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	26.8	Xtrriage
Anisotropy	0.235	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 66.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.53$, $\langle L^2 \rangle = 0.36$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	7215	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PGO, GOL, RGL, NA, CL, PO4, PGR, FC0

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.81	66/6332 (1.0%)	1.19	8/8586 (0.1%)
2	B	5.20	4/17 (23.5%)	6.61	3/21 (14.3%)
All	All	1.83	70/6349 (1.1%)	1.24	11/8607 (0.1%)

The worst 5 of 70 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2	ARG	CZ-NH1	13.63	1.51	1.32
2	B	2	ARG	NE-CZ	10.53	1.44	1.33
1	A	663	PRO	N-CA	9.81	1.56	1.47
2	B	2	ARG	CD-NE	9.81	1.59	1.46
1	A	368	ALA	CA-CB	8.63	1.66	1.53

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	ARG	CD-NE-CZ	18.94	150.92	124.40
2	B	2	ARG	NE-CZ-NH2	-17.70	103.27	119.20
2	B	2	ARG	NE-CZ-NH1	14.36	135.86	121.50
1	A	585	ALA	N-CA-C	-5.71	104.97	111.14
1	A	551	SER	N-CA-C	-5.70	105.21	111.82

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6125	0	5995	95	0
2	B	43	0	42	1	0
3	A	60	0	79	11	0
4	A	240	0	382	34	0
5	A	5	0	7	1	0
6	A	15	0	0	1	0
7	A	4	0	0	0	0
8	A	2	0	0	0	0
9	A	721	0	0	21	0
All	All	7215	0	6505	108	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 108 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:712:ILE:CD1	1:A:712:ILE:CG1	1.75	1.60
1:A:448[B]:GLU:HG3	9:A:2471:HOH:O	1.61	1.00
1:A:157[B]:MET:HE1	9:A:2223:HOH:O	1.65	0.96
1:A:448[B]:GLU:CG	9:A:2471:HOH:O	2.16	0.93
1:A:396[B]:MET:HE3	1:A:422:TYR:CE1	2.12	0.85

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	770/751 (102%)	753 (98%)	16 (2%)	1 (0%)	48 30

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	1/4 (25%)	1 (100%)	0	0	100	100
All	All	771/755 (102%)	754 (98%)	16 (2%)	1 (0%)	48	30

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	325	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	675/649 (104%)	667 (99%)	8 (1%)	63	45
2	B	2/2 (100%)	2 (100%)	0	100	100
All	All	677/651 (104%)	669 (99%)	8 (1%)	68	45

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	691[B]	ILE
1	A	691[A]	ILE
1	A	397	GLU
1	A	256[B]	MET
1	A	642	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	661	HIS
1	A	657	GLN
1	A	381	GLN
1	A	340	HIS
1	A	437	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FC0	B	1	2	14,14,15	1.67	4 (28%)	16,17,19	0.97	1 (6%)
2	RGL	B	4	1,2	9,10,10	2.68	3 (33%)	5,11,11	2.70	3 (60%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FC0	B	1	2	-	6/11/11/12	0/1/1/1
2	RGL	B	4	1,2	-	0/8/9/9	-

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	4	RGL	O-C	6.77	1.45	1.20
2	B	1	FC0	CA-N	3.23	1.50	1.45
2	B	4	RGL	CZ-NE	3.08	1.39	1.33
2	B	4	RGL	CG-CB	2.23	1.61	1.52
2	B	1	FC0	CA-C	2.19	1.58	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	4	RGL	NE-CZ-NH2	-5.06	111.99	120.67

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	FC0	O1-C1-N	-2.51	118.84	125.32
2	B	4	RGL	NH1-CZ-NH2	2.38	126.85	120.07
2	B	4	RGL	CB-CG-CD	-2.07	106.11	112.07

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1	FC0	O-C-CA-N
2	B	1	FC0	OXT-C-CA-N
2	B	1	FC0	O-C-CA-CB
2	B	1	FC0	OXT-C-CA-CB
2	B	1	FC0	CA-CB-CG-CD2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1	FC0	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 68 ligands modelled in this entry, 6 are monoatomic - leaving 62 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	1741	-	5,5,5	1.19	0	5,5,5	1.98	2 (40%)
4	PGR	A	1775	-	4,4,4	0.54	0	4,4,4	1.71	1 (25%)
3	GOL	A	1734	-	5,5,5	0.61	0	5,5,5	1.07	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	PGR	A	1747	-	4,4,4	1.01	0	4,4,4	2.41	2 (50%)
4	PGR	A	1753	-	4,4,4	1.72	1 (25%)	4,4,4	0.60	0
4	PGR	A	1766	-	4,4,4	1.31	0	4,4,4	0.76	0
4	PGR	A	1770	-	4,4,4	0.96	0	4,4,4	1.70	2 (50%)
4	PGR	A	1754	-	4,4,4	0.80	0	4,4,4	1.40	1 (25%)
4	PGR	A	1762	-	4,4,4	0.43	0	4,4,4	1.40	1 (25%)
4	PGR	A	1748	-	4,4,4	0.69	0	4,4,4	1.02	0
4	PGR	A	1755	-	4,4,4	1.72	2 (50%)	4,4,4	0.59	0
4	PGR	A	1764	-	4,4,4	1.36	1 (25%)	4,4,4	1.05	0
3	GOL	A	1733	-	5,5,5	0.51	0	5,5,5	1.71	1 (20%)
4	PGR	A	1779	-	4,4,4	0.71	0	4,4,4	1.20	0
4	PGR	A	1784	-	4,4,4	1.84	2 (50%)	4,4,4	1.22	0
4	PGR	A	1756	-	4,4,4	0.50	0	4,4,4	1.44	1 (25%)
4	PGR	A	1746	-	4,4,4	1.88	1 (25%)	4,4,4	1.03	0
4	PGR	A	1790	-	4,4,4	1.36	0	4,4,4	1.16	0
4	PGR	A	1763	-	4,4,4	1.33	0	4,4,4	2.03	2 (50%)
6	PO4	A	1793	-	4,4,4	1.67	1 (25%)	6,6,6	1.12	1 (16%)
4	PGR	A	1745	-	4,4,4	0.74	0	4,4,4	1.52	1 (25%)
4	PGR	A	1768	-	4,4,4	0.86	0	4,4,4	1.08	0
4	PGR	A	1787	-	4,4,4	2.43	2 (50%)	4,4,4	2.01	2 (50%)
4	PGR	A	1765	-	4,4,4	0.81	0	4,4,4	1.87	1 (25%)
3	GOL	A	1736	-	5,5,5	0.72	0	5,5,5	0.85	0
3	GOL	A	1739	-	5,5,5	0.54	0	5,5,5	1.04	0
3	GOL	A	1732	-	5,5,5	1.62	1 (20%)	5,5,5	1.48	1 (20%)
4	PGR	A	1778	-	4,4,4	0.92	0	4,4,4	1.77	2 (50%)
4	PGR	A	1786	-	4,4,4	0.98	0	4,4,4	1.60	1 (25%)
6	PO4	A	1791	-	4,4,4	2.69	1 (25%)	6,6,6	1.82	2 (33%)
4	PGR	A	1773	-	4,4,4	1.08	0	4,4,4	0.79	0
4	PGR	A	1750	-	4,4,4	1.59	1 (25%)	4,4,4	0.80	0
3	GOL	A	1735	-	5,5,5	0.79	0	5,5,5	2.02	1 (20%)
4	PGR	A	1780	-	4,4,4	0.70	0	4,4,4	1.27	0
6	PO4	A	1792	-	4,4,4	1.26	0	6,6,6	0.72	0
4	PGR	A	1758	-	4,4,4	1.26	0	4,4,4	1.21	0
3	GOL	A	1737	-	5,5,5	0.38	0	5,5,5	0.55	0
4	PGR	A	1772	-	4,4,4	0.56	0	4,4,4	1.22	0
3	GOL	A	1738	-	5,5,5	0.79	0	5,5,5	0.61	0
4	PGR	A	1785	-	4,4,4	1.07	0	4,4,4	1.77	2 (50%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	PGR	A	1774	-	4,4,4	1.15	0	4,4,4	1.79	1 (25%)
4	PGR	A	1788	-	4,4,4	0.75	0	4,4,4	2.05	2 (50%)
3	GOL	A	1740	-	5,5,5	0.65	0	5,5,5	1.16	0
4	PGR	A	1769	-	4,4,4	1.12	0	4,4,4	0.84	0
4	PGR	A	1781	-	4,4,4	1.01	0	4,4,4	1.39	1 (25%)
4	PGR	A	1759	-	4,4,4	1.17	1 (25%)	4,4,4	0.56	0
4	PGR	A	1752	-	4,4,4	0.64	0	4,4,4	1.27	1 (25%)
4	PGR	A	1767	-	4,4,4	1.37	1 (25%)	4,4,4	2.24	2 (50%)
4	PGR	A	1771	-	4,4,4	1.66	1 (25%)	4,4,4	1.61	1 (25%)
4	PGR	A	1783	-	4,4,4	2.19	2 (50%)	4,4,4	0.40	0
4	PGR	A	1749	-	4,4,4	1.03	0	4,4,4	1.91	1 (25%)
4	PGR	A	1760	-	4,4,4	1.64	0	4,4,4	0.59	0
4	PGR	A	1789	-	4,4,4	1.12	0	4,4,4	1.88	2 (50%)
4	PGR	A	1757	-	4,4,4	1.08	0	4,4,4	1.17	1 (25%)
4	PGR	A	1776	-	4,4,4	1.87	2 (50%)	4,4,4	0.38	0
4	PGR	A	1744	-	4,4,4	0.83	0	4,4,4	1.33	1 (25%)
4	PGR	A	1751	-	4,4,4	2.28	2 (50%)	4,4,4	0.53	0
5	PGO	A	1743	-	4,4,4	1.07	0	4,4,4	2.53	2 (50%)
4	PGR	A	1782	-	4,4,4	1.36	1 (25%)	4,4,4	0.75	0
4	PGR	A	1777	-	4,4,4	1.26	1 (25%)	4,4,4	0.84	0
4	PGR	A	1761	-	4,4,4	1.80	1 (25%)	4,4,4	0.29	0
4	PGR	A	1742	-	4,4,4	1.30	0	4,4,4	1.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	1741	-	-	3/4/4/4	-
4	PGR	A	1775	-	-	0/2/2/2	-
3	GOL	A	1734	-	-	4/4/4/4	-
4	PGR	A	1747	-	-	2/2/2/2	-
4	PGR	A	1753	-	-	1/2/2/2	-
4	PGR	A	1766	-	-	2/2/2/2	-
4	PGR	A	1770	-	-	2/2/2/2	-
4	PGR	A	1754	-	-	2/2/2/2	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PGR	A	1762	-	-	2/2/2/2	-
4	PGR	A	1748	-	-	0/2/2/2	-
4	PGR	A	1755	-	-	0/2/2/2	-
4	PGR	A	1764	-	-	2/2/2/2	-
3	GOL	A	1733	-	-	4/4/4/4	-
4	PGR	A	1779	-	-	2/2/2/2	-
4	PGR	A	1784	-	-	1/2/2/2	-
4	PGR	A	1756	-	-	0/2/2/2	-
4	PGR	A	1746	-	-	2/2/2/2	-
4	PGR	A	1790	-	-	2/2/2/2	-
4	PGR	A	1763	-	-	1/2/2/2	-
4	PGR	A	1745	-	-	0/2/2/2	-
4	PGR	A	1768	-	-	2/2/2/2	-
4	PGR	A	1787	-	-	1/2/2/2	-
4	PGR	A	1765	-	-	0/2/2/2	-
3	GOL	A	1736	-	-	3/4/4/4	-
3	GOL	A	1739	-	-	4/4/4/4	-
3	GOL	A	1732	-	-	1/4/4/4	-
4	PGR	A	1778	-	-	0/2/2/2	-
4	PGR	A	1786	-	-	0/2/2/2	-
4	PGR	A	1773	-	-	1/2/2/2	-
4	PGR	A	1750	-	-	0/2/2/2	-
3	GOL	A	1735	-	-	4/4/4/4	-
4	PGR	A	1780	-	-	0/2/2/2	-
4	PGR	A	1758	-	-	0/2/2/2	-
3	GOL	A	1737	-	-	1/4/4/4	-
4	PGR	A	1772	-	-	1/2/2/2	-
3	GOL	A	1738	-	-	2/4/4/4	-
4	PGR	A	1785	-	-	2/2/2/2	-
4	PGR	A	1774	-	-	2/2/2/2	-
4	PGR	A	1788	-	-	2/2/2/2	-
3	GOL	A	1740	-	-	4/4/4/4	-
4	PGR	A	1769	-	-	1/2/2/2	-
4	PGR	A	1781	-	-	2/2/2/2	-
4	PGR	A	1759	-	-	2/2/2/2	-
4	PGR	A	1752	-	-	0/2/2/2	-
4	PGR	A	1767	-	-	2/2/2/2	-
4	PGR	A	1771	-	-	0/2/2/2	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PGR	A	1783	-	-	2/2/2/2	-
4	PGR	A	1749	-	-	1/2/2/2	-
4	PGR	A	1760	-	-	0/2/2/2	-
4	PGR	A	1789	-	-	1/2/2/2	-
4	PGR	A	1757	-	-	2/2/2/2	-
4	PGR	A	1776	-	-	0/2/2/2	-
5	PGO	A	1743	-	1/1/1/1	1/2/2/2	-
4	PGR	A	1744	-	-	0/2/2/2	-
4	PGR	A	1751	-	-	1/2/2/2	-
4	PGR	A	1782	-	-	2/2/2/2	-
4	PGR	A	1777	-	-	2/2/2/2	-
4	PGR	A	1761	-	-	0/2/2/2	-
4	PGR	A	1742	-	-	2/2/2/2	-

The worst 5 of 25 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	1791	PO4	P-O4	-5.11	1.39	1.54
4	A	1746	PGR	O1-C1	3.59	1.57	1.42
4	A	1787	PGR	C1-C2	3.56	1.64	1.46
4	A	1751	PGR	O2-C2	3.19	1.56	1.43
4	A	1771	PGR	O2-C2	3.16	1.56	1.43

The worst 5 of 42 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1735	GOL	O1-C1-C2	-3.88	92.91	110.38
5	A	1743	PGO	O2-C2-C3	3.56	124.78	109.45
4	A	1747	PGR	O1-C1-C2	-3.52	95.85	114.54
5	A	1743	PGO	C3-C2-C1	3.37	124.65	110.80
3	A	1741	GOL	O2-C2-C3	3.23	122.54	109.18

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	A	1743	PGO	C2

5 of 83 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1733	GOL	O1-C1-C2-O2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	1733	GOL	O1-C1-C2-C3
3	A	1733	GOL	O2-C2-C3-O3
3	A	1734	GOL	C1-C2-C3-O3
3	A	1739	GOL	C1-C2-C3-O3

There are no ring outliers.

28 monomers are involved in 46 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1741	GOL	3	0
3	A	1734	GOL	1	0
4	A	1747	PGR	1	0
4	A	1753	PGR	2	0
4	A	1770	PGR	1	0
4	A	1764	PGR	1	0
4	A	1784	PGR	2	0
4	A	1756	PGR	3	0
4	A	1746	PGR	1	0
4	A	1768	PGR	1	0
3	A	1732	GOL	1	0
4	A	1778	PGR	1	0
6	A	1791	PO4	1	0
4	A	1750	PGR	2	0
3	A	1735	GOL	1	0
4	A	1780	PGR	1	0
4	A	1758	PGR	1	0
4	A	1772	PGR	1	0
3	A	1738	GOL	2	0
3	A	1740	GOL	3	0
4	A	1781	PGR	1	0
4	A	1752	PGR	3	0
4	A	1767	PGR	3	0
4	A	1749	PGR	2	0
4	A	1760	PGR	1	0
5	A	1743	PGO	1	0
4	A	1782	PGR	6	0
4	A	1777	PGR	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	721/751 (96%)	-0.27	10 (1%) 73 78	16, 36, 52, 79	51 (7%)
2	B	2/4 (50%)	0.41	0 100 100	38, 38, 38, 49	0
All	All	723/755 (95%)	-0.27	10 (1%) 73 78	16, 36, 52, 79	51 (7%)

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	192	SER	3.0
1	A	389[A]	THR	2.5
1	A	136	ALA	2.3
1	A	135	VAL	2.3
1	A	397	GLU	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	FC0	B	1	14/15	0.89	0.18	62,84,89,89	0
2	RGL	B	4	11/11	0.99	0.05	28,31,33,36	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands i

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	PGR	A	1789	5/5	0.58	0.35	67,69,73,75	5
4	PGR	A	1773	5/5	0.75	0.28	86,91,93,94	0
4	PGR	A	1785	5/5	0.78	0.28	67,67,70,73	5
4	PGR	A	1763	5/5	0.81	0.24	67,77,80,82	0
4	PGR	A	1779	5/5	0.81	0.22	78,86,92,93	0
4	PGR	A	1767	5/5	0.82	0.23	67,71,76,76	0
4	PGR	A	1790	5/5	0.82	0.23	69,76,77,78	0
4	PGR	A	1760	5/5	0.83	0.23	71,75,81,83	0
3	GOL	A	1735	6/6	0.84	0.20	64,74,78,78	0
4	PGR	A	1750	5/5	0.84	0.24	68,74,75,75	0
4	PGR	A	1783	5/5	0.85	0.20	61,62,66,66	0
3	GOL	A	1739	6/6	0.85	0.26	57,69,71,72	6
4	PGR	A	1784	5/5	0.86	0.21	49,52,55,57	5
4	PGR	A	1781	5/5	0.86	0.18	70,70,72,73	0
4	PGR	A	1770	5/5	0.87	0.25	63,69,70,71	5
4	PGR	A	1787	5/5	0.87	0.21	65,73,78,79	0
4	PGR	A	1759	5/5	0.87	0.24	89,90,92,92	0
4	PGR	A	1768	5/5	0.87	0.21	81,84,85,86	0
4	PGR	A	1748	5/5	0.88	0.21	85,88,91,93	0
4	PGR	A	1766	5/5	0.88	0.20	60,73,75,79	0
3	GOL	A	1733	6/6	0.88	0.20	75,81,82,84	0
4	PGR	A	1774	5/5	0.88	0.17	41,56,67,69	0
4	PGR	A	1782	5/5	0.89	0.21	48,61,67,70	4
4	PGR	A	1769	5/5	0.89	0.22	59,69,70,70	4
4	PGR	A	1786	5/5	0.89	0.24	87,87,89,89	0
6	PO4	A	1793	5/5	0.89	0.20	53,53,56,58	5
3	GOL	A	1736	6/6	0.90	0.20	71,82,85,89	0
4	PGR	A	1753	5/5	0.90	0.17	63,67,69,72	0
4	PGR	A	1754	5/5	0.90	0.23	87,88,89,90	0
4	PGR	A	1778	5/5	0.91	0.18	73,75,78,78	0
4	PGR	A	1761	5/5	0.91	0.17	55,58,64,71	0
4	PGR	A	1758	5/5	0.91	0.17	77,78,80,81	0
4	PGR	A	1788	5/5	0.91	0.21	63,70,73,73	5
4	PGR	A	1765	5/5	0.91	0.15	35,53,58,58	0
3	GOL	A	1737	6/6	0.91	0.17	68,70,73,73	6
4	PGR	A	1751	5/5	0.91	0.17	54,55,58,61	0
4	PGR	A	1780	5/5	0.92	0.22	67,70,71,72	5

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GOL	A	1732	6/6	0.92	0.15	44,46,49,50	0
4	PGR	A	1772	5/5	0.92	0.12	79,82,83,85	0
4	PGR	A	1762	5/5	0.92	0.17	69,70,71,72	0
4	PGR	A	1771	5/5	0.93	0.14	43,46,51,53	5
4	PGR	A	1755	5/5	0.93	0.14	51,59,61,64	0
4	PGR	A	1757	5/5	0.93	0.14	52,59,64,67	0
4	PGR	A	1749	5/5	0.93	0.14	53,59,62,62	0
4	PGR	A	1775	5/5	0.93	0.16	42,53,61,65	0
4	PGR	A	1776	5/5	0.93	0.15	41,52,58,65	0
4	PGR	A	1742	5/5	0.93	0.13	41,41,46,50	0
4	PGR	A	1745	5/5	0.93	0.14	51,55,60,65	0
4	PGR	A	1747	5/5	0.93	0.16	44,57,61,61	0
6	PO4	A	1792	5/5	0.93	0.25	74,75,76,77	5
3	GOL	A	1734	6/6	0.93	0.14	50,67,70,70	0
4	PGR	A	1752	5/5	0.94	0.10	54,60,65,71	0
3	GOL	A	1740	6/6	0.94	0.16	46,48,53,54	6
4	PGR	A	1777	5/5	0.95	0.14	56,59,62,63	0
4	PGR	A	1744	5/5	0.95	0.11	47,48,51,53	0
3	GOL	A	1738	6/6	0.95	0.13	46,49,53,64	5
4	PGR	A	1764	5/5	0.95	0.12	48,54,60,61	0
6	PO4	A	1791	5/5	0.96	0.13	47,50,56,58	5
4	PGR	A	1746	5/5	0.96	0.12	34,43,56,61	0
3	GOL	A	1741	6/6	0.96	0.09	42,48,52,56	6
4	PGR	A	1756	5/5	0.97	0.09	54,58,63,63	0
7	CL	A	1799	1/1	0.98	0.11	66,66,66,66	0
5	PGO	A	1743	5/5	0.99	0.05	30,34,41,43	0
8	NA	A	1795	1/1	0.99	0.11	47,47,47,47	0
7	CL	A	1798	1/1	1.00	0.03	42,42,42,42	0
7	CL	A	1794	1/1	1.00	0.06	45,45,45,45	0
7	CL	A	1796	1/1	1.00	0.15	46,46,46,46	0
8	NA	A	1797	1/1	1.00	0.11	34,34,34,34	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.