



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 15, 2026 – 12:53 AM UTC

PDB ID : 2XEF / pdb_00002xef
Title : Human glutamate carboxypeptidase II in complex with Antibody- Recruiting Molecule ARM-P8
Authors : Zhang, A.X.; Murelli, R.P.; Barinka, C.; Michel, J.; Cocleaza, A.; Jorgensen, W.L.; Lubkowski, J.; Spiegel, D.A.
Deposited on : 2010-05-14
Resolution : 1.59 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	709	5% 86%11%..
2	B	2	50%50%
2	C	2	100%
2	D	2	100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	E	4	25% 50% 25%

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 6687 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GLUTAMATE CARBOXYPEPTIDASE 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	694	Total	C	N	O	S	0	53	0
			5798	3733	960	1081	24			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	42	ARG	-	expression tag	UNP Q04609
A	43	SER	-	expression tag	UNP Q04609
A	593	GLN	LEU	conflict	UNP Q04609

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	C	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	D	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	4	Total	C	N	O	0	0	0
			50	28	2	20			

- Molecule 4 is N-{[(1S)-1-carboxy-5-{4-[25-(2,4-dinitroanilino)-2,5,8,11,14,17,20,23-octaoxa pentacosan-1-yl]-1H-1,2,3-triazol-1-yl}pentyl]carbamoyl}-L-glutamic acid (CCD ID: AR8) (formula: C₃₇H₅₈N₈O₁₉).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	35	1
			128	74	16	38		

- Molecule 5 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Zn	0	0
			2	2		


- Molecule 6 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Ca	0	0
			1	1		

- Molecule 7 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total Cl 1 1	0	0

- Molecule 8 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total C N O 14 8 1 5	0	0
8	A	1	Total C N O 14 8 1 5	0	0
8	A	1	Total C N O 14 8 1 5	0	0


- Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	581	Total O 581 581	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: GLUTAMATE CARBOXYPEPTIDASE 2



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



NAG1
NAG2

- Molecule 3: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E: 

NAG1
NAG2
BMA3
MAN4

4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	101.84Å 130.03Å 158.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.59 30.00 – 1.59	Depositor EDS
% Data completeness (in resolution range)	96.9 (30.00-1.59) 97.3 (30.00-1.59)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.13 (at 1.58Å)	Xtriage
Refinement program	REFMAC 5.4.0057	Depositor
R, R_{free}	0.161 , 0.183 0.187 , 0.207	Depositor DCC
R_{free} test set	2037 reflections (1.48%)	wwPDB-VP
Wilson B-factor (Å ²)	21.0	Xtriage
Anisotropy	0.040	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 56.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6687	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AR8, NAG, ZN, BMA, MAN, CA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.92	1/6104 (0.0%)	0.95	1/8269 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	533	GLY	N-CA	5.27	1.51	1.45

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	55	LYS	N-CA-CB	5.39	119.66	110.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5798	0	5699	80	1
2	B	28	0	25	2	0
2	C	28	0	25	0	0
2	D	28	0	25	0	0
3	E	50	0	43	0	1
4	A	128	0	114	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	2	0	0	0	0
6	A	1	0	0	0	0
7	A	1	0	0	0	0
8	A	42	0	39	1	0
9	A	581	0	0	22	1
All	All	6687	0	5970	86	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (86) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:660[A]:VAL:O	1:A:664[A]:MET:HG2	1.16	1.28
1:A:593[B]:GLN:NE2	1:A:594:PRO:HD2	1.81	0.95
1:A:660[A]:VAL:O	1:A:664[A]:MET:CG	2.12	0.94
1:A:89[B]:ASN:HD21	1:A:378:ARG:HH21	1.00	0.92
1:A:733:TYR:HE2	9:A:2549:HOH:O	1.61	0.83
1:A:89[B]:ASN:ND2	1:A:378:ARG:HH21	1.75	0.83
1:A:593[B]:GLN:NE2	1:A:593[B]:GLN:HA	1.93	0.80
1:A:641:GLU:HG3	9:A:2447:HOH:O	1.84	0.78
1:A:89[B]:ASN:HD21	1:A:378:ARG:NH2	1.81	0.73
1:A:400:ARG:O	1:A:404[B]:THR:HG23	1.89	0.73
1:A:620[B]:GLN:OE1	9:A:2426:HOH:O	2.07	0.71
1:A:131:ILE:HG22	1:A:137:GLU:HG2	1.74	0.70
1:A:593[B]:GLN:HE22	1:A:594:PRO:HD2	1.58	0.66
1:A:612:TYR:CZ	1:A:616:MET:HG3	2.30	0.66
1:A:613[B]:SER:OG	9:A:2420:HOH:O	2.15	0.65
1:A:593[B]:GLN:HE21	1:A:593[B]:GLN:CA	2.11	0.64
1:A:593[B]:GLN:HE21	1:A:594:PRO:HD2	1.59	0.63
1:A:88:GLN:NE2	9:A:2027:HOH:O	2.30	0.63
1:A:593[B]:GLN:NE2	1:A:594:PRO:CD	2.60	0.62
1:A:236:ALA:O	1:A:239[A]:VAL:HG13	1.99	0.62
1:A:733:TYR:CE2	9:A:2549:HOH:O	2.43	0.62
1:A:593[B]:GLN:NE2	1:A:593[B]:GLN:CA	2.52	0.61
1:A:169:MET:HE2	1:A:347:HIS:NE2	2.15	0.61
1:A:603[B]:VAL:CG1	1:A:607:TYR:CE2	2.84	0.60
1:A:90[B]:PHE:CZ	1:A:94:LYS:HD3	2.37	0.59
1:A:688[B]:ARG:NH2	9:A:2495:HOH:O	2.35	0.59
1:A:236:ALA:HB3	1:A:239[A]:VAL:HG11	1.86	0.58
1:A:89[B]:ASN:C	1:A:89[B]:ASN:HD22	2.12	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:591[B]:ILE:HD11	9:A:2074:HOH:O	2.05	0.56
1:A:412:PRO:HA	1:A:589[B]:ASN:OD1	2.06	0.56
1:A:308:LYS:HB2	9:A:2242:HOH:O	2.07	0.55
1:A:232:ALA:O	1:A:545:LYS:HE2	2.06	0.55
1:A:90[B]:PHE:HE2	1:A:94:LYS:HZ3	1.54	0.55
1:A:586[B]:GLU:OE2	1:A:590[B]:SER:OG	2.23	0.55
1:A:517:SER:HB2	1:A:694:PRO:HG3	1.89	0.54
1:A:645[B]:LYS:HE3	9:A:2453:HOH:O	2.07	0.54
1:A:656[B]:SER:O	1:A:658[B]:PRO:HD3	2.08	0.53
1:A:648:GLU:O	1:A:652:ASP:HB3	2.08	0.52
9:A:2033:HOH:O	2:B:2:NAG:H81	2.08	0.52
1:A:603[B]:VAL:HG12	1:A:607:TYR:CE2	2.43	0.52
1:A:603[B]:VAL:HG13	1:A:607:TYR:CZ	2.44	0.52
1:A:603[B]:VAL:CG1	1:A:607:TYR:CZ	2.94	0.50
1:A:586[B]:GLU:CD	1:A:590[B]:SER:OG	2.55	0.50
1:A:242:TYR:OH	9:A:2165:HOH:O	2.18	0.50
4:A:1770[A]:AR8:HBE2	4:A:1770[A]:AR8:HAU2	1.93	0.50
1:A:131:ILE:CG2	1:A:137:GLU:HG2	2.42	0.49
1:A:236:ALA:HB3	1:A:239[A]:VAL:CG1	2.42	0.49
1:A:90[B]:PHE:CE2	1:A:94:LYS:NZ	2.81	0.48
1:A:174[A]:LEU:HG	1:A:202:ILE:HG22	1.95	0.48
1:A:89[B]:ASN:ND2	1:A:89[B]:ASN:C	2.72	0.48
1:A:475:HIS:CE1	1:A:509[B]:MET:CE	2.97	0.47
1:A:367:GLU:OE1	1:A:662[B]:ARG:NH1	2.47	0.47
1:A:676:ILE:HG23	1:A:688[B]:ARG:HB3	1.95	0.47
1:A:88:GLN:CD	9:A:2027:HOH:O	2.58	0.47
1:A:57:ASN:C	1:A:57:ASN:HD22	2.23	0.46
1:A:58[B]:MET:HG3	1:A:585:PHE:CG	2.51	0.46
1:A:593[B]:GLN:HE21	1:A:594:PRO:CD	2.26	0.45
1:A:308:LYS:NZ	9:A:2245:HOH:O	2.48	0.44
1:A:586[B]:GLU:HG3	1:A:590[B]:SER:OG	2.18	0.44
1:A:169:MET:HA	1:A:344:MET:O	2.18	0.44
1:A:347:HIS:CE1	9:A:2066:HOH:O	2.70	0.44
1:A:468:PRO:HG2	1:A:603[A]:VAL:HG21	1.99	0.44
1:A:347:HIS:HE1	9:A:2066:HOH:O	2.02	0.43
1:A:488:PHE:CE1	1:A:496:SER:HA	2.53	0.43
1:A:618:HIS:HE1	9:A:2525:HOH:O	2.02	0.43
1:A:309[B]:MET:HE3	1:A:309[B]:MET:HB3	1.92	0.43
1:A:468:PRO:CG	1:A:603[A]:VAL:HG21	2.49	0.43
1:A:137:GLU:OE2	1:A:345:HIS:CE1	2.72	0.42
1:A:514[A]:LYS:HG3	9:A:2358:HOH:O	2.18	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:325:VAL:HB	1:A:326:PRO:HD2	2.00	0.42
1:A:246:TRP:CD1	8:A:1760:NAG:H83	2.55	0.42
1:A:544:ASN:HB3	1:A:546:PHE:CE2	2.54	0.41
1:A:645[B]:LYS:HA	1:A:645[B]:LYS:HD3	1.94	0.41
1:A:551:LEU:HD22	1:A:556:TYR:HB2	2.02	0.41
1:A:688[B]:ARG:CZ	9:A:2490:HOH:O	2.67	0.41
4:A:1770[B]:AR8:OAC	4:A:1770[B]:AR8:CCC	2.67	0.41
1:A:137:GLU:OE2	1:A:345:HIS:HE1	2.03	0.41
1:A:258:ILE:HD13	1:A:294:VAL:HB	2.03	0.41
1:A:591[B]:ILE:HG13	9:A:2400:HOH:O	2.21	0.41
1:A:617:LYS:NZ	9:A:2425:HOH:O	2.54	0.41
1:A:661[A]:LEU:O	1:A:664[A]:MET:HB2	2.21	0.41
4:A:1770[A]:AR8:OAK	4:A:1770[A]:AR8:NBO	2.47	0.40
9:A:2564:HOH:O	2:B:2:NAG:H83	2.21	0.40
1:A:57:ASN:ND2	1:A:57:ASN:N	2.69	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:A:2469:HOH:O	9:A:2469:HOH:O[2_565]	1.04	1.16
1:A:276[A]:GLU:OE1	3:E:3:BMA:O2[2_565]	1.96	0.24

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	743/709 (105%)	725 (98%)	17 (2%)	1 (0%)	48 27

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	382	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	644/605 (106%)	633 (98%)	11 (2%)	53 30

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	55	LYS
1	A	57	ASN
1	A	239[A]	VAL
1	A	239[B]	VAL
1	A	507	SER
1	A	537	TYR
1	A	545	LYS
1	A	547	SER
1	A	600	TYR
1	A	652	ASP
1	A	719	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	57	ASN
1	A	95	GLN
1	A	97	GLN
1	A	303	GLN
1	A	345	HIS
1	A	618	HIS
1	A	667	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

10 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAG	B	1	2,1	14,14,15	0.48	0	17,19,21	1.36	2 (11%)
2	NAG	B	2	2	14,14,15	0.61	0	17,19,21	1.13	1 (5%)
2	NAG	C	1	2,1	14,14,15	0.54	0	17,19,21	0.97	0
2	NAG	C	2	2	14,14,15	0.41	0	17,19,21	0.80	0
2	NAG	D	1	2,1	14,14,15	0.80	0	17,19,21	1.35	2 (11%)
2	NAG	D	2	2	14,14,15	0.47	0	17,19,21	1.29	1 (5%)
3	NAG	E	1	3,1	14,14,15	0.82	1 (7%)	17,19,21	1.25	3 (17%)
3	NAG	E	2	3	14,14,15	0.83	1 (7%)	17,19,21	0.98	0
3	BMA	E	3	3	11,11,12	0.55	0	15,15,17	1.03	1 (6%)
3	MAN	E	4	3	11,11,12	0.57	0	15,15,17	0.96	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	2/6/23/26	0/1/1/1
2	NAG	C	1	2,1	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	2	2	-	2/6/23/26	0/1/1/1
2	NAG	D	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	D	2	2	-	2/6/23/26	0/1/1/1
3	NAG	E	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	E	2	3	-	3/6/23/26	0/1/1/1
3	BMA	E	3	3	-	0/2/19/22	0/1/1/1
3	MAN	E	4	3	-	0/2/19/22	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	2	NAG	O5-C1	-2.75	1.39	1.43
3	E	1	NAG	C1-C2	2.18	1.55	1.52

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	NAG	C2-N2-C7	3.20	127.19	122.90
2	B	1	NAG	C2-N2-C7	-3.09	118.76	122.90
2	D	1	NAG	O5-C1-C2	-2.84	106.89	111.29
3	E	1	NAG	O5-C1-C2	-2.82	106.93	111.29
2	B	1	NAG	O5-C1-C2	-2.71	107.09	111.29
3	E	3	BMA	O3-C3-C2	-2.62	104.71	110.05
3	E	1	NAG	O4-C4-C5	-2.59	102.94	109.32
2	D	2	NAG	C8-C7-N2	2.46	120.20	116.12
2	D	1	NAG	O5-C5-C4	-2.42	104.95	110.83
3	E	1	NAG	C2-N2-C7	-2.23	119.91	122.90

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	2	NAG	C8-C7-N2-C2
2	B	2	NAG	O7-C7-N2-C2
2	D	1	NAG	C8-C7-N2-C2
2	D	1	NAG	O7-C7-N2-C2
2	D	2	NAG	C8-C7-N2-C2
2	D	2	NAG	O7-C7-N2-C2
3	E	2	NAG	C8-C7-N2-C2
3	E	2	NAG	O7-C7-N2-C2

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	C	2	NAG	C4-C5-C6-O6
2	C	2	NAG	O5-C5-C6-O6
3	E	2	NAG	C4-C5-C6-O6


There are no ring outliers.


2 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	2	NAG	2	0
3	E	3	BMA	0	1

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 4 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	AR8	A	1770[B]	-	65,65,65	2.94	13 (20%)	73,79,79	1.48	11 (15%)
8	NAG	A	1757	1	14,14,15	0.63	0	17,19,21	1.28	2 (11%)
8	NAG	A	1759	1	14,14,15	0.37	0	17,19,21	2.23	6 (35%)
4	AR8	A	1770[A]	5	65,65,65	2.66	8 (12%)	73,79,79	1.51	10 (13%)
8	NAG	A	1760	1	14,14,15	0.62	0	17,19,21	1.54	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	AR8	A	1770[B]	-	-	12/61/65/65	0/2/2/2
8	NAG	A	1757	1	-	0/6/23/26	0/1/1/1
8	NAG	A	1759	1	-	4/6/23/26	0/1/1/1
4	AR8	A	1770[A]	5	-	18/61/65/65	0/2/2/2
8	NAG	A	1760	1	-	0/6/23/26	0/1/1/1

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1770[B]	AR8	OAE-NCK	13.42	1.45	1.22
4	A	1770[A]	AR8	OAE-NCK	12.42	1.44	1.22
4	A	1770[B]	AR8	OAF-NCL	11.79	1.43	1.22
4	A	1770[A]	AR8	OAF-NCL	10.93	1.41	1.22
4	A	1770[B]	AR8	CCG-NCL	-6.58	1.33	1.45
4	A	1770[B]	AR8	NCJ-NBN	-6.47	1.24	1.34
4	A	1770[A]	AR8	NCJ-NBN	-6.25	1.24	1.34
4	A	1770[A]	AR8	CCG-NCL	-5.86	1.35	1.45
4	A	1770[B]	AR8	CCE-NCK	-4.83	1.33	1.45
4	A	1770[B]	AR8	NBM-NBN	-4.68	1.24	1.32
4	A	1770[B]	AR8	CBL-CCD	-4.37	1.40	1.49
4	A	1770[A]	AR8	NBM-NBN	-4.34	1.24	1.32
4	A	1770[A]	AR8	CCE-NCK	-4.24	1.35	1.45
4	A	1770[A]	AR8	CBL-CCD	-4.12	1.41	1.49
4	A	1770[B]	AR8	CAR-NBO	3.92	1.53	1.45
4	A	1770[B]	AR8	OBR-CAS	3.07	1.55	1.42
4	A	1770[B]	AR8	OBW-CBC	2.99	1.55	1.42
4	A	1770[B]	AR8	OBS-CAU	2.16	1.51	1.42
4	A	1770[B]	AR8	CAO-CCD	-2.12	1.33	1.36
4	A	1770[B]	AR8	CBB-CBC	2.09	1.59	1.49
4	A	1770[A]	AR8	OBT-CAX	2.08	1.51	1.42

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	A	1759	NAG	C1-O5-C5	7.04	121.62	112.19
4	A	1770[B]	AR8	OBV-CBL-CCD	5.27	123.17	110.08
4	A	1770[A]	AR8	OBV-CBL-CCD	4.93	122.32	110.08
4	A	1770[A]	AR8	CB-CA-N	-4.79	101.43	110.91
4	A	1770[B]	AR8	CB-CA-N	-4.68	101.64	110.91
4	A	1770[A]	AR8	NCJ-NBN-NBM	3.81	110.44	107.02
4	A	1770[B]	AR8	NCJ-NBN-NBM	3.68	110.33	107.02
8	A	1757	NAG	C1-C2-N2	-3.21	105.38	110.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	A	1760	NAG	C1-O5-C5	3.15	116.41	112.19
4	A	1770[B]	AR8	CAR-NBO-CCF	-2.96	116.16	123.35
4	A	1770[B]	AR8	CCB-CCI-NBQ	-2.75	104.18	110.57
4	A	1770[A]	AR8	CAQ-CAP-CB	-2.68	103.51	113.62
8	A	1760	NAG	C2-N2-C7	-2.68	119.31	122.90
4	A	1770[A]	AR8	CBL-CCD-NBM	2.63	125.84	121.49
8	A	1760	NAG	O3-C3-C4	-2.61	104.23	110.38
8	A	1759	NAG	C6-C5-C4	-2.59	106.67	113.02
4	A	1770[A]	AR8	CAN-CCG-CCF	-2.57	119.20	121.55
8	A	1760	NAG	O5-C1-C2	-2.51	107.40	111.29
4	A	1770[A]	AR8	CCG-CCF-NBO	-2.47	119.15	123.38
8	A	1759	NAG	C4-C3-C2	-2.44	107.44	111.02
8	A	1759	NAG	C8-C7-N2	2.41	120.12	116.12
4	A	1770[A]	AR8	CBL-CCD-CAO	-2.39	126.10	130.22
4	A	1770[B]	AR8	CAS-OBR-CAT	-2.35	102.97	113.26
4	A	1770[B]	AR8	CBL-CCD-NBM	2.18	125.11	121.49
8	A	1759	NAG	C1-C2-N2	2.17	113.84	110.43
8	A	1759	NAG	C2-N2-C7	2.11	125.73	122.90
4	A	1770[B]	AR8	CAM-CCF-NBO	-2.10	118.28	121.75
4	A	1770[B]	AR8	OAE-NCK-CCE	2.08	121.68	118.82
4	A	1770[B]	AR8	CBK-NCJ-CAO	-2.05	125.01	128.83
4	A	1770[A]	AR8	OAC-CCB-CCI	2.04	120.43	113.51
4	A	1770[B]	AR8	CAQ-CAP-CB	-2.04	105.93	113.62
8	A	1757	NAG	O3-C3-C2	-2.02	105.21	109.40
4	A	1770[A]	AR8	CBK-NCJ-CAO	-2.01	125.07	128.83

There are no chirality outliers.

All (34) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1770[A]	AR8	CCD-CBL-OBY-CBG
4	A	1770[B]	AR8	CCD-CBL-OBY-CBG
4	A	1770[A]	AR8	CAW-CAV-OBS-CAU
4	A	1770[B]	AR8	OBT-CAX-CAY-OBV
4	A	1770[B]	AR8	OBR-CAT-CAU-OBS
4	A	1770[B]	AR8	NBO-CAR-CAS-OBR
4	A	1770[B]	AR8	OBX-CBF-CBG-OBV
4	A	1770[A]	AR8	OBS-CAV-CAW-OBT
8	A	1759	NAG	C8-C7-N2-C2
8	A	1759	NAG	O7-C7-N2-C2
4	A	1770[A]	AR8	OBX-CBF-CBG-OBV
4	A	1770[B]	AR8	CAS-CAR-NBO-CCF

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	A	1770[A]	AR8	OBT-CAX-CAY-OBV
4	A	1770[A]	AR8	NBO-CAR-CAS-OBV
4	A	1770[B]	AR8	OBV-CAZ-CBA-OBV
4	A	1770[A]	AR8	OBV-CBD-CBE-OBX
8	A	1759	NAG	C4-C5-C6-O6
4	A	1770[A]	AR8	CAY-CAX-OBV-CAW
4	A	1770[A]	AR8	CBD-CBE-OBX-CBF
4	A	1770[A]	AR8	CAU-CAT-OBV-CAS
4	A	1770[A]	AR8	CAT-CAU-OBS-CAV
4	A	1770[B]	AR8	CBB-CBC-OBV-CBD
4	A	1770[A]	AR8	CBB-CBC-OBV-CBD
4	A	1770[A]	AR8	CAZ-CBA-OBV-CBB
4	A	1770[A]	AR8	CBE-CBD-OBV-CBC
4	A	1770[B]	AR8	OBV-CBD-CBE-OBX
4	A	1770[A]	AR8	CBF-CBG-OBV-CBL
4	A	1770[A]	AR8	CAR-CAS-OBV-CAT
8	A	1759	NAG	O5-C5-C6-O6
4	A	1770[B]	AR8	CBC-CBB-OBV-CBA
4	A	1770[B]	AR8	CAU-CAT-OBV-CAS
4	A	1770[A]	AR8	CBC-CBB-OBV-CBA
4	A	1770[B]	AR8	OBS-CAV-CAW-OBV
4	A	1770[A]	AR8	C-CA-CB-CAP


There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1770[B]	AR8	1	0
4	A	1770[A]	AR8	3	0
8	A	1760	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	694/709 (97%)	0.13	32 (4%) 37 39	8, 23, 42, 62	54 (7%)

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	656[A]	SER	7.9
1	A	488	PHE	4.5
1	A	506	PHE	4.3
1	A	546	PHE	4.0
1	A	541	TRP	3.9
1	A	545	LYS	3.8
1	A	652	ASP	3.5
1	A	335	GLY	3.5
1	A	135	GLY	3.4
1	A	153	ASN	3.4
1	A	653	PHE	3.3
1	A	152	GLU	3.2
1	A	616	MET	3.1
1	A	154	VAL	2.9
1	A	196	CYS	2.9
1	A	55	LYS	2.8
1	A	547	SER	2.7
1	A	544	ASN	2.6
1	A	169	MET	2.6
1	A	155	SER	2.6
1	A	134	ASP	2.5
1	A	597	CYS	2.4
1	A	507	SER	2.4
1	A	192	MET	2.4
1	A	486	GLU	2.4
1	A	336	ASN	2.3
1	A	189	GLU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	332	GLY	2.3
1	A	719	VAL	2.2
1	A	279[A]	TYR	2.2
1	A	334	THR	2.0
1	A	504	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	B	2	14/15	0.75	0.15	38,44,48,51	0
2	NAG	C	2	14/15	0.82	0.12	51,55,59,60	0
2	NAG	D	2	14/15	0.85	0.12	31,36,42,43	0
3	NAG	E	2	14/15	0.85	0.12	32,35,45,47	0
3	MAN	E	4	11/12	0.86	0.12	38,42,45,47	0
2	NAG	C	1	14/15	0.88	0.11	35,39,43,48	0
3	BMA	E	3	11/12	0.91	0.09	31,35,36,38	0
2	NAG	D	1	14/15	0.91	0.10	24,27,34,40	0
2	NAG	B	1	14/15	0.92	0.09	28,35,40,43	0
3	NAG	E	1	14/15	0.93	0.09	19,26,31,40	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain B:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain C:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
8	NAG	A	1759	14/15	0.62	0.18	54,61,65,66	0
8	NAG	A	1757	14/15	0.76	0.15	39,47,51,51	0
8	NAG	A	1760	14/15	0.83	0.13	28,43,47,48	0
4	AR8	A	1770[B]	64/64	0.84	0.26	18,56,74,74	64
4	AR8	A	1770[A]	64/64	0.84	0.26	16,66,75,76	64
7	CL	A	1754	1/1	0.99	0.25	6,6,6,6	0
5	ZN	A	1751	1/1	0.99	0.20	6,6,6,6	0
5	ZN	A	1752	1/1	1.00	0.19	5,5,5,5	0
6	CA	A	1753	1/1	1.00	0.15	4,4,4,4	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different

orientation to approximate a three-dimensional view.




Electron density around AR8 A 1770 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around AR8 A 1770 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.