



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 01:05 AM UTC

PDB ID : 4XEB / pdb_00004xeb
Title : The structure of P. funiculosum Cel7A
Authors : Alahuhta, P.M.; Lunin, V.V.
Deposited on : 2014-12-23
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	436	
2	B	2	
3	C	6	

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 4316 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glucanase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	430	Total	C	N	O	S	0	19	0
			3319	2040	556	698	25			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	348	SER	ASN	conflict	UNP F1CYZ0

- Molecule 2 is an oligosaccharide called beta-D-glucopyranose-(1-4)-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	B	2	Total	C	O	0	0	0
			23	12	11			


- Molecule 3 is an oligosaccharide called beta-D-glucopyranose-(1-4)-beta-D-glucopyranose-(1-4)-beta-D-glucopyranose-(1-4)-beta-D-glucopyranose-(1-4)-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
3	C	6	Total	C	O	0	1	0
			70	38	32			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula:

C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	3	Total	K	0	0
			3	3		

- Molecule 6 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Na	0	0
			1	1		

- Molecule 7 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Cl	0	0
			1	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	820	Total 857	O 857	0	37


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glucanase



- Molecule 2: beta-D-glucopyranose-(1-4)-beta-D-glucopyranose



- Molecule 3: beta-D-glucopyranose-(1-4)-beta-D-glucopyranose-(1-4)-beta-D-glucopyranose-(1-4)-beta-D-glucopyranose-(1-4)-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	70.97Å 61.84Å 85.85Å 90.00° 98.77° 90.00°	Depositor
Resolution (Å)	50.00 – 1.70 50.00 – 1.70	Depositor EDS
% Data completeness (in resolution range)	95.5 (50.00-1.70) 95.5 (50.00-1.70)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.41 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.138 , 0.203 (Not available) , 0.195	Depositor DCC
R_{free} test set	1188 reflections (2.93%)	wwPDB-VP
Wilson B-factor (Å ²)	10.4	Xtriage
Anisotropy	0.138	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 59.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4316	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PCA, NAG, K, CL, BGC, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.11	3/3386 (0.1%)	1.04	1/4625 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	243	THR	CA-C	8.17	1.64	1.52
1	A	190	THR	C-O	-5.57	1.20	1.25
1	A	314	ILE	CA-C	5.02	1.58	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	132	HIS	N-CA-C	5.64	118.94	112.57

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3319	0	3031	38	0
2	B	23	0	21	0	0
3	C	70	0	56	0	0
4	A	42	0	39	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	3	0	0	0	0
6	A	1	0	0	0	0
7	A	1	0	0	0	0
8	A	857	0	0	17	0
All	All	4316	0	3147	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:69:GLN:HE21	1:A:157[B]:ASN:HD21	1.20	0.86
1:A:336:ALA:O	1:A:340:THR:HG23	1.86	0.76
1:A:41[B]:VAL:HG22	1:A:50:CYS:HB2	1.68	0.74
1:A:46[A]:THR:HG23	8:A:1145:HOH:O	1.85	0.73
1:A:69:GLN:HG3	1:A:157[B]:ASN:CG	2.20	0.66
1:A:45:ASN:OD1	4:A:503:NAG:C2	2.44	0.65
1:A:69:GLN:NE2	1:A:157[B]:ASN:HD21	1.96	0.61
1:A:45:ASN:OD1	4:A:503:NAG:H2	2.01	0.61
1:A:342:GLY:HA2	8:A:675[A]:HOH:O	2.01	0.60
1:A:46[A]:THR:CG2	8:A:1145:HOH:O	2.48	0.60
1:A:187[B]:GLU:HB2	8:A:917[B]:HOH:O	2.00	0.60
1:A:202:ASN:OD1	8:A:610:HOH:O	2.17	0.57
1:A:183:GLN:NE2	8:A:613:HOH:O	2.32	0.55
1:A:224:PRO:HD2	1:A:258:CYS:O	2.09	0.52
1:A:69:GLN:HG2	1:A:157[B]:ASN:OD1	2.14	0.48
1:A:132:HIS:CG	1:A:418[A]:TYR:CE2	3.01	0.47
1:A:69:GLN:HG3	1:A:157[B]:ASN:ND2	2.30	0.47
1:A:200:ILE:HD11	1:A:244:TYR:CD1	2.50	0.47
1:A:244:TYR:CE2	8:A:1177[A]:HOH:O	2.56	0.47
1:A:344:THR:HA	8:A:1115:HOH:O	2.15	0.46
1:A:51:TYR:CD1	1:A:178[A]:LYS:HE2	2.51	0.46
1:A:69:GLN:CG	1:A:157[B]:ASN:OD1	2.64	0.46
1:A:320[A]:LYS:NZ	8:A:611:HOH:O	2.29	0.45
1:A:336:ALA:O	1:A:340:THR:CG2	2.60	0.45
1:A:413[B]:GLN:HG2	8:A:976:HOH:O	2.17	0.45
1:A:275:LYS:NZ	8:A:643:HOH:O	2.50	0.44
1:A:41[A]:VAL:HG23	8:A:637:HOH:O	2.16	0.44
1:A:244:TYR:N	8:A:645:HOH:O	2.50	0.44
1:A:194:ASN:HB2	8:A:1209:HOH:O	2.17	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:385:TYR:HA	1:A:386:PRO:C	2.44	0.42
1:A:45:ASN:OD1	4:A:503:NAG:N2	2.52	0.42
1:A:218[B]:ILE:HD12	1:A:399:THR:OG1	2.20	0.42
1:A:178[B]:LYS:HE2	8:A:776:HOH:O	2.19	0.42
1:A:46[B]:THR:HG21	8:A:973:HOH:O	2.19	0.42
1:A:157[A]:ASN:HB2	1:A:182:GLY:O	2.20	0.41
1:A:178[B]:LYS:NZ	8:A:637:HOH:O	2.48	0.41
1:A:302:GLU:HA	1:A:329:ILE:O	2.22	0.40
1:A:69:GLN:CG	1:A:157[B]:ASN:CG	2.93	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	445/436 (102%)	436 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	367/350 (105%)	359 (98%)	8 (2%)	45	29

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	41[A]	VAL
1	A	41[B]	VAL
1	A	74	ASP
1	A	137	LEU
1	A	222	LEU
1	A	243	THR
1	A	340	THR
1	A	376	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	69	GLN
1	A	115	GLN
1	A	132	HIS
1	A	288	GLN
1	A	377	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	PCA	A	1	1	7,8,9	1.20	0	9,10,12	1.01	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PCA	A	1	1	-	0/0/11/13	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates ⓘ

9 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	BGC	B	1	2	12,12,12	0.78	0	17,17,17	1.13	1 (5%)
2	BGC	B	2	2	11,11,12	0.73	0	15,15,17	1.03	1 (6%)
3	BGC	C	1[A]	-	12,12,12	0.75	0	17,17,17	1.01	1 (5%)
3	BGC	C	1[B]	-	12,12,12	0.75	0	17,17,17	1.07	1 (5%)
3	BGC	C	2	3	11,11,12	0.78	0	15,15,17	1.41	3 (20%)
3	BGC	C	3	3	11,11,12	0.98	1 (9%)	15,15,17	0.90	0
3	BGC	C	4	3	11,11,12	1.14	0	15,15,17	0.82	0
3	BGC	C	5	3	11,11,12	0.98	1 (9%)	15,15,17	1.08	1 (6%)
3	BGC	C	6	3	11,11,12	1.02	1 (9%)	15,15,17	1.11	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	B	1	2	-	0/2/22/22	0/1/1/1
2	BGC	B	2	2	-	0/2/19/22	0/1/1/1
3	BGC	C	1[A]	-	-	0/2/22/22	0/1/1/1
3	BGC	C	1[B]	-	-	0/2/22/22	0/1/1/1
3	BGC	C	2	3	-	0/2/19/22	0/1/1/1
3	BGC	C	3	3	-	0/2/19/22	0/1/1/1
3	BGC	C	4	3	-	0/2/19/22	0/1/1/1
3	BGC	C	5	3	-	0/2/19/22	0/1/1/1
3	BGC	C	6	3	-	0/2/19/22	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	6	BGC	O5-C1	-2.52	1.39	1.43
3	C	3	BGC	O5-C1	-2.36	1.39	1.43
3	C	5	BGC	O4-C4	2.22	1.48	1.43

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	2	BGC	C1-C2-C3	3.03	114.06	109.64
3	C	5	BGC	O6-C6-C5	-2.87	101.57	111.33
3	C	2	BGC	C1-O5-C5	-2.81	108.42	112.19
3	C	1[A]	BGC	O5-C1-C2	2.51	114.70	110.30
3	C	1[B]	BGC	O5-C1-C2	2.51	114.70	110.30
2	B	1	BGC	O5-C1-C2	-2.43	106.03	110.30
3	C	6	BGC	C1-O5-C5	2.42	115.44	112.19
3	C	2	BGC	C3-C4-C5	-2.31	106.05	110.23
2	B	2	BGC	O3-C3-C2	2.06	114.26	110.05


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 5 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul

statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	NAG	A	503	1	14,14,15	1.14	2 (14%)	17,19,21	3.03	6 (35%)
4	NAG	A	501	1	14,14,15	0.93	1 (7%)	17,19,21	1.93	6 (35%)
4	NAG	A	502	1	14,14,15	0.77	0	17,19,21	1.74	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	503	1	-	2/6/23/26	0/1/1/1
4	NAG	A	501	1	-	1/6/23/26	0/1/1/1
4	NAG	A	502	1	-	1/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	503	NAG	C1-C2	3.14	1.56	1.52
4	A	501	NAG	O5-C1	-2.41	1.39	1.43
4	A	503	NAG	O5-C5	-2.14	1.39	1.43

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	503	NAG	C1-O5-C5	6.94	121.48	112.19
4	A	503	NAG	O5-C1-C2	-6.94	100.56	111.29
4	A	502	NAG	O5-C1-C2	-4.91	103.70	111.29
4	A	503	NAG	C8-C7-N2	4.54	123.65	116.12
4	A	501	NAG	C2-N2-C7	-4.29	117.15	122.90
4	A	503	NAG	C2-N2-C7	4.00	128.26	122.90
4	A	501	NAG	O7-C7-C8	3.02	127.43	122.05
4	A	502	NAG	C4-C3-C2	-2.86	106.83	111.02
4	A	501	NAG	C1-O5-C5	2.83	115.98	112.19
4	A	503	NAG	O5-C5-C6	-2.75	102.31	107.66

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	503	NAG	O7-C7-C8	-2.58	117.46	122.05
4	A	501	NAG	O3-C3-C2	-2.36	104.50	109.40
4	A	501	NAG	C4-C3-C2	-2.34	107.59	111.02
4	A	501	NAG	O7-C7-N2	-2.16	118.17	121.98
4	A	502	NAG	O4-C4-C5	-2.12	104.09	109.32
4	A	502	NAG	C2-N2-C7	-2.11	120.07	122.90

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	502	NAG	C1-C2-N2-C7
4	A	503	NAG	C8-C7-N2-C2
4	A	503	NAG	O7-C7-N2-C2
4	A	501	NAG	C1-C2-N2-C7

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	503	NAG	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	429/436 (98%)	-0.51	3 (0%) 84 86	4, 10, 21, 38	20 (4%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	244	TYR	3.5
1	A	193	ALA	3.4
1	A	243	THR	2.5

6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	PCA	A	1	8/9	0.97	0.05	7,8,8,8	0

6.3 Carbohydrates

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	BGC	C	6	11/12	0.94	0.07	15,20,24,27	0
3	BGC	C	5	11/12	0.95	0.06	9,11,13,14	0
2	BGC	B	1	12/12	0.96	0.05	11,12,17,18	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	BGC	C	1[B]	12/12	0.98	0.04	7,7,8,9	3
3	BGC	C	3	11/12	0.98	0.04	6,6,7,8	0
3	BGC	C	4	11/12	0.98	0.04	7,7,9,9	0
2	BGC	B	2	11/12	0.98	0.04	10,12,14,14	0
3	BGC	C	1[A]	12/12	0.98	0.04	7,7,8,8	3
3	BGC	C	2	11/12	0.99	0.03	6,7,8,8	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	A	503	14/15	0.62	0.18	37,44,47,48	14
4	NAG	A	502	14/15	0.65	0.18	28,32,36,36	14
4	NAG	A	501	14/15	0.82	0.12	10,12,12,12	14
6	NA	A	509	1/1	0.84	0.16	50,50,50,50	0
5	K	A	508	1/1	0.97	0.08	32,32,32,32	0
5	K	A	507	1/1	0.99	0.06	19,19,19,19	0
7	CL	A	510	1/1	0.99	0.12	25,25,25,25	0
5	K	A	506	1/1	1.00	0.04	14,14,14,14	0

6.5 Other polymers [i](#)

There are no such residues in this entry.