



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:22 PM UTC

PDB ID : 7XEN / pdb_00007xen
Title : SufS with L-penicillamine
Authors : Nakamura, R.; Fujishiro, T.
Deposited on : 2022-03-31
Resolution : 2.47 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.47 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7589 (2.50-2.46)
Clashscore	190562	8295 (2.50-2.46)
Ramachandran outliers	187476	8164 (2.50-2.46)
Sidechain outliers	187428	8166 (2.50-2.46)
RSRZ outliers	180081	7593 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	419	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3310 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Cysteine desulfurase SufS.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	392	3052	1938	512	589	13	0	0	0

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	MET	-	initiating methionine	UNP O32164
A	-1	GLY	-	expression tag	UNP O32164
A	0	HIS	-	expression tag	UNP O32164
A	407	VAL	-	expression tag	UNP O32164
A	408	ASP	-	expression tag	UNP O32164
A	409	LEU	-	expression tag	UNP O32164
A	410	GLU	-	expression tag	UNP O32164
A	411	HIS	-	expression tag	UNP O32164
A	412	HIS	-	expression tag	UNP O32164
A	413	HIS	-	expression tag	UNP O32164
A	414	HIS	-	expression tag	UNP O32164
A	415	HIS	-	expression tag	UNP O32164
A	416	HIS	-	expression tag	UNP O32164

- Molecule 2 is (2 {R})-3-methyl-2-[({E})-[2-methyl-3-oxidanyl-5-(phosphonooxymethyl)pyridin-4-yl]methylideneamino]-3-sulfanyl-butanoic acid (CCD ID: 9ZN) (formula: C₁₃H₁₉N₂O₇PS) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
2	A	1	Total	C	N	O	P	S	0	0
			24	13	2	7	1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	234	Total	O	0	0
			234	234		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cysteine desulfurase SufS



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	94.20Å 94.20Å 133.20Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.10 – 2.47 47.10 – 2.47	Depositor EDS
% Data completeness (in resolution range)	98.3 (47.10-2.47) 98.3 (47.10-2.47)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.66 (at 2.48Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.161 , 0.211 0.162 , 0.211	Depositor DCC
R_{free} test set	1235 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	80.1	Xtriage
Anisotropy	0.044	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 63.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.016 for -h,-k,l	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	3310	wwPDB-VP
Average B, all atoms (Å ²)	85.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 9ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.84	2/3114 (0.1%)	1.58	39/4228 (0.9%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	170	HIS	CE1-NE2	5.92	1.38	1.32
1	A	14	HIS	CE1-NE2	5.12	1.37	1.32

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	229	THR	CB-CA-C	-9.24	91.46	111.78
1	A	31	SER	CA-C-N	-8.11	106.69	122.02
1	A	31	SER	C-N-CA	-8.11	106.69	122.02
1	A	18	ASN	CB-CA-C	-7.78	101.86	111.82
1	A	301	ARG	CB-CA-C	7.64	123.83	110.85
1	A	45	TYR	CA-C-N	7.44	133.24	122.23
1	A	45	TYR	C-N-CA	7.44	133.24	122.23
1	A	30	THR	CB-CA-C	7.14	120.67	110.24
1	A	386	GLU	CB-CG-CD	7.04	124.57	112.60
1	A	47	GLN	CB-CA-C	6.77	120.85	109.48
1	A	35	ARG	CB-CA-C	6.70	122.07	110.68

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	65	ASP	CA-C-N	6.60	127.26	119.94
1	A	65	ASP	C-N-CA	6.60	127.26	119.94
1	A	340	HIS	CA-CB-CG	-6.50	107.30	113.80
1	A	160	THR	CA-CB-OG1	-6.50	99.85	109.60
1	A	47	GLN	CA-C-N	6.39	130.60	122.16
1	A	47	GLN	C-N-CA	6.39	130.60	122.16
1	A	49	ASN	CA-CB-CG	6.39	118.99	112.60
1	A	177	THR	CB-CA-C	-6.36	99.99	109.90
1	A	389	ASP	CA-CB-CG	-5.94	106.66	112.60
1	A	349	ASP	CB-CA-C	5.91	120.17	110.88
1	A	371	VAL	CA-C-N	-5.89	112.07	122.26
1	A	371	VAL	C-N-CA	-5.89	112.07	122.26
1	A	211	GLN	CB-CA-C	5.82	120.57	110.79
1	A	186	LYS	CB-CA-C	5.80	120.09	110.81
1	A	397	LYS	CB-CA-C	5.70	121.51	110.46
1	A	230	GLY	CA-C-N	-5.68	114.51	122.75
1	A	230	GLY	C-N-CA	-5.68	114.51	122.75
1	A	160	THR	CB-CA-C	-5.68	100.42	109.80
1	A	371	VAL	O-C-N	-5.66	118.04	122.97
1	A	272	GLU	CB-CG-CD	5.57	122.07	112.60
1	A	32	GLN	CB-CA-C	5.31	123.46	111.78
1	A	298	GLU	CB-CA-C	5.30	119.59	110.79
1	A	32	GLN	CB-CG-CD	5.26	121.54	112.60
1	A	251	GLY	CA-C-N	-5.19	113.47	122.14
1	A	251	GLY	C-N-CA	-5.19	113.47	122.14
1	A	65	ASP	CA-CB-CG	5.16	117.76	112.60
1	A	229	THR	N-CA-CB	5.14	118.36	110.49
1	A	316	GLN	N-CA-C	-5.10	106.57	112.89

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	274	GLY	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3052	0	3009	64	0
2	A	24	0	0	2	0
3	A	234	0	0	7	0
All	All	3310	0	3009	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:249:PHE:HA	1:A:253:MET:HE2	1.37	1.05
1:A:250:GLY:H	1:A:253:MET:CE	1.70	1.04
1:A:359:HIS:HA	1:A:366:MET:HE2	1.40	1.03
1:A:250:GLY:H	1:A:253:MET:HE2	1.37	0.88
1:A:366:MET:HE3	1:A:373:ALA:HA	1.60	0.84
1:A:249:PHE:CA	1:A:253:MET:HE2	2.08	0.83
1:A:250:GLY:H	1:A:253:MET:HE3	1.45	0.79
1:A:138:THR:HG23	3:A:687:HOH:O	1.82	0.79
1:A:121:HIS:NE2	2:A:501:9ZN:SG	2.55	0.78
1:A:250:GLY:N	1:A:253:MET:HE2	1.99	0.76
1:A:386:GLU:HB3	3:A:804:HOH:O	1.88	0.72
1:A:366:MET:CE	1:A:373:ALA:HA	2.20	0.70
2:A:501:9ZN:SG	2:A:501:9ZN:CAO	2.79	0.70
1:A:249:PHE:HA	1:A:253:MET:CE	2.19	0.69
1:A:146:GLU:HA	1:A:146:GLU:OE1	1.92	0.69
1:A:250:GLY:N	1:A:253:MET:CE	2.50	0.68
1:A:337:ASP:O	1:A:338:ASP:HB2	1.93	0.68
1:A:366:MET:HE3	1:A:373:ALA:CA	2.25	0.65
1:A:180:PRO:O	1:A:184:MET:HG3	1.97	0.64
1:A:401:TYR:C	1:A:401:TYR:HD1	2.06	0.64
1:A:183:GLU:HG2	3:A:758:HOH:O	1.97	0.63
1:A:12:ILE:HG21	1:A:32:GLN:HG3	1.80	0.63
1:A:1:MET:N	3:A:601:HOH:O	2.32	0.61
1:A:401:TYR:HD1	1:A:401:TYR:O	1.83	0.61
1:A:366:MET:HE1	1:A:374:THR:HG23	1.84	0.60
1:A:181:ILE:HD11	1:A:213:LEU:HD22	1.85	0.59
1:A:401:TYR:C	1:A:401:TYR:CD1	2.81	0.59
1:A:48:TYR:N	1:A:48:TYR:CD1	2.71	0.58
1:A:381:LEU:H	1:A:381:LEU:HD23	1.69	0.58
1:A:364:PRO:HB2	3:A:690:HOH:O	2.03	0.58
1:A:222:SER:HB2	1:A:284:LEU:HD13	1.84	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:144:LEU:HD13	1:A:148:GLY:HA2	1.87	0.56
1:A:119:MET:HE2	1:A:144:LEU:HD21	1.88	0.56
1:A:112:ASP:O	1:A:137:ALA:HB1	2.07	0.55
1:A:383:ASN:HA	1:A:387:GLU:OE1	2.08	0.53
1:A:348:LEU:HB3	1:A:353:ILE:HB	1.93	0.50
1:A:223:HIS:HD2	3:A:643:HOH:O	1.94	0.49
1:A:181:ILE:CD1	1:A:213:LEU:HD22	2.41	0.49
1:A:223:HIS:HB3	1:A:229:THR:HA	1.96	0.48
1:A:249:PHE:C	1:A:253:MET:HE2	2.39	0.48
1:A:41:LEU:O	1:A:45:TYR:CD2	2.66	0.47
1:A:223:HIS:H	1:A:223:HIS:CD2	2.32	0.47
1:A:117:THR:OG1	1:A:119:MET:HG2	2.16	0.46
1:A:9:GLN:HG3	3:A:660:HOH:O	2.14	0.46
1:A:155:VAL:HG23	1:A:187:ILE:HG21	1.99	0.45
1:A:225:MET:O	1:A:226:CYS:HB2	2.17	0.44
1:A:389:ASP:HA	1:A:392:VAL:HG22	1.99	0.44
1:A:44:TYR:O	1:A:45:TYR:CD1	2.71	0.44
1:A:159:VAL:HG13	1:A:163:THR:CB	2.48	0.44
1:A:160:THR:C	1:A:162:ASN:H	2.26	0.43
1:A:1:MET:HB2	1:A:298:GLU:HG3	2.00	0.43
1:A:381:LEU:HD23	1:A:381:LEU:N	2.32	0.42
1:A:17:VAL:O	1:A:18:ASN:CB	2.68	0.42
1:A:100:ALA:O	1:A:104:ALA:HB3	2.19	0.42
1:A:163:THR:O	1:A:193:ALA:HB2	2.20	0.42
1:A:237:LYS:O	1:A:238:LYS:C	2.59	0.42
1:A:238:LYS:NZ	1:A:242:GLU:OE2	2.48	0.42
1:A:381:LEU:H	1:A:381:LEU:CD2	2.33	0.42
1:A:155:VAL:O	1:A:159:VAL:HG23	2.21	0.41
1:A:395:LEU:HA	1:A:395:LEU:HD23	1.79	0.41
1:A:17:VAL:O	1:A:18:ASN:HB2	2.19	0.41
1:A:318:ASP:O	1:A:399:LYS:HE2	2.20	0.41
1:A:160:THR:HG22	1:A:162:ASN:H	1.86	0.41
1:A:1:MET:HE3	1:A:6:ILE:CD1	2.51	0.40
1:A:159:VAL:HG13	1:A:163:THR:HB	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	388/419 (93%)	368 (95%)	20 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	325/349 (93%)	288 (89%)	37 (11%)	5	10

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	GLN
1	A	12	ILE
1	A	18	ASN
1	A	30	THR
1	A	35	ARG
1	A	43	LYS
1	A	48	TYR
1	A	49	ASN
1	A	65	ASP
1	A	114	VAL
1	A	115	VAL
1	A	117	THR
1	A	132	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	133	LYS
1	A	144	LEU
1	A	146	GLU
1	A	155	VAL
1	A	156	ARG
1	A	158	THR
1	A	164	LYS
1	A	181	ILE
1	A	186	LYS
1	A	222	SER
1	A	229	THR
1	A	231	VAL
1	A	254	ILE
1	A	277	ILE
1	A	301	ARG
1	A	318	ASP
1	A	320	VAL
1	A	332	VAL
1	A	364	PRO
1	A	367	LYS
1	A	371	VAL
1	A	397	LYS
1	A	401	TYR
1	A	403	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	9	GLN
1	A	16	GLN
1	A	20	HIS
1	A	211	GLN
1	A	223	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	9ZN	A	501	-	21,24,24	1.94	6 (28%)	27,36,36	2.38	11 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	9ZN	A	501	-	-	8/20/21/21	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	9ZN	CAS-CAR	-4.33	1.39	1.50
2	A	501	9ZN	CAQ-CAJ	-4.30	1.43	1.50
2	A	501	9ZN	CA-N	-3.61	1.43	1.46
2	A	501	9ZN	PAU-OAX	2.90	1.59	1.50
2	A	501	9ZN	CAL-CAO	-2.74	1.40	1.46
2	A	501	9ZN	CAO-N	2.15	1.31	1.27

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	9ZN	CAL-CAO-N	-5.25	112.42	122.73
2	A	501	9ZN	CA-N-CAO	5.17	124.61	116.89
2	A	501	9ZN	OAT-CAS-CAR	-4.08	101.71	109.36
2	A	501	9ZN	CAL-CAK-CAJ	3.73	122.24	120.14
2	A	501	9ZN	OAV-PAU-OAX	-3.18	98.44	110.83
2	A	501	9ZN	CAS-CAR-CAM	-2.69	114.97	119.36
2	A	501	9ZN	CAK-CAL-CAR	-2.69	116.11	118.28
2	A	501	9ZN	OAW-PAU-OAV	2.55	117.36	107.80
2	A	501	9ZN	OAW-PAU-OAT	2.50	113.20	106.67
2	A	501	9ZN	CAK-CAL-CAO	2.33	124.61	120.40
2	A	501	9ZN	CG1-CB-CG2	2.26	115.45	109.54

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	501	9ZN	CB-CA-N-CAO
2	A	501	9ZN	N-CA-CB-CG2
2	A	501	9ZN	N-CA-CB-CG1
2	A	501	9ZN	N-CA-CB-SG
2	A	501	9ZN	C-CA-CB-CG2
2	A	501	9ZN	C-CA-CB-CG1
2	A	501	9ZN	CAL-CAR-CAS-OAT
2	A	501	9ZN	CAM-CAR-CAS-OAT

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	9ZN	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	392/419 (93%)	-0.65	3 (0%) 82 80	57, 80, 125, 173	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	63	ALA	4.2
1	A	48	TYR	3.4
1	A	49	ASN	2.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	9ZN	A	501	24/24	0.97	0.08	54,74,87,117	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different

orientation to approximate a three-dimensional view.

Electron density around 9ZN A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.