



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2026 – 07:44 PM UTC

PDB ID : 5XFY / pdb_00005xfy
Title : Crystal structure of a novel PET hydrolase S131A mutant from Ideonella sakaiensis 201-F6
Authors : Han, X.; Liu, W.D.; Zheng, Y.Y.; Chen, C.C.; Guo, R.T.
Deposited on : 2017-04-11
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.40-1.40)
Clashscore	190562	2660 (1.40-1.40)
Ramachandran outliers	187476	2611 (1.40-1.40)
Sidechain outliers	187428	2610 (1.40-1.40)
RSRZ outliers	180081	2561 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	261	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2142 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Poly(ethylene terephthalate) hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	261	Total	C	N	O	S	0	4	0
			1931	1195	343	382	11			

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	131	ALA	SER	engineered mutation	UNP A0A0K8P6T7

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	190	Total	O	0	0
			190	190		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Poly(ethylene terephthalate) hydrolase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	114.19Å 50.33Å 40.98Å 90.00° 92.24° 90.00°	Depositor
Resolution (Å)	25.00 – 1.40 25.00 – 1.40	Depositor EDS
% Data completeness (in resolution range)	99.6 (25.00-1.40) 99.6 (25.00-1.40)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	9.39 (at 1.40Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.134 , 0.159 0.141 , 0.163	Depositor DCC
R_{free} test set	2306 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	14.6	Xtriage
Anisotropy	0.053	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.45 , 50.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.021 for -h,-k,l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	2142	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	2.22	108/1988 (5.4%)	1.07	8/2708 (0.3%)

All (108) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	55[A]	VAL	CA-C	14.35	1.68	1.52
1	A	55[B]	VAL	CA-C	14.35	1.68	1.52
1	A	185[A]	SER	CA-C	12.01	1.68	1.52
1	A	185[B]	SER	CA-C	12.01	1.68	1.52
1	A	154	ALA	C-O	-10.89	1.18	1.23
1	A	187	LEU	C-O	-8.87	1.16	1.24
1	A	177	ASP	C-O	-8.61	1.13	1.24
1	A	32	SER	C-O	-8.46	1.14	1.23
1	A	41	TYR	C-O	-8.44	1.15	1.24
1	A	42	PRO	C-O	-7.84	1.16	1.23
1	A	28	VAL	C-O	-7.68	1.16	1.24
1	A	52	ILE	C-O	-7.64	1.16	1.24
1	A	55[A]	VAL	C-O	-7.50	1.15	1.24
1	A	55[B]	VAL	C-O	-7.50	1.15	1.24
1	A	77	PHE	C-O	-7.50	1.14	1.23
1	A	146	SER	C-O	-7.49	1.14	1.24
1	A	54	ILE	C-O	-7.42	1.16	1.24
1	A	228	TRP	C-O	-7.06	1.16	1.24
1	A	127	VAL	C-O	-7.05	1.16	1.24
1	A	185[A]	SER	C-O	-6.91	1.15	1.24
1	A	185[B]	SER	C-O	-6.91	1.15	1.24
1	A	219	ALA	C-O	-6.89	1.16	1.24
1	A	192	SER	C-O	-6.75	1.15	1.24
1	A	189	ILE	C-O	-6.67	1.16	1.24
1	A	70	PRO	C-O	-6.65	1.16	1.24
1	A	60	ALA	C-O	-6.55	1.15	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	243	ALA	C-O	-6.52	1.16	1.24
1	A	183	ASN	C-O	-6.44	1.15	1.24
1	A	200	PHE	C-O	-6.44	1.16	1.23
1	A	181	PRO	C-O	-6.42	1.16	1.23
1	A	94	ARG	C-O	-6.36	1.16	1.24
1	A	188	PRO	C-O	-6.35	1.16	1.24
1	A	170	LEU	C-O	-6.30	1.16	1.24
1	A	6	GLY	C-O	-6.29	1.15	1.24
1	A	124	ARG	C-O	-6.28	1.16	1.23
1	A	245	GLU	C-O	-6.22	1.16	1.23
1	A	222	GLY	C-O	-6.20	1.16	1.23
1	A	131	ALA	C-O	-6.17	1.16	1.24
1	A	159	SER	C-O	-6.16	1.16	1.23
1	A	120	VAL	C-O	-6.12	1.17	1.24
1	A	167	VAL	C-O	-6.11	1.16	1.24
1	A	220	LEU	C-O	-6.11	1.17	1.24
1	A	95	SER	C-O	-6.09	1.17	1.24
1	A	169	THR	C-O	-6.09	1.17	1.24
1	A	68	TRP	C-O	-6.08	1.16	1.24
1	A	225	GLY	C-O	-6.05	1.16	1.23
1	A	59	THR	C-O	-6.05	1.16	1.24
1	A	11	ALA	C-O	-6.04	1.17	1.24
1	A	130	TRP	C-O	-6.00	1.16	1.24
1	A	218	GLN	C-O	-5.99	1.16	1.24
1	A	14	LEU	C-O	-5.97	1.16	1.24
1	A	201	LEU	C-O	-5.96	1.17	1.24
1	A	62	GLN	C-O	-5.95	1.16	1.24
1	A	78	VAL	C-O	-5.89	1.17	1.24
1	A	241	THR	C-O	-5.88	1.17	1.24
1	A	223	LYS	C-O	-5.85	1.17	1.24
1	A	79	VAL	C-O	-5.84	1.17	1.24
1	A	13	SER	C-O	-5.80	1.16	1.24
1	A	80	ILE	C-O	-5.80	1.17	1.24
1	A	112	SER	C-O	-5.79	1.16	1.24
1	A	63	SER	C-O	-5.76	1.16	1.24
1	A	193	MET	C-O	-5.65	1.17	1.24
1	A	153	GLN	C-O	-5.62	1.17	1.24
1	A	227	ALA	C-O	-5.62	1.17	1.24
1	A	186	ALA	C-O	-5.61	1.17	1.24
1	A	237	THR	C-O	-5.60	1.16	1.24
1	A	182	VAL	C-O	-5.59	1.17	1.24
1	A	129	GLY	C-O	-5.53	1.16	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	209	SER	C-O	-5.52	1.17	1.24
1	A	190	TYR	C-O	-5.50	1.17	1.24
1	A	135	GLY	C-O	-5.50	1.17	1.23
1	A	231	ARG	C-O	-5.49	1.17	1.24
1	A	136	GLY	C-O	-5.48	1.17	1.23
1	A	30	ARG	C-O	-5.44	1.18	1.24
1	A	229	MET	C-O	-5.44	1.17	1.24
1	A	81	THR	C-O	-5.39	1.17	1.24
1	A	138	LEU	C-O	-5.39	1.17	1.24
1	A	69	GLY	C-O	-5.38	1.16	1.24
1	A	203	ILE	C-O	-5.36	1.18	1.24
1	A	98	GLN	C-O	-5.35	1.18	1.24
1	A	156	TRP	C-O	-5.34	1.17	1.23
1	A	144	ASN	C-O	-5.33	1.18	1.24
1	A	75	HIS	C-O	-5.32	1.16	1.23
1	A	101	ALA	C-O	-5.32	1.17	1.24
1	A	180	ALA	C-O	-5.30	1.18	1.24
1	A	71	ARG	C-O	-5.29	1.18	1.24
1	A	149	ALA	C-O	-5.29	1.17	1.23
1	A	206	GLY	C-O	-5.28	1.18	1.23
1	A	23	VAL	C-O	-5.28	1.18	1.24
1	A	162	PHE	C-O	-5.26	1.16	1.23
1	A	260	CYS	C-O	-5.23	1.17	1.23
1	A	132	MET	C-O	-5.21	1.17	1.24
1	A	66	LYS	C-O	-5.17	1.17	1.24
1	A	118	GLY	C-O	-5.17	1.17	1.23
1	A	93	SER	C-O	-5.12	1.18	1.24
1	A	133	GLY	C-O	-5.12	1.17	1.23
1	A	128	MET	C-O	-5.12	1.17	1.23
1	A	230	LYS	C-O	-5.11	1.18	1.24
1	A	161	ASN	C-O	-5.11	1.17	1.23
1	A	168	PRO	C-O	-5.08	1.17	1.23
1	A	134	GLY	C-O	-5.08	1.17	1.23
1	A	143	ASN	C-O	-5.08	1.17	1.24
1	A	123	ALA	C-O	-5.07	1.17	1.24
1	A	72	LEU	C-O	-5.07	1.17	1.24
1	A	150	ALA	C-O	-5.06	1.18	1.23
1	A	21	PHE	C-O	-5.05	1.18	1.23
1	A	97	GLN	C-O	-5.01	1.18	1.24
1	A	96	SER	C-O	-5.00	1.18	1.24

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	17[A]	SER	N-CA-C	7.48	121.50	112.38
1	A	17[B]	SER	N-CA-C	7.48	121.50	112.38
1	A	35	GLY	N-CA-C	6.38	120.20	112.49
1	A	216[A]	SER	N-CA-C	6.24	120.58	113.15
1	A	216[B]	SER	N-CA-C	6.24	120.58	113.15
1	A	82	ILE	N-CA-C	5.67	117.84	108.99
1	A	197	ALA	N-CA-C	-5.37	101.91	109.96
1	A	245	GLU	N-CA-C	5.32	118.00	110.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1931	0	1874	19	1
2	A	15	0	0	1	0
3	A	6	0	8	0	0
4	A	190	0	0	2	2
All	All	2142	0	1882	19	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:55[B]:VAL:HG21	4:A:535:HOH:O	1.70	0.89
1:A:85:ASN:H	1:A:97:GLN:HE22	1.20	0.85
1:A:119:LYS:NZ	4:A:401:HOH:O	2.18	0.75
1:A:161:ASN:HD22	1:A:161:ASN:C	2.01	0.68
1:A:85:ASN:N	1:A:97:GLN:HE22	1.97	0.60
1:A:246:ASN:C	1:A:246:ASN:HD22	2.10	0.59
1:A:199:GLN:HE22	1:A:224:LYS:NZ	2.02	0.58
1:A:85:ASN:H	1:A:97:GLN:NE2	1.95	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:55[B]:VAL:HG23	1:A:133:GLY:HA3	1.99	0.45
1:A:55[A]:VAL:HG21	1:A:98:GLN:HB2	1.99	0.44
1:A:131:ALA:HB1	2:A:301:SO4:S	2.58	0.44
1:A:161:ASN:HD22	1:A:163:SER:H	1.66	0.43
1:A:161:ASN:ND2	1:A:163:SER:H	2.17	0.42
1:A:55[B]:VAL:HG23	1:A:133:GLY:CA	2.50	0.42
1:A:55[B]:VAL:HG11	1:A:98:GLN:HB2	2.02	0.42
1:A:1:ASN:C	1:A:1:ASN:HD22	2.28	0.42
1:A:131:ALA:HA	1:A:154:ALA:O	2.19	0.41
1:A:199:GLN:HE22	1:A:224:LYS:HZ1	1.68	0.41
1:A:1:ASN:ND2	1:A:3:TYR:H	2.18	0.41

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:259:ASN:ND2	1:A:259:ASN:ND2[2_556]	1.58	0.62
4:A:495:HOH:O	4:A:569:HOH:O[1_554]	1.77	0.43
4:A:403:HOH:O	4:A:570:HOH:O[2_556]	2.06	0.14

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	263/261 (101%)	259 (98%)	4 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	210/206 (102%)	208 (99%)	2 (1%)	68 40

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	32	SER
1	A	246	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1	ASN
1	A	62	GLN
1	A	85	ASN
1	A	90	GLN
1	A	97	GLN
1	A	143	ASN
1	A	161	ASN
1	A	199	GLN
1	A	204	ASN
1	A	218	GLN
1	A	246	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	304	-	5,5,5	0.10	0	5,5,5	0.83	0
2	SO4	A	301	-	4,4,4	0.20	0	6,6,6	0.27	0
2	SO4	A	302	-	4,4,4	0.50	0	6,6,6	0.20	0
2	SO4	A	303	-	4,4,4	0.50	0	6,6,6	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	304	-	-	1/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	304	GOL	C1-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	261/261 (100%)	-0.00	1 (0%) 88 91	8, 15, 24, 48	4 (1%)

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	260	CYS	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	A	302	5/5	0.73	0.14	50,54,59,85	0
3	GOL	A	304	6/6	0.87	0.12	32,36,41,43	0
2	SO4	A	303	5/5	0.94	0.08	25,30,39,41	0
2	SO4	A	301	5/5	0.99	0.05	13,15,15,17	0

6.5 Other polymers [i](#)

There are no such residues in this entry.