



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:56 PM UTC

PDB ID : 4XGH / pdb_00004xgh
Title : X-ray Crystal Structure of Citrate Synthase from Burkholderia thailandensis
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2014-12-30
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	454	13% 82% • 14%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3133 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Citrate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	390	2889	1834	493	539	23	0	1	0

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	expression tag	UNP Q2T7I5
A	-19	ALA	-	expression tag	UNP Q2T7I5
A	-18	HIS	-	expression tag	UNP Q2T7I5
A	-17	HIS	-	expression tag	UNP Q2T7I5
A	-16	HIS	-	expression tag	UNP Q2T7I5
A	-15	HIS	-	expression tag	UNP Q2T7I5
A	-14	HIS	-	expression tag	UNP Q2T7I5
A	-13	HIS	-	expression tag	UNP Q2T7I5
A	-12	MET	-	expression tag	UNP Q2T7I5
A	-11	GLY	-	expression tag	UNP Q2T7I5
A	-10	THR	-	expression tag	UNP Q2T7I5
A	-9	LEU	-	expression tag	UNP Q2T7I5
A	-8	GLU	-	expression tag	UNP Q2T7I5
A	-7	ALA	-	expression tag	UNP Q2T7I5
A	-6	GLN	-	expression tag	UNP Q2T7I5
A	-5	THR	-	expression tag	UNP Q2T7I5
A	-4	GLN	-	expression tag	UNP Q2T7I5
A	-3	GLY	-	expression tag	UNP Q2T7I5
A	-2	PRO	-	expression tag	UNP Q2T7I5
A	-1	GLY	-	expression tag	UNP Q2T7I5
A	0	SER	-	expression tag	UNP Q2T7I5

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	202	Total	O	0	0
			202	202		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Citrate synthase



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	120.25Å 120.25Å 219.65Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.57 – 2.10 48.57 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.1 (48.57-2.10) 99.2 (48.57-2.10)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.64 (at 2.10Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.208 , 0.230 0.209 , 0.232	Depositor DCC
R_{free} test set	1745 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	34.6	Xtriage
Anisotropy	0.537	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 62.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3133	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.28	0/2953	0.72	2/4016 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	286	PRO	CA-C-N	5.82	132.18	121.70
1	A	286	PRO	C-N-CA	5.82	132.18	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2889	0	2695	8	0
2	A	30	0	0	1	0
3	A	12	0	18	0	0
4	A	202	0	0	0	0
All	All	3133	0	2713	8	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (8) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:192:PRO:HB3	1:A:201:ASN:HD21	1.72	0.55
1:A:286:PRO:HA	1:A:287:ASP:CB	2.40	0.52
1:A:323:MET:HG3	1:A:367:VAL:HG23	1.93	0.50
1:A:348:LEU:HD12	1:A:365:PRO:HG3	1.97	0.46
1:A:384:MET:O	1:A:388:ILE:HG12	2.20	0.42
1:A:5:ASP:HB2	1:A:24:TYR:HD1	1.86	0.41
1:A:111:THR:HG22	1:A:167:ILE:HG21	2.03	0.41
1:A:127:ARG:NH2	2:A:504:SO4:O4	2.53	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	381/454 (84%)	369 (97%)	10 (3%)	2 (0%)	24	22

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	235	GLU
1	A	286	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	279/378 (74%)	275 (99%)	4 (1%)	59 67

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	ASP
1	A	173	LEU
1	A	235	GLU
1	A	283	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	234	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	A	506	-	4,4,4	0.23	0	6,6,6	0.07	0
2	SO4	A	505	-	4,4,4	0.23	0	6,6,6	0.07	0
3	EDO	A	507	-	3,3,3	0.43	0	2,2,2	0.37	0
3	EDO	A	509	-	3,3,3	0.43	0	2,2,2	0.35	0
2	SO4	A	502	-	4,4,4	0.24	0	6,6,6	0.08	0
2	SO4	A	503	-	4,4,4	0.24	0	6,6,6	0.07	0
3	EDO	A	508	-	3,3,3	0.43	0	2,2,2	0.39	0
2	SO4	A	504	-	4,4,4	0.24	0	6,6,6	0.08	0
2	SO4	A	501	-	4,4,4	0.23	0	6,6,6	0.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	507	-	-	0/1/1/1	-
3	EDO	A	508	-	-	0/1/1/1	-
3	EDO	A	509	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	504	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	390/454 (85%)	0.91	61 (15%) 5 5	20, 44, 111, 151	1 (0%)

All (61) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	341	LEU	7.4
1	A	153	SER	6.0
1	A	7	LYS	5.7
1	A	210	PRO	5.5
1	A	267	PRO	5.5
1	A	6	VAL	4.9
1	A	266	GLY	4.7
1	A	316	TYR	4.7
1	A	286	PRO	4.5
1	A	340	PRO	4.5
1	A	348	LEU	4.4
1	A	284	GLY	4.3
1	A	270	GLY	4.2
1	A	8	ALA	4.2
1	A	367	VAL	4.1
1	A	328	TYR	3.8
1	A	369	PHE	3.8
1	A	285	THR	3.8
1	A	151	HIS	3.8
1	A	339	ASP	3.7
1	A	16	SER	3.5
1	A	364	TYR	3.5
1	A	347	GLN	3.4
1	A	338	ASP	3.4
1	A	345	ALA	3.4
1	A	337	HIS	3.4
1	A	209	ASN	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	370	TYR	3.3
1	A	292	PHE	3.3
1	A	10	LEU	3.2
1	A	289	ILE	3.1
1	A	279	MET	3.1
1	A	280	LEU	3.1
1	A	288	ASN	3.0
1	A	363	LEU	3.0
1	A	276	ALA	2.9
1	A	342	PHE	2.8
1	A	15	ASN	2.7
1	A	211	CYS	2.7
1	A	373	ILE	2.7
1	A	283	ILE	2.6
1	A	405	ILE	2.5
1	A	293	ILE	2.5
1	A	40	GLY	2.5
1	A	22	PRO	2.5
1	A	344	LEU	2.4
1	A	413	GLY	2.4
1	A	162	ARG	2.4
1	A	350	LYS	2.3
1	A	39	TYR	2.3
1	A	5	ASP	2.3
1	A	294	LYS	2.3
1	A	213	GLU	2.2
1	A	412	ILE	2.2
1	A	159	PRO	2.2
1	A	58	ASN	2.2
1	A	318	PRO	2.1
1	A	12	PHE	2.1
1	A	19	VAL	2.1
1	A	152	ASP	2.1
1	A	13	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SO4	A	506	5/5	0.77	0.19	133,133,134,134	0
3	EDO	A	509	4/4	0.78	0.28	70,71,73,75	0
3	EDO	A	508	4/4	0.80	0.21	68,69,70,71	0
2	SO4	A	502	5/5	0.80	0.16	113,114,114,115	0
2	SO4	A	504	5/5	0.84	0.16	91,91,91,91	5
2	SO4	A	503	5/5	0.84	0.12	124,124,124,125	0
2	SO4	A	505	5/5	0.85	0.13	90,90,91,92	0
3	EDO	A	507	4/4	0.85	0.16	68,69,71,72	0
2	SO4	A	501	5/5	0.96	0.09	44,46,53,53	0

6.5 Other polymers [i](#)

There are no such residues in this entry.