



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 07:03 PM UTC

PDB ID : 4XGX / pdb_00004xgx
Title : Crystal structure of Escherichia coli Flavin trafficking protein, an FMN transferase, Y60N mutant, ADP-inhibited
Authors : Tomchick, D.R.; Brautigam, C.A.; Deka, R.K.; Norgard, M.V.
Deposited on : 2015-01-02
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	340	13% 82% 15%
1	B	340	7% 80% 6% 14%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 9560 atoms, of which 4566 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FAD:protein FMN transferase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	290	Total	C	H	N	O	S	0	3	0
			4462	1400	2233	383	435	11			
1	B	293	Total	C	H	N	O	S	0	8	0
			4568	1435	2285	390	446	12			


There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP P0AB85
A	60	ASN	TYR	engineered mutation	UNP P0AB85
A	333	LEU	-	expression tag	UNP P0AB85
A	334	GLU	-	expression tag	UNP P0AB85
A	335	HIS	-	expression tag	UNP P0AB85
A	336	HIS	-	expression tag	UNP P0AB85
A	337	HIS	-	expression tag	UNP P0AB85
A	338	HIS	-	expression tag	UNP P0AB85
A	339	HIS	-	expression tag	UNP P0AB85
A	340	HIS	-	expression tag	UNP P0AB85
B	1	MET	-	initiating methionine	UNP P0AB85
B	60	ASN	TYR	engineered mutation	UNP P0AB85
B	333	LEU	-	expression tag	UNP P0AB85
B	334	GLU	-	expression tag	UNP P0AB85
B	335	HIS	-	expression tag	UNP P0AB85
B	336	HIS	-	expression tag	UNP P0AB85
B	337	HIS	-	expression tag	UNP P0AB85
B	338	HIS	-	expression tag	UNP P0AB85
B	339	HIS	-	expression tag	UNP P0AB85
B	340	HIS	-	expression tag	UNP P0AB85

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Mg 2 2	0	0
2	B	2	Total Mg 2 2	0	0

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
3	A	1	Total 39	C 10	H 12	N 5	O 10	P 2	0	0
3	B	1	Total 39	C 10	H 12	N 5	O 10	P 2	0	0

- Molecule 4 is NITRATE ION (CCD ID: NO3) (formula: NO_3).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	N	O	0	0
			4	1	3		
4	A	1	Total	N	O	0	0
			4	1	3		
4	A	1	Total	N	O	0	0
			4	1	3		
4	A	1	Total	N	O	0	0
			4	1	3		
4	A	1	Total	N	O	0	0
			4	1	3		
4	B	1	Total	N	O	0	0
			4	1	3		
4	B	1	Total	N	O	0	0
			4	1	3		
4	B	1	Total	N	O	0	0
			4	1	3		

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	H	O	0	0
			10	2	6	2		
5	A	1	Total	C	H	O	0	0
			10	2	6	2		
5	B	1	Total	C	H	O	0	0
			10	2	6	2		
5	B	1	Total	C	H	O	0	0
			10	2	6	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	160	Total	O	0	0
			160	160		
6	B	212	Total	O	0	0
			212	212		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: FAD:protein FMN transferase



- Molecule 1: FAD:protein FMN transferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	54.58Å 56.98Å 224.77Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.26 – 1.90 28.26 – 1.90	Depositor EDS
% Data completeness (in resolution range)	88.4 (28.26-1.90) 88.4 (28.26-1.90)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.44 (at 1.91Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.180 , 0.214 0.183 , 0.216	Depositor DCC
R_{free} test set	2495 reflections (4.43%)	wwPDB-VP
Wilson B-factor (Å ²)	19.1	Xtriage
Anisotropy	0.186	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 49.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.026 for k,h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9560	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ADP, NO3, MG, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.36	0/2278	0.64	0/3086
1	B	0.39	0/2353	0.65	0/3185
All	All	0.38	0/4631	0.64	0/6271

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2229	2233	2226	6	0
1	B	2283	2285	2271	11	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	27	12	12	1	0
3	B	27	12	12	0	0
4	A	20	0	0	0	0
4	B	16	0	0	0	0
5	A	8	12	12	0	0
5	B	8	12	12	0	0
6	A	160	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	212	0	0	2	0
All	All	4994	4566	4545	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:207:PRO:O	1:B:209:ARG:NH1	2.33	0.62
1:A:14:GLU:O	1:A:33:GLY:N	2.33	0.56
1:A:178:ARG:NH2	6:A:504:HOH:O	2.37	0.56
1:B:241:ASN:ND2	6:B:505:HOH:O	2.39	0.53
1:B:189:LEU:HD13	1:B:198:SER:HB3	1.92	0.52
1:B:21:LYS:NZ	6:B:508:HOH:O	2.43	0.52
1:A:110:VAL:HA	1:A:254:ILE:CD1	2.42	0.50
1:B:110:VAL:HA	1:B:254:ILE:HD12	1.93	0.49
1:A:23[A]:MET:HE2	1:A:240:ARG:HH21	1.79	0.47
1:A:23[A]:MET:HE1	1:A:168:GLY:HA3	1.97	0.47
1:B:63[A]:ASP:OD1	1:B:68:ARG:NH2	2.53	0.42
1:B:25:THR:OG1	1:B:26:PHE:N	2.51	0.42
1:B:115:PHE:CD2	1:B:252:HIS:N	2.88	0.41
1:B:212:ILE:HD13	1:B:323:PHE:HZ	1.85	0.41
1:A:163:ASP:OD2	3:A:403:ADP:O2'	2.26	0.41
1:B:166:THR:OG1	1:B:284:THR:OG1	2.33	0.41
1:B:23[A]:MET:HE1	1:B:168:GLY:HA3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	285/340 (84%)	277 (97%)	8 (3%)	0	100	100
1	B	293/340 (86%)	286 (98%)	6 (2%)	1 (0%)	36	29
All	All	578/680 (85%)	563 (97%)	14 (2%)	1 (0%)	43	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	264	HIS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	246/288 (85%)	245 (100%)	1 (0%)	84	87
1	B	254/288 (88%)	251 (99%)	3 (1%)	63	63
All	All	500/576 (87%)	496 (99%)	4 (1%)	73	75

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	269	VAL
1	B	166	THR
1	B	252	HIS
1	B	280	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	146	ASN
1	A	147	GLN
1	A	150	GLN
1	A	151	GLN
1	A	154	GLN
1	B	150	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	151	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 19 ligands modelled in this entry, 4 are monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NO3	A	408	-	1,3,3	0.64	0	0,3,3	-	-
4	NO3	A	404	-	1,3,3	0.41	0	0,3,3	-	-
5	EDO	B	409	-	3,3,3	0.43	0	2,2,2	0.38	0
4	NO3	A	405	-	1,3,3	0.50	0	0,3,3	-	-
4	NO3	B	407	-	1,3,3	0.50	0	0,3,3	-	-
5	EDO	A	409	-	3,3,3	0.37	0	2,2,2	0.39	0
4	NO3	A	407	-	1,3,3	0.55	0	0,3,3	-	-
5	EDO	A	410	-	3,3,3	0.38	0	2,2,2	0.42	0
3	ADP	A	403	2	28,29,29	1.31	4 (14%)	43,45,45	1.94	12 (27%)
4	NO3	B	404	-	1,3,3	0.54	0	0,3,3	-	-
4	NO3	B	405	-	1,3,3	0.55	0	0,3,3	-	-
5	EDO	B	408	-	3,3,3	0.42	0	2,2,2	0.40	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ADP	B	403	2	28,29,29	1.32	5 (17%)	43,45,45	1.98	11 (25%)
4	NO3	A	406	-	1,3,3	0.50	0	0,3,3	-	-
4	NO3	B	406	-	1,3,3	0.56	0	0,3,3	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	B	409	-	-	0/1/1/1	-
5	EDO	A	410	-	-	0/1/1/1	-
5	EDO	A	409	-	-	0/1/1/1	-
3	ADP	A	403	2	-	4/16/32/32	0/3/3/3
5	EDO	B	408	-	-	0/1/1/1	-
3	ADP	B	403	2	-	6/16/32/32	0/3/3/3

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	403	ADP	C5-C4	4.05	1.46	1.39
3	A	403	ADP	C5-C4	3.89	1.46	1.39
3	A	403	ADP	PA-O3A	2.71	1.62	1.59
3	B	403	ADP	PA-O3A	2.49	1.62	1.59
3	A	403	ADP	C8-N7	2.24	1.36	1.31
3	B	403	ADP	C5-C6	2.20	1.47	1.41
3	B	403	ADP	C8-N7	2.18	1.35	1.31
3	B	403	ADP	C4-N9	-2.09	1.33	1.37
3	A	403	ADP	C5-C6	2.07	1.46	1.41

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	403	ADP	C5-C4-N3	-6.02	118.42	126.72
3	A	403	ADP	C5-C4-N3	-5.83	118.69	126.72
3	B	403	ADP	N3-C4-N9	5.32	136.22	127.17
3	A	403	ADP	N3-C4-N9	5.16	135.95	127.17
3	B	403	ADP	C2-N3-C4	4.15	121.96	111.83
3	A	403	ADP	C2-N3-C4	4.09	121.82	111.83
3	B	403	ADP	C4-N9-C8	3.74	109.67	105.74
3	A	403	ADP	N3-C2-N1	-3.64	123.08	128.58

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	403	ADP	C4-N9-C8	3.61	109.53	105.74
3	B	403	ADP	N3-C2-N1	-3.61	123.12	128.58
3	A	403	ADP	C5-C6-N6	-2.96	115.95	123.29
3	B	403	ADP	C4-C5-N7	-2.56	107.65	110.58
3	B	403	ADP	N9-C8-N7	-2.39	110.54	113.94
3	A	403	ADP	C4-C5-N7	-2.37	107.88	110.58
3	A	403	ADP	N9-C8-N7	-2.35	110.61	113.94
3	B	403	ADP	C5-C6-N6	-2.27	117.66	123.29
3	B	403	ADP	O4'-C1'-N9	-2.26	103.75	108.09
3	A	403	ADP	N6-C6-N1	2.25	123.39	118.38
3	B	403	ADP	C6-C5-N7	2.19	136.30	132.09
3	B	403	ADP	C5-N7-C8	2.12	106.78	103.45
3	A	403	ADP	C6-C5-N7	2.11	136.16	132.09
3	A	403	ADP	O3B-PB-O2B	2.09	115.62	107.80
3	A	403	ADP	C5-N7-C8	2.03	106.63	103.45

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	403	ADP	PA-O3A-PB-O1B
3	A	403	ADP	C2'-C1'-N9-C8
3	B	403	ADP	PA-O3A-PB-O2B
3	B	403	ADP	C2'-C1'-N9-C8
3	B	403	ADP	C5'-O5'-PA-O1A
3	A	403	ADP	C2'-C1'-N9-C4
3	B	403	ADP	C2'-C1'-N9-C4
3	B	403	ADP	O4'-C1'-N9-C8
3	A	403	ADP	PA-O3A-PB-O2B
3	A	403	ADP	O4'-C1'-N9-C8



There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	403	ADP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	290/340 (85%)	0.42	43 (14%) 5 6	9, 30, 83, 114	3 (1%)
1	B	293/340 (86%)	-0.00	25 (8%) 16 17	8, 23, 67, 111	8 (2%)
All	All	583/680 (85%)	0.21	68 (11%) 9 10	8, 26, 79, 114	11 (1%)

All (68) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	122	VAL	7.1
1	A	122	VAL	6.6
1	B	265	ASN	4.9
1	B	264	HIS	4.8
1	A	13	THR	4.8
1	A	203	GLY	4.5
1	A	312	GLY	4.4
1	B	121	PRO	4.2
1	A	36	ALA	4.0
1	A	263	GLU	3.9
1	B	311	GLU	3.8
1	A	329	SER	3.8
1	A	315	PHE	3.7
1	A	37	LYS	3.6
1	A	265	ASN	3.6
1	B	314	SER	3.5
1	A	34	ILE	3.5
1	B	252	HIS	3.5
1	A	257	GLN	3.4
1	A	33	GLY	3.3
1	A	212	ILE	3.3
1	A	206	LEU	3.3
1	B	115	PHE	3.3
1	A	114	GLY	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	241	ASN	3.1
1	B	310	LYS	3.1
1	B	266	LEU	3.0
1	B	263	GLU	2.9
1	B	329	SER	2.9
1	B	206	LEU	2.9
1	B	212	ILE	2.8
1	B	213	GLN	2.8
1	B	312	GLY	2.8
1	A	251	SER	2.7
1	A	60	ASN	2.7
1	B	214	LYS	2.7
1	B	224	ALA	2.7
1	A	17	VAL	2.7
1	A	313	ASP	2.7
1	A	224	ALA	2.6
1	B	315	PHE	2.6
1	A	63	ASP	2.5
1	B	241	ASN	2.5
1	A	314	SER	2.5
1	A	14	GLU	2.5
1	A	201	MET	2.5
1	A	209	ARG	2.4
1	A	316	LYS	2.3
1	B	60	ASN	2.3
1	A	267	VAL	2.3
1	A	225[A]	VAL	2.2
1	A	16	THR	2.2
1	A	128	GLU	2.2
1	A	207	PRO	2.2
1	A	62	LYS	2.2
1	A	124	ILE	2.2
1	B	114	GLY	2.2
1	A	310	LYS	2.2
1	B	316	LYS	2.2
1	A	266	LEU	2.1
1	A	159	ASP	2.1
1	A	40	ALA	2.1
1	B	201[A]	MET	2.1
1	A	15	VAL	2.1
1	B	225	VAL	2.1
1	A	202	ASN	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	328	VAL	2.0
1	A	317	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	EDO	B	409	4/4	0.74	0.20	52,62,67,67	0
4	NO3	A	408	4/4	0.77	0.19	43,47,50,52	0
5	EDO	B	408	4/4	0.83	0.19	29,43,52,63	0
5	EDO	A	410	4/4	0.86	0.17	40,48,53,57	0
5	EDO	A	409	4/4	0.87	0.18	40,50,61,71	0
4	NO3	A	405	4/4	0.88	0.10	36,50,50,55	0
4	NO3	B	404	4/4	0.88	0.12	44,48,50,53	0
4	NO3	A	407	4/4	0.89	0.12	47,47,52,53	0
4	NO3	B	407	4/4	0.89	0.16	54,58,61,67	0
3	ADP	A	403	27/27	0.90	0.11	9,36,64,70	0
2	MG	B	402	1/1	0.90	0.13	39,39,39,39	0
4	NO3	B	405	4/4	0.91	0.18	39,41,48,52	0
4	NO3	A	406	4/4	0.91	0.19	41,45,49,53	0
3	ADP	B	403	27/27	0.92	0.14	8,27,74,161	0
2	MG	A	402	1/1	0.94	0.16	40,40,40,40	0
4	NO3	B	406	4/4	0.95	0.08	23,26,29,31	0
4	NO3	A	404	4/4	0.96	0.09	21,26,26,34	0
2	MG	B	401	1/1	0.97	0.09	22,22,22,22	0
2	MG	A	401	1/1	0.99	0.07	21,21,21,21	0


The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ADP A 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.