



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:51 PM UTC

PDB ID : 7XGL / pdb_00007xgl
Title : Quinolate Phosphoribosyl Transferase (QAPRTase) from Streptomyces pyridomyceticus NRRL B-2517 in Apo form
Authors : Zhou, Z.; Yang, X.; Huang, T.; Wang, X.; Liang, R.; Zheng, J.; Dai, S.; Lin, S.; Deng, Z.
Deposited on : 2022-04-05
Resolution : 2.11 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.11 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	8290 (2.14-2.10)
Clashscore	190562	8817 (2.14-2.10)
Ramachandran outliers	187476	8738 (2.14-2.10)
Sidechain outliers	187428	8739 (2.14-2.10)
RSRZ outliers	180081	8294 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	308	
1	B	308	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	B	402	-	-	X	-

2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 4700 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Quinolate Phosphoribosyl Transferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	297	Total	C	N	O	S	0	0	0
			2173	1340	405	421	7			
1	B	294	Total	C	N	O	S	0	2	0
			2174	1339	404	424	7			

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is hexanedioic acid (CCD ID: 0L1) (formula: $C_6H_{10}O_4$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			10	6	4		

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Cl	0	0
			2	2		
5	B	1	Total	Cl	0	0
			1	1		

- Molecule 6 is TRIS-HYDROXYMETHYL-METHYL-AMMONIUM (CCD ID: 144) (formula: C₄H₁₂NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	B	1	Total	C	N	O	0	0
			8	4	1	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	160	Total	O	0	0
			160	160		
7	B	124	Total	O	0	0
			124	124		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Quinolate Phosphoribosyl Transferase



- Molecule 1: Quinolate Phosphoribosyl Transferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	120.05Å 150.93Å 55.18Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.73 – 2.11 37.73 – 2.11	Depositor EDS
% Data completeness (in resolution range)	99.2 (37.73-2.11) 99.4 (37.73-2.11)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.41 (at 2.10Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.176 , 0.207 0.176 , 0.206	Depositor DCC
R_{free} test set	2986 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	27.6	Xtriage
Anisotropy	0.045	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 34.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4700	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4, OL1, 144, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.29	0/2198	0.46	0/2994
1	B	0.28	0/2198	0.45	0/2992
All	All	0.29	0/4396	0.46	0/5986

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2173	0	2195	16	0
1	B	2174	0	2188	19	0
2	A	15	0	0	0	0
2	B	15	0	0	1	0
3	A	6	0	8	0	0
3	B	12	0	16	9	0
4	A	10	0	8	3	0
5	A	2	0	0	0	0
5	B	1	0	0	0	0
6	B	8	0	12	3	0
7	A	160	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	B	124	0	0	2	0
All	All	4700	0	4427	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:117:ARG:HH21	4:A:405:0L1:H5	1.54	0.71
1:A:154:LEU:HD13	1:B:154:LEU:HD13	1.76	0.66
1:B:186:ASN:ND2	3:B:402:GOL:H12	2.11	0.66
1:B:186:ASN:HD22	3:B:402:GOL:H31	1.61	0.64
1:A:154:LEU:HD11	6:B:403:144:H41	1.81	0.62
1:A:85:GLU:HG2	1:A:106:THR:HB	1.86	0.56
1:A:62:GLN:NE2	1:A:292:ASP:H	2.05	0.53
1:A:173:HIS:HB3	7:A:516:HOH:O	2.08	0.53
1:A:288:ALA:O	6:B:403:144:H22	2.07	0.53
1:A:181:ILE:HD13	1:A:201:ALA:HB1	1.91	0.52
1:B:186:ASN:HB2	3:B:402:GOL:C3	2.40	0.52
1:B:186:ASN:CG	3:B:402:GOL:H12	2.35	0.52
1:B:260:SER:HB3	1:B:280:ALA:HB3	1.93	0.51
1:A:117:ARG:NH2	4:A:405:0L1:H5	2.23	0.50
1:B:186:ASN:HB2	3:B:402:GOL:H31	1.93	0.50
1:B:287:SER:HA	6:B:403:144:H21	1.94	0.48
1:A:57:VAL:HG13	1:A:101:VAL:HG13	1.95	0.48
1:B:186:ASN:ND2	3:B:402:GOL:H31	2.28	0.47
1:B:232:MET:HE3	7:B:509:HOH:O	2.13	0.47
1:A:181:ILE:HD11	1:A:205:THR:OG1	2.17	0.44
1:A:167:ARG:NE	1:A:167:ARG:HA	2.33	0.44
4:A:405:0L1:H4	3:B:402:GOL:O2	2.17	0.44
3:B:401:GOL:O1	2:B:406:SO4:O1	2.30	0.43
1:B:150:SER:HB3	7:B:570:HOH:O	2.17	0.43
1:B:174:ARG:HG2	1:B:176:GLY:O	2.18	0.43
1:B:202[B]:ARG:NH2	1:B:229:ASP:OD2	2.51	0.42
1:A:298:ALA:HA	1:A:299:PRO:HD3	1.83	0.42
1:B:148:ARG:NE	1:B:258:GLU:OE2	2.51	0.42
1:B:28:VAL:HG13	1:B:75:VAL:HG22	2.01	0.42
1:B:182:LEU:HG	1:B:184:LYS:HE2	2.01	0.42
1:B:60:SER:HA	1:B:293:LEU:HD23	2.01	0.41
1:B:269:ALA:O	1:B:273:ARG:HG2	2.19	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:61:ARG:NE	7:A:509:HOH:O	2.53	0.41
1:A:86:VAL:HG22	1:A:105:VAL:HG12	2.02	0.40
1:A:62:GLN:HE21	1:A:292:ASP:H	1.70	0.40
1:B:186:ASN:HB2	3:B:402:GOL:H32	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	295/308 (96%)	288 (98%)	5 (2%)	2 (1%)	18	15
1	B	294/308 (96%)	289 (98%)	5 (2%)	0	100	100
All	All	589/616 (96%)	577 (98%)	10 (2%)	2 (0%)	36	36

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	4	GLU
1	A	5	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	217/228 (95%)	212 (98%)	5 (2%)	44	50
1	B	218/228 (96%)	214 (98%)	4 (2%)	51	59
All	All	435/456 (95%)	426 (98%)	9 (2%)	47	54

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	42	VAL
1	A	53	VAL
1	A	212	VAL
1	A	252	ARG
1	A	254	ARG
1	B	4	GLU
1	B	173	HIS
1	B	174	ARG
1	B	256	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	62	GLN
1	A	235	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 14 ligands modelled in this entry, 3 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	B	406	-	4,4,4	0.15	0	6,6,6	0.24	0
2	SO4	A	401	-	4,4,4	0.21	0	6,6,6	0.48	0
2	SO4	B	404	-	4,4,4	0.27	0	6,6,6	0.21	0
2	SO4	B	405	-	4,4,4	0.18	0	6,6,6	0.24	0
4	OL1	A	405	-	9,9,9	1.27	1 (11%)	10,10,10	1.25	1 (10%)
2	SO4	A	403	-	4,4,4	0.22	0	6,6,6	0.50	0
3	GOL	B	402	-	5,5,5	1.50	2 (40%)	5,5,5	0.78	0
2	SO4	A	402	-	4,4,4	0.30	0	6,6,6	0.20	0
3	GOL	A	404	-	5,5,5	1.00	0	5,5,5	0.98	0
3	GOL	B	401	-	5,5,5	0.87	0	5,5,5	1.10	0
6	144	B	403	-	1,7,7	1.26	0	3,9,9	0.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	OL1	A	405	-	-	5/7/7/7	-
3	GOL	B	402	-	-	2/4/4/4	-
3	GOL	A	404	-	-	2/4/4/4	-
3	GOL	B	401	-	-	2/4/4/4	-
6	144	B	403	-	-	0/0/9/9	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	405	OL1	O2-C2	2.17	1.29	1.22
3	B	402	GOL	C3-C2	2.12	1.59	1.51
3	B	402	GOL	O2-C2	-2.04	1.37	1.43

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	405	OL1	O3-C7-C6	2.24	121.08	114.00

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	404	GOL	C1-C2-C3-O3
3	B	401	GOL	C1-C2-C3-O3
3	A	404	GOL	O2-C2-C3-O3
3	B	402	GOL	O1-C1-C2-O2
3	B	402	GOL	O1-C1-C2-C3
3	B	401	GOL	O2-C2-C3-O3
4	A	405	OL1	C3-C4-C5-C6
4	A	405	OL1	C5-C6-C7-O3
4	A	405	OL1	O1-C2-C3-C4
4	A	405	OL1	C5-C6-C7-O4
4	A	405	OL1	O2-C2-C3-C4

There are no ring outliers.

5 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	406	SO4	1	0
4	A	405	OL1	3	0
3	B	402	GOL	8	0
3	B	401	GOL	1	0
6	B	403	144	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	297/308 (96%)	-0.22	11 (3%) 45 48	17, 27, 45, 76	0
1	B	294/308 (95%)	-0.14	11 (3%) 45 48	15, 28, 46, 76	2 (0%)
All	All	591/616 (95%)	-0.18	22 (3%) 45 48	15, 28, 46, 76	2 (0%)

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	3	ALA	4.9
1	A	253	THR	4.5
1	B	4	GLU	4.4
1	A	4	GLU	4.0
1	A	299	PRO	3.9
1	A	252	ARG	3.5
1	B	173	HIS	3.4
1	A	254	ARG	3.0
1	B	254	ARG	3.0
1	B	8	PRO	3.0
1	A	5	ALA	2.9
1	A	144	HIS	2.9
1	B	144	HIS	2.9
1	B	209	PRO	2.6
1	B	95	ARG	2.6
1	B	252	ARG	2.4
1	A	8	PRO	2.3
1	B	191	GLY	2.2
1	B	192	GLY	2.1
1	B	5	ALA	2.1
1	A	298	ALA	2.1
1	A	9	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	B	402	6/6	0.81	0.38	43,45,46,51	0
4	0L1	A	405	10/10	0.81	0.18	35,45,51,52	0
5	CL	A	407	1/1	0.81	0.25	65,65,65,65	0
6	144	B	403	8/8	0.91	0.23	28,38,40,41	0
3	GOL	A	404	6/6	0.92	0.18	33,40,45,47	0
5	CL	B	407	1/1	0.92	0.24	48,48,48,48	0
5	CL	A	406	1/1	0.92	0.27	59,59,59,59	0
3	GOL	B	401	6/6	0.94	0.12	40,44,48,52	0
2	SO4	B	404	5/5	0.95	0.14	45,47,60,62	0
2	SO4	A	401	5/5	0.97	0.06	26,26,32,38	0
2	SO4	A	403	5/5	0.98	0.08	24,28,32,32	0
2	SO4	B	406	5/5	0.98	0.06	29,33,36,38	0
2	SO4	B	405	5/5	0.99	0.05	29,31,34,36	0
2	SO4	A	402	5/5	1.00	0.03	23,25,26,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.