



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:19 AM UTC

PDB ID : 4XHI / pdb_00004xhi
Title : Crystal structure of native Thosea asigna virus RNA-dependent RNA polymerase (RdRP) at 2.15 Angstrom resolution
Authors : Ferrero, D.S.; Buxaderas, M.; Rodriguez, J.F.; Verdaguer, N.
Deposited on : 2015-01-05
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	705	
1	B	705	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 11380 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RNA-dependent RNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	663	Total	C	N	O	S	0	3	0
			5265	3342	914	983	26			
1	B	664	Total	C	N	O	S	0	1	0
			5261	3340	910	984	27			

There are 62 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-30	MET	-	initiating methionine	UNP Q6A562
A	-29	SER	-	expression tag	UNP Q6A562
A	-28	TYR	-	expression tag	UNP Q6A562
A	-27	TYR	-	expression tag	UNP Q6A562
A	-26	HIS	-	expression tag	UNP Q6A562
A	-25	HIS	-	expression tag	UNP Q6A562
A	-24	HIS	-	expression tag	UNP Q6A562
A	-23	HIS	-	expression tag	UNP Q6A562
A	-22	HIS	-	expression tag	UNP Q6A562
A	-21	HIS	-	expression tag	UNP Q6A562
A	-20	ASP	-	expression tag	UNP Q6A562
A	-19	TYR	-	expression tag	UNP Q6A562
A	-18	ASP	-	expression tag	UNP Q6A562
A	-17	ILE	-	expression tag	UNP Q6A562
A	-16	PRO	-	expression tag	UNP Q6A562
A	-15	THR	-	expression tag	UNP Q6A562
A	-14	THR	-	expression tag	UNP Q6A562
A	-13	GLU	-	expression tag	UNP Q6A562
A	-12	ASN	-	expression tag	UNP Q6A562
A	-11	LEU	-	expression tag	UNP Q6A562
A	-10	TYR	-	expression tag	UNP Q6A562
A	-9	PHE	-	expression tag	UNP Q6A562
A	-8	GLN	-	expression tag	UNP Q6A562
A	-7	GLY	-	expression tag	UNP Q6A562
A	-6	ALA	-	expression tag	UNP Q6A562

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	MET	-	expression tag	UNP Q6A562
A	-4	GLY	-	expression tag	UNP Q6A562
A	-3	ALA	-	expression tag	UNP Q6A562
A	-2	MET	-	expression tag	UNP Q6A562
A	-1	GLY	-	expression tag	UNP Q6A562
A	0	ILE	-	expression tag	UNP Q6A562
B	-30	MET	-	initiating methionine	UNP Q6A562
B	-29	SER	-	expression tag	UNP Q6A562
B	-28	TYR	-	expression tag	UNP Q6A562
B	-27	TYR	-	expression tag	UNP Q6A562
B	-26	HIS	-	expression tag	UNP Q6A562
B	-25	HIS	-	expression tag	UNP Q6A562
B	-24	HIS	-	expression tag	UNP Q6A562
B	-23	HIS	-	expression tag	UNP Q6A562
B	-22	HIS	-	expression tag	UNP Q6A562
B	-21	HIS	-	expression tag	UNP Q6A562
B	-20	ASP	-	expression tag	UNP Q6A562
B	-19	TYR	-	expression tag	UNP Q6A562
B	-18	ASP	-	expression tag	UNP Q6A562
B	-17	ILE	-	expression tag	UNP Q6A562
B	-16	PRO	-	expression tag	UNP Q6A562
B	-15	THR	-	expression tag	UNP Q6A562
B	-14	THR	-	expression tag	UNP Q6A562
B	-13	GLU	-	expression tag	UNP Q6A562
B	-12	ASN	-	expression tag	UNP Q6A562
B	-11	LEU	-	expression tag	UNP Q6A562
B	-10	TYR	-	expression tag	UNP Q6A562
B	-9	PHE	-	expression tag	UNP Q6A562
B	-8	GLN	-	expression tag	UNP Q6A562
B	-7	GLY	-	expression tag	UNP Q6A562
B	-6	ALA	-	expression tag	UNP Q6A562
B	-5	MET	-	expression tag	UNP Q6A562
B	-4	GLY	-	expression tag	UNP Q6A562
B	-3	ALA	-	expression tag	UNP Q6A562
B	-2	MET	-	expression tag	UNP Q6A562
B	-1	GLY	-	expression tag	UNP Q6A562
B	0	ILE	-	expression tag	UNP Q6A562

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	366	Total	O	0	0
			366	366		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	357	Total	O	0	0
			357	357		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: RNA-dependent RNA polymerase



• Molecule 1: RNA-dependent RNA polymerase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	134.88Å 150.82Å 100.61Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.61 – 2.15 39.61 – 2.15	Depositor EDS
% Data completeness (in resolution range)	88.9 (39.61-2.15) 88.9 (39.61-2.15)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.45 (at 2.16Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, R_{free}	0.175 , 0.216 0.178 , 0.216	Depositor DCC
R_{free} test set	4979 reflections (4.45%)	wwPDB-VP
Wilson B-factor (Å ²)	19.6	Xtriage
Anisotropy	0.060	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 36.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	11380	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.01% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/5394	0.69	2/7305 (0.0%)
1	B	0.28	0/5383	0.67	0/7289
All	All	0.28	0/10777	0.68	2/14594 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	642	HIS	CA-C-N	7.68	127.73	119.89
1	A	642	HIS	C-N-CA	7.68	127.73	119.89

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	641	ASP	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5265	0	5230	30	0
1	B	5261	0	5225	29	0
2	A	35	0	0	0	0
2	B	30	0	0	1	0
3	A	36	0	48	1	0
3	B	30	0	40	2	0
4	A	366	0	0	2	0
4	B	357	0	0	6	0
All	All	11380	0	10543	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:641:ASP:OD1	1:B:642:HIS:N	2.05	0.88
1:A:163:GLN:NE2	4:A:1157:HOH:O	2.32	0.62
1:B:636:SER:HB2	1:B:643:PRO:HA	1.84	0.60
1:B:553:THR:H	1:B:556:MET:HE3	1.67	0.59
1:A:371:LYS:NZ	3:A:712:GOL:O3	2.37	0.58
1:A:164:ARG:NH1	1:A:372[A]:GLN:OE1	2.37	0.57
1:A:328:ASP:OD1	1:A:640[A]:ARG:NH1	2.37	0.57
1:A:81:GLN:NE2	1:A:163:GLN:OE1	2.36	0.56
1:B:54:ARG:NH1	2:B:703:SO4:O3	2.39	0.56
1:B:45:ALA:O	4:B:801:HOH:O	2.18	0.55
1:B:350:GLY:HA3	3:B:705:GOL:H11	1.90	0.54
1:A:549:ARG:HB2	1:A:552:GLU:HG2	1.90	0.54
1:A:317:TYR:HA	1:A:349:TYR:CD1	2.43	0.54
1:A:391:ASP:HA	1:A:394:VAL:HG22	1.91	0.53
1:B:532:LEU:HB2	1:B:537:TRP:CE2	2.44	0.53
1:A:641:ASP:HB3	1:A:643:PRO:HD3	1.91	0.52
1:A:332:TRP:CE2	1:A:345:ARG:HD2	2.45	0.52
1:B:110:HIS:H	3:B:708:GOL:H2	1.76	0.51
1:A:605:ARG:NH2	1:A:621:GLU:OE1	2.41	0.50
1:A:225:ARG:O	1:B:22:ARG:HD2	2.12	0.50
1:B:332:TRP:CE2	1:B:345:ARG:HD2	2.47	0.49
1:B:642:HIS:O	4:B:1155:HOH:O	2.20	0.49
1:A:22:ARG:HD2	1:B:225:ARG:O	2.12	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:521:ARG:NH1	4:B:1128:HOH:O	2.44	0.48
1:B:566:ARG:NH2	1:B:624:ILE:O	2.40	0.47
1:A:532:LEU:HB2	1:A:537:TRP:CE2	2.50	0.47
1:B:633:TRP:O	1:B:636:SER:OG	2.31	0.47
1:B:142:ASP:O	1:B:146:GLU:HG3	2.14	0.46
1:B:391:ASP:HA	1:B:394:VAL:HG22	1.97	0.46
1:B:598:ARG:NH1	4:B:934:HOH:O	2.39	0.46
1:B:476:ILE:HG12	1:B:487:ILE:HB	1.97	0.46
1:A:414:MET:HA	1:A:418:SER:OG	2.15	0.46
1:A:13:LEU:HD21	1:B:212:ALA:HB2	1.97	0.45
1:B:588:ASN:ND2	4:B:801:HOH:O	2.49	0.45
1:B:457:ASP:O	1:B:461:GLN:HG2	2.17	0.44
1:A:164:ARG:HE	1:A:372[A]:GLN:HE22	1.66	0.44
1:A:118:VAL:O	1:A:119:LYS:HD3	2.17	0.44
1:A:515:LEU:HD13	1:A:530:PRO:HB3	2.00	0.43
4:A:1162:HOH:O	1:B:662:LYS:HE2	2.18	0.43
1:B:599:VAL:O	1:B:604:GLY:HA2	2.19	0.43
1:B:372:GLN:NE2	4:B:1081:HOH:O	2.49	0.43
1:A:457:ASP:O	1:A:461:GLN:HG2	2.19	0.42
1:A:553:THR:HG1	1:A:556:MET:HG3	1.83	0.42
1:A:228:LEU:HB3	1:A:229:PRO:HD3	2.02	0.41
1:A:119:LYS:HD3	1:A:119:LYS:HA	1.62	0.41
1:A:164:ARG:HE	1:A:372[A]:GLN:NE2	2.19	0.41
1:A:650:PHE:HA	1:A:651:PRO:HD2	1.94	0.41
1:B:373:MET:HE1	1:B:451:SER:HB2	2.02	0.41
1:A:163:GLN:O	1:A:167:MET:HG3	2.21	0.41
1:A:599:VAL:O	1:A:604:GLY:HA2	2.19	0.41
1:B:522:ARG:HB2	1:B:635:VAL:HG11	2.02	0.41
1:A:617:ASP:N	1:A:617:ASP:OD1	2.53	0.41
1:B:252:ARG:NH2	1:B:413:GLU:OE1	2.47	0.41
1:B:628:TYR:HA	1:B:629:PRO:HD3	1.88	0.40
1:A:593:GLU:O	1:A:597:MET:HB3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	664/705 (94%)	647 (97%)	16 (2%)	1 (0%)	43	44
1	B	663/705 (94%)	648 (98%)	13 (2%)	2 (0%)	36	34
All	All	1327/1410 (94%)	1295 (98%)	29 (2%)	3 (0%)	43	44

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	641	ASP
1	B	317	TYR
1	B	372	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	565/598 (94%)	560 (99%)	5 (1%)	70	77
1	B	564/598 (94%)	560 (99%)	4 (1%)	76	82
All	All	1129/1196 (94%)	1120 (99%)	9 (1%)	73	79

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	119	LYS
1	A	208	ILE
1	A	213	ILE
1	A	430	LYS
1	A	619	VAL
1	B	126	VAL
1	B	192	GLN
1	B	341	LYS
1	B	619	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	168	GLN
1	A	192	GLN
1	B	28	GLN
1	B	168	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

24 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	710	-	5,5,5	0.37	0	5,5,5	0.36	0
2	SO4	B	711	-	4,4,4	0.24	0	6,6,6	0.07	0
3	GOL	A	711	-	5,5,5	0.36	0	5,5,5	0.34	0
3	GOL	B	706	-	5,5,5	0.37	0	5,5,5	0.39	0
2	SO4	B	701	-	4,4,4	0.23	0	6,6,6	0.06	0
2	SO4	A	706	-	4,4,4	0.23	0	6,6,6	0.07	0
2	SO4	A	702	-	4,4,4	0.23	0	6,6,6	0.08	0
2	SO4	B	710	-	4,4,4	0.23	0	6,6,6	0.07	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	B	702	-	4,4,4	0.23	0	6,6,6	0.08	0
2	SO4	B	709	-	4,4,4	0.23	0	6,6,6	0.09	0
3	GOL	B	704	-	5,5,5	0.36	0	5,5,5	0.32	0
2	SO4	A	704	-	4,4,4	0.23	0	6,6,6	0.08	0
3	GOL	A	707	-	5,5,5	0.39	0	5,5,5	0.32	0
2	SO4	A	701	-	4,4,4	0.23	0	6,6,6	0.06	0
3	GOL	A	712	-	5,5,5	0.36	0	5,5,5	0.34	0
2	SO4	A	703	-	4,4,4	0.24	0	6,6,6	0.08	0
3	GOL	A	708	-	5,5,5	0.37	0	5,5,5	0.33	0
2	SO4	B	703	-	4,4,4	0.23	0	6,6,6	0.07	0
3	GOL	B	708	-	5,5,5	0.36	0	5,5,5	0.32	0
2	SO4	A	713	-	4,4,4	0.24	0	6,6,6	0.07	0
3	GOL	A	709	-	5,5,5	0.37	0	5,5,5	0.32	0
3	GOL	B	705	-	5,5,5	0.37	0	5,5,5	0.39	0
2	SO4	A	705	-	4,4,4	0.24	0	6,6,6	0.08	0
3	GOL	B	707	-	5,5,5	0.38	0	5,5,5	0.31	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	708	-	-	2/4/4/4	-
3	GOL	A	710	-	-	2/4/4/4	-
3	GOL	A	709	-	-	2/4/4/4	-
3	GOL	B	705	-	-	2/4/4/4	-
3	GOL	A	711	-	-	2/4/4/4	-
3	GOL	A	712	-	-	2/4/4/4	-
3	GOL	B	707	-	-	2/4/4/4	-
3	GOL	B	706	-	-	2/4/4/4	-
3	GOL	B	704	-	-	2/4/4/4	-
3	GOL	A	708	-	-	2/4/4/4	-
3	GOL	A	707	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (22) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	707	GOL	O1-C1-C2-C3
3	A	709	GOL	O1-C1-C2-C3
3	A	712	GOL	O1-C1-C2-C3
3	B	704	GOL	O1-C1-C2-C3
3	B	706	GOL	O1-C1-C2-C3
3	B	707	GOL	O1-C1-C2-C3
3	B	708	GOL	O1-C1-C2-C3
3	A	708	GOL	O1-C1-C2-C3
3	A	710	GOL	O1-C1-C2-C3
3	A	711	GOL	O1-C1-C2-C3
3	B	705	GOL	O1-C1-C2-C3
3	A	707	GOL	O1-C1-C2-O2
3	A	709	GOL	O1-C1-C2-O2
3	A	710	GOL	O1-C1-C2-O2
3	A	711	GOL	O1-C1-C2-O2
3	B	705	GOL	O1-C1-C2-O2
3	B	706	GOL	O1-C1-C2-O2
3	B	707	GOL	O1-C1-C2-O2
3	B	708	GOL	O1-C1-C2-O2
3	A	712	GOL	O1-C1-C2-O2
3	B	704	GOL	O1-C1-C2-O2
3	A	708	GOL	O1-C1-C2-O2

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	712	GOL	1	0
2	B	703	SO4	1	0
3	B	708	GOL	1	0
3	B	705	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	663/705 (94%)	-0.11	27 (4%) 41 46	9, 18, 47, 101	3 (0%)
1	B	664/705 (94%)	0.02	31 (4%) 36 41	11, 22, 48, 96	1 (0%)
All	All	1327/1410 (94%)	-0.04	58 (4%) 39 44	9, 20, 48, 101	4 (0%)

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	548	TYR	11.3
1	B	642	HIS	6.4
1	A	10	PRO	6.3
1	A	642	HIS	5.6
1	B	307	MET	5.6
1	A	11	THR	5.4
1	B	10	PRO	5.1
1	B	643	PRO	5.1
1	A	643	PRO	4.8
1	A	641	ASP	4.6
1	A	307	MET	4.6
1	B	126	VAL	4.5
1	A	349	TYR	4.5
1	B	11	THR	4.5
1	B	552	GLU	3.9
1	B	641	ASP	3.7
1	A	549	ARG	3.6
1	B	12	ARG	3.6
1	B	548	TYR	3.5
1	B	550	SER	3.5
1	B	644	CYS	3.4
1	B	618	GLY	3.4
1	A	127	ASN	3.3
1	B	54	ARG	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	617	ASP	3.3
1	B	125	GLY	3.2
1	B	127	ASN	3.2
1	B	551	LYS	3.2
1	A	551	LYS	3.1
1	A	550	SER	3.0
1	B	549	ARG	3.0
1	B	13	LEU	3.0
1	A	347	ALA	3.0
1	B	496	LEU	3.0
1	A	128	THR	2.7
1	B	553	THR	2.7
1	B	523	GLU	2.6
1	A	101	ASP	2.5
1	A	125	GLY	2.4
1	B	121	GLN	2.4
1	B	101	ASP	2.4
1	A	126	VAL	2.4
1	A	644	CYS	2.4
1	A	12	ARG	2.3
1	A	619	VAL	2.3
1	A	41	ASP	2.2
1	A	119	LYS	2.2
1	B	128	THR	2.2
1	B	619	VAL	2.2
1	A	552	GLU	2.2
1	A	640[A]	ARG	2.2
1	B	62	ARG	2.1
1	A	672	LYS	2.1
1	A	23	ALA	2.1
1	A	617	ASP	2.1
1	B	488	LYS	2.1
1	B	600	GLN	2.0
1	B	640	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	A	707	6/6	0.71	0.22	41,52,57,58	0
3	GOL	B	704	6/6	0.74	0.22	38,50,53,54	0
3	GOL	A	711	6/6	0.75	0.21	44,52,59,61	0
3	GOL	B	707	6/6	0.77	0.24	56,61,64,66	0
3	GOL	A	712	6/6	0.81	0.16	49,50,54,55	0
3	GOL	B	706	6/6	0.82	0.17	34,43,48,54	0
3	GOL	A	710	6/6	0.82	0.18	38,42,48,50	0
2	SO4	B	710	5/5	0.83	0.14	68,70,72,73	0
2	SO4	B	703	5/5	0.83	0.13	89,92,93,95	0
3	GOL	B	705	6/6	0.84	0.19	53,53,55,56	0
3	GOL	A	708	6/6	0.87	0.12	37,40,43,48	0
3	GOL	A	709	6/6	0.88	0.16	35,47,49,50	0
2	SO4	A	703	5/5	0.88	0.16	66,68,69,74	0
3	GOL	B	708	6/6	0.88	0.21	46,48,59,64	0
2	SO4	B	702	5/5	0.92	0.15	39,48,56,57	0
2	SO4	A	705	5/5	0.93	0.18	54,58,64,64	0
2	SO4	B	711	5/5	0.94	0.12	50,53,57,58	0
2	SO4	A	701	5/5	0.97	0.08	24,29,37,42	0
2	SO4	B	709	5/5	0.97	0.11	28,30,40,41	0
2	SO4	A	706	5/5	0.97	0.08	35,38,41,43	0
2	SO4	A	713	5/5	0.97	0.09	28,34,39,43	0
2	SO4	A	704	5/5	0.97	0.13	37,37,42,43	0
2	SO4	A	702	5/5	0.98	0.07	27,31,35,41	0
2	SO4	B	701	5/5	0.98	0.06	29,35,39,39	0

6.5 Other polymers ⓘ

There are no such residues in this entry.