



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 17, 2026 – 10:08 PM UTC

PDB ID : 8XIS / pdb_00008xis
Title : Cellodextrin phosphorylase from Clostridium thermocellum mutant - all cysteine residues were substituted with serines
Authors : Kuga, T.; Sunagawa, N.; Igarashi, K.
Deposited on : 2023-12-19
Resolution : 1.68 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.68 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1054 (1.68-1.68)
Clashscore	190562	1078 (1.68-1.68)
Ramachandran outliers	187476	1068 (1.68-1.68)
Sidechain outliers	187428	1067 (1.68-1.68)
RSRZ outliers	180081	1055 (1.68-1.68)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	992	 85% 14% ..
1	B	992	 87% 12% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	ACT	A	1004	-	-	X	-
5	ACT	B	1004	-	-	X	-
5	ACT	B	1006	-	-	X	-
5	ACT	B	1014	-	-	X	-
6	PEG	B	1012	-	-	X	-

2 Entry composition [i](#)


There are 8 unique types of molecules in this entry. The entry contains 32463 atoms, of which 14633 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cellodextrin phosphorylase variant.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	984	Total	C	H	N	O	S	190	19	0
			15225	5142	7192	1336	1533	22			
1	B	984	Total	C	H	N	O	S	188	22	0
			15305	5155	7258	1333	1537	22			

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is beta-D-glucopyranose (CCD ID: BGC) (formula: $C_6H_{12}O_6$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			12	6	6		
3	B	1	Total	C	O	0	0
			12	6	6		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			14	3	8	3		
4	A	1	Total	C	H	O	0	0
			14	3	8	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			14	3	8	3		
4	A	1	Total	C	H	O	0	0
			14	3	8	3		
4	A	1	Total	C	H	O	0	0
			14	3	8	3		
4	A	1	Total	C	H	O	0	0
			14	3	8	3		
4	B	1	Total	C	H	O	0	0
			14	3	8	3		
4	B	1	Total	C	H	O	0	0
			14	3	8	3		
4	B	1	Total	C	H	O	0	0
			14	3	8	3		
4	B	1	Total	C	H	O	0	0
			14	3	8	3		
4	B	1	Total	C	H	O	0	0
			14	3	8	3		
4	B	1	Total	C	H	O	0	0
			14	3	8	3		
4	B	1	Total	C	H	O	0	0
			14	3	8	3		

- Molecule 5 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	H	O	0	0
			7	2	3	2		
5	A	1	Total	C	H	O	0	0
			7	2	3	2		
5	A	1	Total	C	H	O	0	0
			7	2	3	2		
5	A	1	Total	C	H	O	0	0
			7	2	3	2		
5	A	1	Total	C	H	O	0	0
			7	2	3	2		
5	B	1	Total	C	H	O	0	0
			7	2	3	2		
5	B	1	Total	C	H	O	0	0
			7	2	3	2		
5	B	1	Total	C	H	O	0	0
			7	2	3	2		
5	B	1	Total	C	H	O	0	0
			7	2	3	2		

- Molecule 6 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	H	O	0	0
			17	4	10	3		
6	A	1	Total	C	H	O	0	0
			17	4	10	3		
6	B	1	Total	C	H	O	0	0
			17	4	10	3		

- Molecule 7 is CHLORIDE ION (CCD ID: CL) (formula: Cl) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	2	Total	Cl	0	0
			2	2		
7	B	2	Total	Cl	0	0
			2	2		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	793	Total	O	0	0
			793	793		
8	B	764	Total	O	0	0
			764	764		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: cellodextrin phosphorylase variant



- Molecule 1: cellodextrin phosphorylase variant



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	82.28Å 88.52Å 88.54Å 98.41° 110.69° 110.65°	Depositor
Resolution (Å)	43.44 – 1.68 43.44 – 1.68	Depositor EDS
% Data completeness (in resolution range)	97.0 (43.44-1.68) 96.9 (43.44-1.68)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.24 (at 1.68Å)	Xtriage
Refinement program	PHENIX 1.21rc1_5127	Depositor
R, R_{free}	0.188 , 0.230 0.188 , 0.230	Depositor DCC
R_{free} test set	11541 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å ²)	23.4	Xtriage
Anisotropy	0.224	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 40.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.479 for -h,-l,-k	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	32463	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CL, ACT, PEG, BGC, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.46	0/8211	0.62	0/11108
1	B	0.46	0/8225	0.62	0/11132
All	All	0.46	0/16436	0.62	0/22240

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
All	All	0	4

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	494	GLY	Peptide
1	A	495[B]	TYR	Peptide
1	B	494	GLY	Peptide
1	B	495[B]	TYR	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8033	7192	7884	120	0
1	B	8047	7258	7896	116	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	A	12	0	12	0	0
3	B	12	0	12	0	0
4	A	36	48	48	4	0
4	B	54	72	72	7	0
5	A	24	18	18	4	0
5	B	20	15	15	9	0
6	A	14	20	20	2	0
6	B	7	10	10	6	0
7	A	2	0	0	1	0
7	B	2	0	0	0	0
8	A	793	0	0	26	2
8	B	764	0	0	37	2
All	All	17830	14633	15987	242	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 242 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:699:LYS:NZ	1:B:703:ASP:OD2	1.68	1.22
1:B:828:LYS:O	8:B:1101:HOH:O	1.77	1.02
1:A:717[B]:ARG:NH1	7:A:1014:CL:CL	2.30	1.01
1:A:64:SER:O	1:A:68:LYS:HD3	1.69	0.90
1:A:633:ILE:HD11	8:A:1856:HOH:O	1.77	0.84

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:A:1757:HOH:O	8:B:1781:HOH:O[1_444]	2.12	0.08
8:A:1607:HOH:O	8:B:1831:HOH:O[1_444]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1001/992 (101%)	958 (96%)	40 (4%)	3 (0%)	36	21
1	B	1004/992 (101%)	963 (96%)	39 (4%)	2 (0%)	43	27
All	All	2005/1984 (101%)	1921 (96%)	79 (4%)	5 (0%)	36	27

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	798	ASP
1	A	800	ALA
1	B	497	GLU
1	A	796	ALA
1	B	799	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	869/859 (101%)	863 (99%)	6 (1%)	76	62
1	B	872/859 (102%)	867 (99%)	5 (1%)	78	67

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	1741/1718 (101%)	1730 (99%)	11 (1%)	81	67

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	649	THR
1	B	729[A]	THR
1	B	793	ASN
1	B	729[B]	THR
1	A	729[B]	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	58	GLN
1	B	458	GLN
1	B	219	ASN
1	B	612	HIS
1	A	692	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 37 ligands modelled in this entry, 4 are monoatomic - leaving 33 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	ACT	A	1005	-	3,3,3	1.46	1 (33%)	3,3,3	1.22	0
5	ACT	B	1002	-	3,3,3	1.38	1 (33%)	3,3,3	1.53	1 (33%)
3	BGC	B	1001	-	12,12,12	0.44	0	17,17,17	0.97	1 (5%)
4	GOL	A	1002	-	5,5,5	0.32	0	5,5,5	0.59	0
4	GOL	A	1016	-	5,5,5	0.16	0	5,5,5	0.29	0
6	PEG	A	1012	-	6,6,6	0.27	0	5,5,5	0.28	0
4	GOL	B	1011	-	5,5,5	0.30	0	5,5,5	0.40	0
4	GOL	A	1007	-	5,5,5	0.44	0	5,5,5	1.00	1 (20%)
5	ACT	B	1006	-	3,3,3	1.40	1 (33%)	3,3,3	1.13	0
4	GOL	B	1010	-	5,5,5	0.28	0	5,5,5	0.40	0
5	ACT	A	1004	-	3,3,3	1.29	0	3,3,3	1.62	1 (33%)
4	GOL	B	1005	-	5,5,5	0.23	0	5,5,5	0.53	0
5	ACT	B	1014	-	3,3,3	1.41	0	3,3,3	1.23	0
2	PO4	A	1000	-	4,4,4	2.09	1 (25%)	6,6,6	1.51	1 (16%)
5	ACT	A	1011	-	3,3,3	1.35	0	3,3,3	1.07	0
5	ACT	A	1006	-	3,3,3	1.48	1 (33%)	3,3,3	1.06	0
3	BGC	A	1001	-	12,12,12	0.47	0	17,17,17	1.15	2 (11%)
4	GOL	A	1015	-	5,5,5	0.25	0	5,5,5	0.51	0
2	PO4	B	1000	-	4,4,4	1.82	2 (50%)	6,6,6	1.62	1 (16%)
5	ACT	B	1004	-	3,3,3	1.38	0	3,3,3	1.35	0
5	ACT	A	1003	-	3,3,3	1.51	1 (33%)	3,3,3	1.13	0
5	ACT	A	1009	-	3,3,3	1.44	1 (33%)	3,3,3	0.75	0
4	GOL	B	1008	-	5,5,5	0.36	0	5,5,5	0.47	0
5	ACT	B	1007	-	3,3,3	1.39	1 (33%)	3,3,3	1.20	0
4	GOL	A	1008	-	5,5,5	0.30	0	5,5,5	0.42	0
4	GOL	B	1017	-	5,5,5	0.10	0	5,5,5	0.48	0
6	PEG	A	1010	-	6,6,6	0.34	0	5,5,5	0.22	0
4	GOL	B	1018	-	5,5,5	0.26	0	5,5,5	0.36	0
4	GOL	A	1017	-	5,5,5	0.29	0	5,5,5	0.40	0
4	GOL	B	1009	-	5,5,5	0.37	0	5,5,5	0.63	0
6	PEG	B	1012	-	6,6,6	0.53	0	5,5,5	0.37	0
4	GOL	B	1003	-	5,5,5	0.15	0	5,5,5	0.80	0
4	GOL	B	1013	-	5,5,5	0.35	0	5,5,5	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BGC	B	1001	-	-	2/2/22/22	0/1/1/1
4	GOL	A	1002	-	-	4/4/4/4	-
4	GOL	A	1016	-	-	4/4/4/4	-
4	GOL	B	1003	-	-	4/4/4/4	-
4	GOL	B	1011	-	-	4/4/4/4	-
4	GOL	A	1007	-	-	2/4/4/4	-
4	GOL	B	1010	-	-	2/4/4/4	-
4	GOL	B	1005	-	-	0/4/4/4	-
3	BGC	A	1001	-	-	2/2/22/22	0/1/1/1
4	GOL	A	1015	-	-	3/4/4/4	-
4	GOL	B	1008	-	-	0/4/4/4	-
4	GOL	A	1008	-	-	2/4/4/4	-
4	GOL	B	1017	-	-	0/4/4/4	-
6	PEG	A	1010	-	-	2/4/4/4	-
4	GOL	A	1017	-	-	2/4/4/4	-
4	GOL	B	1009	-	-	2/4/4/4	-
4	GOL	B	1018	-	-	2/4/4/4	-
6	PEG	B	1012	-	-	3/4/4/4	-
6	PEG	A	1012	-	-	2/4/4/4	-
4	GOL	B	1013	-	-	2/4/4/4	-

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1000	PO4	P-O1	3.97	1.59	1.50
2	B	1000	PO4	P-O1	2.83	1.57	1.50
5	A	1003	ACT	CH3-C	2.12	1.57	1.49
5	A	1006	ACT	CH3-C	2.08	1.57	1.49
5	B	1002	ACT	CH3-C	2.08	1.57	1.49

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1000	PO4	O3-P-O2	3.22	117.93	107.91
3	A	1001	BGC	C1-O5-C5	3.06	119.58	113.65
2	A	1000	PO4	O4-P-O3	2.86	116.80	107.91

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1001	BGC	C1-O5-C5	2.64	118.75	113.65
3	A	1001	BGC	O5-C1-C2	2.32	114.37	110.30

There are no chirality outliers.

5 of 44 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1002	GOL	O1-C1-C2-C3
4	A	1002	GOL	C1-C2-C3-O3
4	A	1007	GOL	C1-C2-C3-O3
4	A	1007	GOL	O2-C2-C3-O3
4	A	1016	GOL	C1-C2-C3-O3


There are no ring outliers.


16 monomers are involved in 32 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1002	GOL	1	0
4	A	1016	GOL	1	0
4	A	1007	GOL	1	0
5	B	1006	ACT	2	0
4	B	1010	GOL	1	0
5	A	1004	ACT	2	0
5	B	1014	ACT	2	0
5	B	1004	ACT	5	0
5	A	1003	ACT	1	0
5	A	1009	ACT	1	0
4	A	1008	GOL	1	0
4	B	1017	GOL	1	0
6	A	1010	PEG	2	0
4	B	1009	GOL	2	0
6	B	1012	PEG	6	0
4	B	1003	GOL	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	984/992 (99%)	-0.94	0 100 100	10, 31, 58, 104	20 (2%)
1	B	984/992 (99%)	-0.95	0 100 100	9, 32, 57, 84	22 (2%)
All	All	1968/1984 (99%)	-0.94	0 100 100	9, 31, 58, 104	42 (2%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	B	1017	6/6	0.94	0.07	60,72,84,85	0
5	ACT	B	1004	4/4	0.95	0.06	45,54,56,62	0
4	GOL	A	1016	6/6	0.96	0.07	40,61,81,81	0
3	BGC	A	1001	12/12	0.96	0.07	39,55,64,66	0
3	BGC	B	1001	12/12	0.96	0.08	39,56,70,73	0
5	ACT	B	1014	4/4	0.96	0.06	41,50,51,53	0
6	PEG	A	1010	7/7	0.96	0.07	31,58,78,78	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	A	1008	6/6	0.97	0.06	34,41,54,54	0
5	ACT	A	1003	4/4	0.97	0.06	43,51,54,57	0
5	ACT	A	1011	4/4	0.97	0.06	59,66,71,71	0
4	GOL	A	1017	6/6	0.97	0.06	49,73,88,101	0
5	ACT	B	1007	4/4	0.97	0.06	45,51,54,54	0
4	GOL	B	1003	6/6	0.97	0.06	27,42,58,64	0
4	GOL	B	1010	6/6	0.97	0.06	48,61,75,75	0
6	PEG	A	1012	7/7	0.97	0.08	58,70,83,83	0
5	ACT	A	1004	4/4	0.98	0.05	35,50,56,56	0
5	ACT	A	1005	4/4	0.98	0.04	41,50,58,59	0
5	ACT	A	1006	4/4	0.98	0.05	46,55,56,57	0
5	ACT	A	1009	4/4	0.98	0.06	43,52,55,55	0
4	GOL	A	1002	6/6	0.98	0.04	41,51,65,74	0
5	ACT	B	1002	4/4	0.98	0.06	26,31,50,54	0
4	GOL	B	1011	6/6	0.98	0.05	43,60,70,72	0
5	ACT	B	1006	4/4	0.98	0.05	42,44,51,51	0
4	GOL	B	1013	6/6	0.98	0.06	36,50,69,72	0
4	GOL	B	1005	6/6	0.98	0.06	50,60,64,69	0
4	GOL	B	1018	6/6	0.98	0.05	56,68,74,77	0
4	GOL	B	1009	6/6	0.98	0.06	35,51,62,62	0
6	PEG	B	1012	7/7	0.98	0.06	27,48,68,68	0
2	PO4	A	1000	5/5	0.99	0.04	27,28,33,39	0
4	GOL	A	1007	6/6	0.99	0.03	25,38,46,55	0
2	PO4	B	1000	5/5	0.99	0.04	25,29,32,34	0
4	GOL	A	1015	6/6	0.99	0.04	29,48,69,69	0
4	GOL	B	1008	6/6	0.99	0.03	23,35,43,43	0
7	CL	A	1014	1/1	0.99	0.04	27,27,27,27	0
7	CL	B	1016	1/1	0.99	0.06	27,27,27,27	0
7	CL	B	1015	1/1	1.00	0.02	17,17,17,17	0
7	CL	A	1013	1/1	1.00	0.02	18,18,18,18	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around BGC A 1001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BGC B 1001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PO4 A 1000:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PO4 B 1000:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CL A 1014:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CL B 1016:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




Electron density around CL B 1015:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around CL A 1013:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.