



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 05:43 PM UTC

PDB ID : 2XJA / pdb_00002xja
Title : Structure of MurE from M.tuberculosis with dipeptide and ADP
Authors : Basavannacharya, C.; Moody, P.R.; Bhakta, S.; Keep, N.
Deposited on : 2010-07-03
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	535	 58% 28% 7% 6%
1	B	535	 56% 30% 7% 7%
1	C	535	 56% 30% 7% 7%
1	D	535	 55% 29% 7% 9%

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 14477 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UDP-N-ACETYLMURAMOYL-L-ALANYL-D-GLUTAMATE--2,6-DIAMINOPIMELATE LIGASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	502	Total	C	N	O	S	0	0	0
			3567	2220	659	680	8			
1	B	500	Total	C	N	O	S	0	0	0
			3556	2211	660	677	8			
1	C	498	Total	C	N	O	S	0	0	0
			3538	2203	655	672	8			
1	D	488	Total	C	N	O	S	0	0	0
			3462	2159	633	662	8			

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	D	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Mg	0	0
			2	2		
3	B	3	Total	Mg	0	0
			3	3		
3	C	2	Total	Mg	0	0
			2	2		
3	D	3	Total	Mg	0	0
			3	3		

- Molecule 4 is URIDINE-5'-DIPHOSPHATE-N-ACETYLMURAMOYL-L-ALANINE-D-GLUTAMATE (CCD ID: UAG) (formula: C₂₈H₄₃N₅O₂₃P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			58	28	5	23	2		
4	B	1	Total	C	N	O	P	0	0
			58	28	5	23	2		
4	C	1	Total	C	N	O	P	0	0
			58	28	5	23	2		

Continued on next page...


Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	D	1	Total	C	N	O	P	0	0
			58	28	5	23	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	O	0	0
			1	1		
5	B	1	Total	O	0	0
			1	1		
5	C	1	Total	O	0	0
			1	1		
5	D	1	Total	O	0	0
			1	1		


● Molecule 1: UDP-N-ACETYLMURAMOYL-L-ALANYL-D-GLUTAMATE-2,6-DIAMINOPI
MELATE LIGASE






• Molecule 1: UDP-N-ACETYLMURAMOYL-L-ALANYL-D-GLUTAMATE--2,6-DIAMINOPI
MELATE LIGASE


Chain C: 56% 30% 7% 7%



• Molecule 1: UDP-N-ACETYLMURAMOYL-L-ALANYL-D-GLUTAMATE--2,6-DIAMINOPI
MELATE LIGASE

Chain D: 55% 29% 7% 9%





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	75.04Å 76.32Å 81.99Å 111.32° 91.42° 92.90°	Depositor
Resolution (Å)	76.30 – 3.00 76.30 – 3.00	Depositor EDS
% Data completeness (in resolution range)	95.9 (76.30-3.00) 95.9 (76.30-3.00)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.39 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.6.0077	Depositor
R, R_{free}	0.189 , 0.258 0.189 , 0.257	Depositor DCC
R_{free} test set	1598 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	62.6	Xtriage
Anisotropy	0.065	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 73.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	14477	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: KCX, UAG, MG, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.95	0/3612	1.12	7/4939 (0.1%)
1	B	1.00	5/3601 (0.1%)	1.13	14/4921 (0.3%)
1	C	0.91	1/3583 (0.0%)	1.07	10/4898 (0.2%)
1	D	0.94	3/3503 (0.1%)	1.10	10/4787 (0.2%)
All	All	0.95	9/14299 (0.1%)	1.11	41/19545 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	C	0	1
All	All	0	3

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	126	ALA	N-CA	6.75	1.51	1.46
1	D	126	ALA	N-CA	6.16	1.50	1.46
1	B	218	VAL	CA-CB	5.54	1.60	1.54
1	B	199	ALA	CA-CB	-5.54	1.47	1.53
1	D	199	ALA	CA-CB	-5.38	1.47	1.53

The worst 5 of 41 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	454	ASP	CA-C-N	10.29	131.40	119.47
1	A	454	ASP	C-N-CA	10.29	131.40	119.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	119	VAL	CA-C-N	8.10	128.11	119.76
1	D	119	VAL	C-N-CA	8.10	128.11	119.76
1	A	119	VAL	CA-C-N	7.71	127.70	119.76

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	115	GLY	Peptide
1	A	116	ARG	Peptide
1	C	116	ARG	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3567	0	3524	149	1
1	B	3556	0	3520	148	0
1	C	3538	0	3504	146	0
1	D	3462	0	3424	147	1
2	A	27	0	12	1	0
2	B	27	0	12	2	0
2	C	27	0	12	4	0
2	D	27	0	12	5	0
3	A	2	0	0	0	0
3	B	3	0	0	0	0
3	C	2	0	0	0	0
3	D	3	0	0	0	0
4	A	58	0	39	5	0
4	B	58	0	39	4	0
4	C	58	0	39	2	0
4	D	58	0	39	7	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
5	D	1	0	0	0	0
All	All	14477	0	14176	564	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 564 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:383:ARG:HG3	1:A:383:ARG:HH11	1.14	1.07
1:C:424:ARG:HG3	1:C:424:ARG:HH11	1.16	1.07
1:A:116:ARG:HG2	1:A:116:ARG:NH1	1.60	1.04
1:D:383:ARG:HG3	1:D:383:ARG:HH11	1.24	1.02
1:B:383:ARG:HG3	1:B:383:ARG:HH11	1.23	1.02

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:269:ASP:O	1:D:269:ASP:O[1_554]	2.10	0.10

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	495/535 (92%)	453 (92%)	39 (8%)	3 (1%)	21	56
1	B	495/535 (92%)	442 (89%)	46 (9%)	7 (1%)	9	36
1	C	493/535 (92%)	448 (91%)	39 (8%)	6 (1%)	10	40
1	D	477/535 (89%)	439 (92%)	35 (7%)	3 (1%)	21	56
All	All	1960/2140 (92%)	1782 (91%)	159 (8%)	19 (1%)	12	45

5 of 19 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	423	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	469	VAL
1	C	116	ARG
1	B	468	GLU
1	B	514	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	341/382 (89%)	281 (82%)	60 (18%)	2	10
1	B	340/382 (89%)	280 (82%)	60 (18%)	2	10
1	C	338/382 (88%)	274 (81%)	64 (19%)	1	9
1	D	332/382 (87%)	269 (81%)	63 (19%)	1	8
All	All	1351/1528 (88%)	1104 (82%)	247 (18%)	2	9

5 of 247 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	498	VAL
1	D	371	GLU
1	C	218	VAL
1	D	369	LEU
1	D	451	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 25 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	333	HIS
1	D	42	GLN
1	D	409	HIS
1	C	409	HIS
1	D	141	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	KCX	C	262	1	10,11,12	1.16	1 (10%)	6,12,14	1.77	1 (16%)
1	KCX	D	262	1	10,11,12	1.22	1 (10%)	6,12,14	1.31	1 (16%)
1	KCX	A	262	1	10,11,12	1.27	2 (20%)	6,12,14	1.24	1 (16%)
1	KCX	B	262	1	10,11,12	1.02	1 (10%)	6,12,14	1.79	2 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	C	262	1	-	0/9/10/12	-
1	KCX	D	262	1	-	0/9/10/12	-
1	KCX	A	262	1	-	1/9/10/12	-
1	KCX	B	262	1	-	1/9/10/12	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	262	KCX	CE-NZ	2.69	1.52	1.46
1	C	262	KCX	OQ1-CX	2.61	1.26	1.21
1	B	262	KCX	OQ1-CX	2.38	1.26	1.21
1	D	262	KCX	CE-NZ	2.16	1.51	1.46
1	A	262	KCX	OQ1-CX	2.00	1.25	1.21

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	262	KCX	OQ1-CX-NZ	-4.16	118.61	124.92
1	B	262	KCX	OQ1-CX-NZ	-3.58	119.48	124.92
1	A	262	KCX	OQ1-CX-NZ	-2.70	120.82	124.92
1	D	262	KCX	OQ1-CX-NZ	-2.31	121.42	124.92
1	B	262	KCX	CE-NZ-CX	-2.29	118.09	121.98

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	262	KCX	CG-CD-CE-NZ
1	A	262	KCX	CG-CD-CE-NZ

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 10 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	UAG	C	1535	3	59,60,60	2.35	7 (11%)	82,88,88	2.45	24 (29%)
4	UAG	D	1536	3	59,60,60	1.75	7 (11%)	82,88,88	2.86	25 (30%)
4	UAG	B	1536	3	59,60,60	1.55	6 (10%)	82,88,88	2.70	28 (34%)
2	ADP	A	1533	3	28,29,29	1.69	5 (17%)	43,45,45	1.87	10 (23%)
2	ADP	B	1534	3	28,29,29	1.60	6 (21%)	43,45,45	2.11	14 (32%)
4	UAG	A	1536	3	59,60,60	1.94	9 (15%)	82,88,88	2.92	39 (47%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ADP	C	1533	3	28,29,29	1.97	5 (17%)	43,45,45	1.87	10 (23%)
2	ADP	D	1534	3	28,29,29	1.82	5 (17%)	43,45,45	1.92	13 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	UAG	C	1535	3	-	10/55/92/92	0/3/3/3
4	UAG	D	1536	3	-	22/55/92/92	0/3/3/3
4	UAG	B	1536	3	-	12/55/92/92	0/3/3/3
2	ADP	A	1533	3	-	6/16/32/32	0/3/3/3
2	ADP	B	1534	3	-	7/16/32/32	0/3/3/3
4	UAG	A	1536	3	-	9/55/92/92	0/3/3/3
2	ADP	C	1533	3	-	5/16/32/32	0/3/3/3
2	ADP	D	1534	3	-	6/16/32/32	0/3/3/3

The worst 5 of 50 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	1535	UAG	PB-O3A	11.54	1.72	1.59
4	C	1535	UAG	PA-O3A	9.89	1.70	1.59
4	D	1536	UAG	PA-O3A	9.03	1.69	1.59
4	B	1536	UAG	PB-O3A	7.37	1.67	1.59
4	A	1536	UAG	PA-O3A	7.17	1.67	1.59

The worst 5 of 163 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	1536	UAG	O2B-PB-O3A	-13.15	71.74	107.27
4	C	1535	UAG	O2B-PB-O3A	-10.28	79.49	107.27
4	B	1536	UAG	O2B-PB-O3A	-9.40	81.86	107.27
4	A	1536	UAG	O3A-PB-O1B	-9.36	82.54	110.70
4	A	1536	UAG	O2B-PB-O3A	-8.64	83.91	107.27

There are no chirality outliers.

5 of 77 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1533	ADP	C5'-O5'-PA-O2A
2	A	1533	ADP	C5'-O5'-PA-O3A
2	B	1534	ADP	C5'-O5'-PA-O1A
2	B	1534	ADP	C5'-O5'-PA-O2A
2	B	1534	ADP	C5'-O5'-PA-O3A


There are no ring outliers.

8 monomers are involved in 30 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	1535	UAG	2	0
4	D	1536	UAG	7	0
4	B	1536	UAG	4	0
2	A	1533	ADP	1	0
2	B	1534	ADP	2	0
4	A	1536	UAG	5	0
2	C	1533	ADP	4	0
2	D	1534	ADP	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand UAG C 1535




Ligand UAG D 1536




Ligand UAG B 1536




Ligand ADP A 1533




Ligand ADP B 1534




Ligand UAG A 1536



Ligand ADP C 1533





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	501/535 (93%)	-0.34	0 100 100	25, 58, 104, 128	0
1	B	499/535 (93%)	-0.36	1 (0%) 91 83	24, 53, 99, 118	0
1	C	497/535 (92%)	-0.22	0 100 100	30, 65, 103, 137	0
1	D	487/535 (91%)	-0.24	1 (0%) 91 83	29, 60, 115, 155	0
All	All	1984/2140 (92%)	-0.29	2 (0%) 92 86	24, 59, 105, 155	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	57	THR	2.2
1	D	392	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	KCX	D	262	12/13	0.95	0.07	33,36,42,43	0
1	KCX	B	262	12/13	0.97	0.06	29,32,37,41	0
1	KCX	C	262	12/13	0.97	0.07	35,36,43,45	0
1	KCX	A	262	12/13	0.97	0.05	33,36,39,39	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MG	D	1537	1/1	0.81	0.25	53,53,53,53	0
3	MG	D	1533	1/1	0.86	0.15	50,50,50,50	0
3	MG	A	1535	1/1	0.88	0.20	37,37,37,37	0
3	MG	B	1537	1/1	0.89	0.19	39,39,39,39	0
3	MG	C	1536	1/1	0.89	0.14	43,43,43,43	0
3	MG	B	1533	1/1	0.90	0.17	30,30,30,30	0
3	MG	B	1535	1/1	0.90	0.10	46,46,46,46	0
2	ADP	A	1533	27/27	0.92	0.10	56,84,93,95	0
4	UAG	D	1536	58/58	0.93	0.08	48,56,69,74	0
4	UAG	A	1536	58/58	0.94	0.08	36,45,53,62	0
4	UAG	B	1536	58/58	0.94	0.07	39,45,52,53	0
4	UAG	C	1535	58/58	0.94	0.07	44,54,69,74	0
2	ADP	D	1534	27/27	0.94	0.07	46,55,64,72	0
2	ADP	C	1533	27/27	0.95	0.07	49,59,65,72	0
2	ADP	B	1534	27/27	0.95	0.08	41,60,69,71	0
3	MG	D	1535	1/1	0.95	0.06	51,51,51,51	0
3	MG	C	1534	1/1	0.95	0.08	60,60,60,60	0
3	MG	A	1534	1/1	0.97	0.07	48,48,48,48	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ADP A 1533:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around UAG D 1536:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around UAG A 1536:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around UAG B 1536:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around UAG C 1535:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ADP D 1534:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ADP C 1533:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ADP B 1534:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.