



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 12:45 PM UTC

PDB ID : 5XJM / pdb_00005xjm
Title : Complex structure of angiotensin II type 2 receptor with Fab
Authors : Asada, H.; Horita, S.; Shimamura, T.; Iwata, S.
Deposited on : 2017-05-02
Resolution : 3.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1466 (3.20-3.20)
Clashscore	190562	1573 (3.20-3.20)
Ramachandran outliers	187476	1548 (3.20-3.20)
Sidechain outliers	187428	1547 (3.20-3.20)
RSRZ outliers	180081	1466 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	19% 76% 13% 11%
2	H	220	2% 79% 21%
3	L	212	% 90% 10%
4	B	8	25% 88% 12%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6335 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Type-2 angiotensin II receptor,Soluble cytochrome b562,Type-2 angiotensin II receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	375	Total	C	N	O	S	0	0	0
			2992	1980	479	511	22			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	34	GLY	-	expression tag	UNP P50052
A	1007	TRP	MET	engineered mutation	UNP P0ABE7
A	1102	ILE	HIS	engineered mutation	UNP P0ABE7
A	1106	LEU	ARG	engineered mutation	UNP P0ABE7
A	347	GLU	-	expression tag	UNP P50052
A	348	ASN	-	expression tag	UNP P50052
A	349	LEU	-	expression tag	UNP P50052
A	350	TYR	-	expression tag	UNP P50052
A	351	PHE	-	expression tag	UNP P50052
A	352	GLN	-	expression tag	UNP P50052

- Molecule 2 is a protein called FabH.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	220	Total	C	N	O	S	0	0	0
			1643	1046	265	323	9			

- Molecule 3 is a protein called FabL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	212	Total	C	N	O	S	0	0	0
			1631	1017	269	336	9			


- Molecule 4 is a protein called Sar1, Ile8-angiotensin II.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	B	8	Total	C	N	O	0	0	0
			69	46	13	10			


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Type-2 angiotensin II receptor,Soluble cytochrome b562,Type-2 angiotensin II receptor





- Molecule 2: FabH



- Molecule 3: FabL



- Molecule 4: Sar1, Ile8-angiotensin II



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	465.39Å 48.65Å 55.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.39 – 3.20 48.39 – 3.20	Depositor EDS
% Data completeness (in resolution range)	99.8 (48.39-3.20) 95.1 (48.39-3.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.40 (at 3.19Å)	Xtriage
Refinement program	PHENIX 1.11.1 _2575	Depositor
R, R_{free}	0.226 , 0.275 0.228 , 0.277	Depositor DCC
R_{free} test set	2000 reflections (9.06%)	wwPDB-VP
Wilson B-factor (Å ²)	44.1	Xtriage
Anisotropy	0.489	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 58.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	6335	wwPDB-VP
Average B, all atoms (Å ²)	69.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SAR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.10	0/3069	0.24	0/4176
2	H	0.10	0/1690	0.34	0/2311
3	L	0.09	0/1669	0.30	0/2269
4	B	0.14	0/66	0.38	0/88
All	All	0.10	0/6494	0.29	0/8844

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2992	0	3053	33	0
2	H	1643	0	1592	27	0
3	L	1631	0	1561	12	0
4	B	69	0	73	1	0
All	All	6335	0	6279	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:266:ILE:O	1:A:270:LEU:HB2	1.91	0.69
2:H:22:CYS:HB3	2:H:79:LEU:HB3	1.75	0.69
2:H:29:PHE:O	2:H:72:ARG:NH2	2.32	0.63
3:L:194:GLU:HG2	3:L:205:VAL:HG22	1.82	0.61
3:L:65:GLY:HA3	3:L:70:TYR:HA	1.84	0.60
2:H:91:THR:HG23	2:H:117:THR:HA	1.84	0.59
2:H:168:SER:HA	2:H:189:THR:O	2.01	0.59
1:A:310:ASN:O	1:A:314:ASN:HB2	2.03	0.59
1:A:94:LEU:HA	1:A:97:LEU:HG	1.86	0.57
3:L:189:ASN:ND2	3:L:211:ASN:OD1	2.37	0.57
2:H:159:VAL:HG22	2:H:204:VAL:HG22	1.86	0.56
1:A:1011:ASN:HA	1:A:1098:ARG:HH12	1.71	0.56
3:L:107:ARG:NH1	3:L:108:ALA:O	2.36	0.56
1:A:119:VAL:O	1:A:123:PHE:HB2	2.05	0.56
1:A:55:PHE:O	1:A:59:PHE:HB2	2.05	0.55
1:A:252:ASP:OD1	1:A:256:LYS:NZ	2.35	0.55
2:H:159:VAL:HA	2:H:203:SER:O	2.07	0.55
1:A:1011:ASN:HA	1:A:1098:ARG:NH1	2.23	0.54
2:H:175:ALA:HA	2:H:184:MET:HB3	1.89	0.54
1:A:55:PHE:O	1:A:59:PHE:CB	2.57	0.53
3:L:36:GLN:HB2	3:L:46:LEU:HD11	1.90	0.52
3:L:49:ASP:HB2	3:L:52:ASN:HD22	1.76	0.51
1:A:144:GLN:HB3	1:A:151:LEU:HD12	1.93	0.51
1:A:266:ILE:O	1:A:270:LEU:CB	2.57	0.51
3:L:33:TYR:HD1	3:L:48:TYR:HA	1.76	0.51
1:A:1045:PRO:HD2	1:A:1062:ARG:HH21	1.76	0.50
1:A:1061:PHE:HZ	1:A:246:LYS:HD3	1.77	0.50
3:L:159:LEU:O	3:L:176:SER:HA	2.11	0.50
1:A:41:ASP:OD2	1:A:107:ARG:NH2	2.45	0.49
2:H:150:LYS:NZ	2:H:178:GLN:OE1	2.46	0.49
1:A:268:CYS:HB3	1:A:310:ASN:HB2	1.93	0.49
2:H:51:ILE:HG13	2:H:58:ILE:HG12	1.94	0.49
1:A:284:MET:HG3	1:A:286:VAL:HG23	1.95	0.48
2:H:17:SER:HA	2:H:83:MET:O	2.14	0.48
1:A:1060:ASP:O	1:A:1101:TYR:OH	2.30	0.47
2:H:35:HIS:HB2	2:H:97:ALA:O	2.15	0.47
2:H:29:PHE:HB3	2:H:77:ASN:HD22	1.80	0.46
2:H:145:LEU:HD13	2:H:217:LEU:HD13	1.97	0.46
1:A:183:ASP:OD1	1:A:184:VAL:N	2.48	0.45
1:A:290:CYS:SG	1:A:291:GLU:N	2.90	0.45
1:A:293:ILE:HA	1:A:296:ILE:HD12	1.99	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:199:PHE:HB2	1:A:204:TYR:CD2	2.52	0.45
2:H:97:ALA:HB1	2:H:107:PHE:HB3	1.99	0.45
3:L:144:ASN:O	3:L:195:ALA:HA	2.17	0.45
2:H:163:SER:HB3	2:H:200:VAL:HG13	1.98	0.45
2:H:12:VAL:HG11	2:H:86:LEU:HD12	1.98	0.44
2:H:36:TRP:HD1	2:H:70:ILE:HD12	1.82	0.44
1:A:154:ARG:C	1:A:157:PRO:HD2	2.43	0.44
1:A:215:LYS:NZ	4:B:8:ILE:O	2.41	0.44
1:A:109:ASP:HA	1:A:192:VAL:HG13	1.99	0.44
1:A:58:GLY:C	1:A:94:LEU:HD13	2.42	0.43
1:A:1021:ASP:OD1	1:A:1088:GLN:NE2	2.38	0.43
1:A:72:GLN:O	1:A:74:GLY:N	2.44	0.43
1:A:1095:LYS:O	1:A:1099:ASN:HB2	2.18	0.43
1:A:1098:ARG:HG3	1:A:1099:ASN:N	2.33	0.43
2:H:191:PRO:HG2	2:H:194:THR:HG23	2.01	0.42
3:L:144:ASN:HB3	3:L:196:THR:OG1	2.19	0.42
2:H:154:PRO:HD2	2:H:208:ALA:HB1	2.02	0.42
2:H:126:PRO:HB2	2:H:149:VAL:HG13	2.01	0.42
3:L:107:ARG:HG2	3:L:108:ALA:N	2.35	0.42
2:H:51:ILE:HD13	2:H:72:ARG:HB2	2.02	0.41
2:H:83:MET:HE1	2:H:116:VAL:HG11	2.03	0.41
1:A:274:VAL:O	1:A:278:LEU:HG	2.20	0.41
2:H:195:TRP:HA	2:H:196:PRO:HA	1.88	0.41
1:A:164:VAL:HB	1:A:165:PRO:HD3	2.04	0.40
2:H:83:MET:HB3	2:H:86:LEU:HD21	2.03	0.40
2:H:161:TRP:CZ3	2:H:202:CYS:HB3	2.56	0.40
2:H:201:THR:HA	2:H:215:LYS:O	2.21	0.40
3:L:137:ASN:HD22	3:L:171:THR:HG21	1.85	0.40
2:H:169:SER:O	2:H:188:VAL:HA	2.20	0.40
1:A:1003:LEU:HG	1:A:1105:TYR:HE1	1.86	0.40
1:A:1061:PHE:CE2	1:A:1104:LYS:HB3	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	371/422 (88%)	354 (95%)	17 (5%)	0	100	100
2	H	218/220 (99%)	204 (94%)	14 (6%)	0	100	100
3	L	210/212 (99%)	201 (96%)	9 (4%)	0	100	100
4	B	6/8 (75%)	6 (100%)	0	0	100	100
All	All	805/862 (93%)	765 (95%)	40 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	322/366 (88%)	321 (100%)	1 (0%)	86	88
2	H	185/185 (100%)	185 (100%)	0	100	100
3	L	187/187 (100%)	187 (100%)	0	100	100
4	B	7/7 (100%)	7 (100%)	0	100	100
All	All	701/745 (94%)	700 (100%)	1 (0%)	88	91

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1102	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	37	GLN
1	A	85	ASN
1	A	1013	ASN
1	A	1103	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	310	ASN
2	H	77	ASN
4	B	6	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SAR	B	1	4	3,4,5	1.33	0	1,3,5	1.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SAR	B	1	4	-	1/1/2/3	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	1	SAR	C-CA-N-CN

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	375/422 (88%)	1.41	80 (21%) 2 2	61, 101, 143, 176	0
2	H	220/220 (100%)	0.21	5 (2%) 61 41	18, 42, 69, 98	0
3	L	212/212 (100%)	0.03	2 (0%) 81 64	14, 28, 49, 87	0
4	B	7/8 (87%)	1.71	2 (28%) 1 1	84, 86, 94, 98	0
All	All	814/862 (94%)	0.73	89 (10%) 10 7	14, 61, 137, 176	0

All (89) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1090	ALA	4.9
1	A	1098	ARG	4.2
1	A	156	ASN	4.1
1	A	1086	GLU	4.0
1	A	256	LYS	3.7
1	A	153	GLN	3.7
1	A	36	SER	3.6
1	A	162	TYR	3.6
1	A	1044	THR	3.6
1	A	150	PHE	3.5
1	A	157	PRO	3.5
1	A	1061	PHE	3.4
1	A	320	PHE	3.3
2	H	138	THR	3.3
1	A	142	ARG	3.3
1	A	1045	PRO	3.3
1	A	74	GLY	3.3
1	A	1081	GLU	3.2
1	A	154	ARG	3.2
1	A	255	LEU	3.2
1	A	151	LEU	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1023	ALA	3.2
1	A	35	CYS	3.1
1	A	1091	ALA	3.1
1	A	158	TRP	3.1
1	A	316	PHE	3.0
1	A	1024	ALA	2.9
1	A	311	SER	2.9
1	A	1026	VAL	2.9
1	A	1074	ASP	2.9
1	A	272	PHE	2.9
1	A	290	CYS	2.9
1	A	271	PRO	2.9
1	A	252	ASP	2.9
1	A	1073	ASP	2.8
1	A	76	LYS	2.8
1	A	268	CYS	2.8
1	A	1075	ALA	2.7
1	A	1063	HIS	2.7
1	A	75	PRO	2.7
1	A	132	ILE	2.7
1	A	1102	ILE	2.7
1	A	38	LYS	2.7
2	H	137	ASP	2.7
1	A	160	ALA	2.7
1	A	96	THR	2.6
1	A	1030	LEU	2.5
4	B	3	VAL	2.5
1	A	1087	ALA	2.5
1	A	1062	ARG	2.5
1	A	1071	GLN	2.5
1	A	152	SER	2.4
1	A	1017	ILE	2.4
1	A	288	ASN	2.4
1	A	247	ASN	2.4
1	A	303	ALA	2.4
1	A	1028	ASP	2.4
2	H	30	SER	2.4
1	A	283	TRP	2.4
1	A	1027	LYS	2.3
1	A	1077	LYS	2.3
1	A	116	MET	2.3
1	A	159	GLN	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	42	LYS	2.3
1	A	251	ARG	2.2
1	A	1099	ASN	2.2
1	A	315	PRO	2.2
1	A	285	GLY	2.2
1	A	1096	THR	2.2
1	A	1076	LEU	2.2
1	A	263	LEU	2.2
1	A	128	MET	2.2
1	A	170	MET	2.2
3	L	102	LYS	2.2
1	A	1069	VAL	2.1
2	H	167	SER	2.1
1	A	37	GLN	2.1
1	A	115	VAL	2.1
1	A	1083	LYS	2.1
1	A	1101	TYR	2.1
1	A	1019	LYS	2.1
3	L	168	LYS	2.1
1	A	248	ARG	2.0
1	A	223	PRO	2.0
1	A	1015	LYS	2.0
4	B	5	ILE	2.0
1	A	216	ASN	2.0
1	A	52	TYR	2.0
2	H	87	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	SAR	B	1	5/6	0.23	0.33	72,73,82,83	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.