



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 07:11 PM UTC

PDB ID : 6XJF / pdb_00006xjf
Title : X-ray crystal structure of Pyrococcus furiosus general transcription factor
TFE-alpha (SeMet labeled protein)
Authors : Murakami, K.S.; Jun, S.H.
Deposited on : 2020-06-23
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1466 (3.20-3.20)
Clashscore	190562	1573 (3.20-3.20)
Ramachandran outliers	187476	1548 (3.20-3.20)
Sidechain outliers	187428	1547 (3.20-3.20)
RSRZ outliers	180081	1466 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	110	
1	B	110	
1	C	110	
1	D	110	
1	E	110	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	110	68%21%5%6%
1	G	110	71%25%. .
1	H	110	60%29%5%6%

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 6810 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcription factor E.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	102	Total	C	N	O	Se	0	0	0
			842	531	147	160	4			
1	B	103	Total	C	N	O	Se	0	0	0
			850	535	149	162	4			
1	C	103	Total	C	N	O	Se	0	0	0
			850	535	149	162	4			
1	D	102	Total	C	N	O	Se	0	0	0
			842	531	147	160	4			
1	E	103	Total	C	N	O	Se	0	0	0
			850	535	149	162	4			
1	F	103	Total	C	N	O	Se	0	0	0
			850	535	149	162	4			
1	G	106	Total	C	N	O	Se	0	0	0
			876	551	154	167	4			
1	H	103	Total	C	N	O	Se	0	0	0
			850	535	149	162	4			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	13	MSE	ILE	conflict	UNP Q8U3H5
A	54	MSE	ILE	conflict	UNP Q8U3H5
A	66	MSE	PHE	conflict	UNP Q8U3H5
B	13	MSE	ILE	conflict	UNP Q8U3H5
B	54	MSE	ILE	conflict	UNP Q8U3H5
B	66	MSE	PHE	conflict	UNP Q8U3H5
C	13	MSE	ILE	conflict	UNP Q8U3H5
C	54	MSE	ILE	conflict	UNP Q8U3H5
C	66	MSE	PHE	conflict	UNP Q8U3H5
D	13	MSE	ILE	conflict	UNP Q8U3H5
D	54	MSE	ILE	conflict	UNP Q8U3H5
D	66	MSE	PHE	conflict	UNP Q8U3H5
E	13	MSE	ILE	conflict	UNP Q8U3H5


Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
E	54	MSE	ILE	conflict	UNP Q8U3H5
E	66	MSE	PHE	conflict	UNP Q8U3H5
F	13	MSE	ILE	conflict	UNP Q8U3H5
F	54	MSE	ILE	conflict	UNP Q8U3H5
F	66	MSE	PHE	conflict	UNP Q8U3H5
G	13	MSE	ILE	conflict	UNP Q8U3H5
G	54	MSE	ILE	conflict	UNP Q8U3H5
G	66	MSE	PHE	conflict	UNP Q8U3H5
H	13	MSE	ILE	conflict	UNP Q8U3H5
H	54	MSE	ILE	conflict	UNP Q8U3H5
H	66	MSE	PHE	conflict	UNP Q8U3H5




- Molecule 1: Transcription factor E



- Molecule 1: Transcription factor E



- Molecule 1: Transcription factor E



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	76.20Å 115.61Å 124.21Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.44 – 3.20 44.44 – 3.20	Depositor EDS
% Data completeness (in resolution range)	98.7 (44.44-3.20) 97.6 (44.44-3.20)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.16 (at 3.19Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, R_{free}	0.237 , 0.304 (Not available) , 0.307	Depositor DCC
R_{free} test set	1359 reflections (7.27%)	wwPDB-VP
Wilson B-factor (Å ²)	84.7	Xtriage
Anisotropy	0.331	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 7.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6810	wwPDB-VP
Average B, all atoms (Å ²)	82.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.55	0/848	1.11	3/1133 (0.3%)
1	B	0.61	0/856	1.12	0/1144
1	C	0.67	0/856	1.20	6/1144 (0.5%)
1	D	0.59	0/848	1.05	4/1133 (0.4%)
1	E	0.57	0/856	1.01	3/1144 (0.3%)
1	F	0.64	0/856	1.00	1/1144 (0.1%)
1	G	0.72	0/882	1.11	3/1177 (0.3%)
1	H	0.56	0/856	1.05	1/1144 (0.1%)
All	All	0.62	0/6858	1.08	21/9163 (0.2%)

There are no bond length outliers.

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	38	GLU	N-CA-CB	-8.07	98.37	110.07
1	G	84	ASP	CA-C-N	-6.80	110.08	122.38
1	G	84	ASP	C-N-CA	-6.80	110.08	122.38
1	D	84	ASP	CA-C-N	-6.66	110.33	122.38
1	D	84	ASP	C-N-CA	-6.66	110.33	122.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	842	0	861	34	0
1	B	850	0	867	23	0
1	C	850	0	867	18	0
1	D	842	0	861	25	0
1	E	850	0	867	21	0
1	F	850	0	867	23	0
1	G	876	0	897	15	1
1	H	850	0	867	34	1
All	All	6810	0	6954	174	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 174 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:65:THR:OG1	1:G:82:ARG:NH2	2.09	0.85
1:D:29:LEU:HD21	1:D:82:ARG:HA	1.64	0.80
1:B:36:THR:HG22	1:B:39:GLU:HG3	1.62	0.78
1:F:13:MSE:HE2	1:F:63:LEU:HD11	1.66	0.75
1:C:18:GLY:HA3	1:C:54:MSE:HE1	1.69	0.75

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:68:ARG:NH2	1:H:74:THR:O[4_534]	2.03	0.17

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	100/110 (91%)	99 (99%)	1 (1%)	0	100	100
1	B	101/110 (92%)	100 (99%)	1 (1%)	0	100	100
1	C	101/110 (92%)	98 (97%)	3 (3%)	0	100	100
1	D	100/110 (91%)	100 (100%)	0	0	100	100
1	E	101/110 (92%)	99 (98%)	2 (2%)	0	100	100
1	F	101/110 (92%)	98 (97%)	3 (3%)	0	100	100
1	G	104/110 (94%)	102 (98%)	2 (2%)	0	100	100
1	H	101/110 (92%)	101 (100%)	0	0	100	100
All	All	809/880 (92%)	797 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	88/90 (98%)	82 (93%)	6 (7%)	14	46
1	B	89/90 (99%)	86 (97%)	3 (3%)	32	64
1	C	89/90 (99%)	82 (92%)	7 (8%)	11	40
1	D	88/90 (98%)	82 (93%)	6 (7%)	14	46
1	E	89/90 (99%)	86 (97%)	3 (3%)	32	64
1	F	89/90 (99%)	84 (94%)	5 (6%)	19	52
1	G	92/90 (102%)	90 (98%)	2 (2%)	45	71
1	H	89/90 (99%)	84 (94%)	5 (6%)	19	52
All	All	713/720 (99%)	676 (95%)	37 (5%)	21	54

5 of 37 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	73	GLU
1	H	66	MSE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	F	78	TYR
1	H	7	ASN
1	C	37	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	F	108	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	98/110 (89%)	-0.18	0 100 100	49, 107, 134, 146	0
1	B	99/110 (90%)	-0.64	0 100 100	49, 69, 90, 94	0
1	C	99/110 (90%)	-0.46	0 100 100	50, 73, 148, 164	0
1	D	98/110 (89%)	-0.47	0 100 100	49, 75, 136, 150	0
1	E	99/110 (90%)	-0.63	0 100 100	56, 80, 105, 120	0
1	F	99/110 (90%)	-0.67	0 100 100	52, 70, 90, 107	0
1	G	102/110 (92%)	-0.45	0 100 100	46, 72, 105, 121	0
1	H	99/110 (90%)	-0.40	0 100 100	60, 96, 147, 167	0
All	All	793/880 (90%)	-0.49	0 100 100	46, 76, 126, 167	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.