



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 03:33 PM UTC

PDB ID : 6XJP / pdb_00006xjp
Title : Crystal Structure of KPT-185 bound to CRM1 (537-DLTVK-541 to GLCEQ)
Authors : Baumhardt, J.M.; Chook, Y.M.
Deposited on : 2020-06-24
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	216	
2	B	140	
3	C	1024	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 22206 atoms, of which 11142 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GTP-binding nuclear protein Ran.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	208	Total	C	H	N	O	S	0	5	0
			3392	1093	1698	291	304	6			

- Molecule 2 is a protein called Ran-specific GTPase-activating protein 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	123	Total	C	H	N	O	S	0	3	0
			2060	652	1031	181	191	5			

- Molecule 3 is a protein called Exportin-1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	1008	Total	C	H	N	O	S	0	42	0
			16684	5317	8401	1364	1557	45			

There are 45 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	-2	GLY	-	expression tag	UNP P30822
C	-1	GLY	-	expression tag	UNP P30822
C	0	SER	-	expression tag	UNP P30822
C	?	-	VAL	deletion	UNP P30822
C	?	-	GLN	deletion	UNP P30822
C	?	-	ARG	deletion	UNP P30822
C	?	-	LEU	deletion	UNP P30822
C	?	-	PRO	deletion	UNP P30822
C	?	-	ALA	deletion	UNP P30822
C	?	-	THR	deletion	UNP P30822
C	?	-	GLU	deletion	UNP P30822
C	?	-	MET	deletion	UNP P30822
C	?	-	SER	deletion	UNP P30822
C	?	-	PRO	deletion	UNP P30822

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	?	-	LEU	deletion	UNP P30822
C	?	-	ILE	deletion	UNP P30822
C	?	-	GLN	deletion	UNP P30822
C	?	-	LEU	deletion	UNP P30822
C	?	-	SER	deletion	UNP P30822
C	?	-	VAL	deletion	UNP P30822
C	?	-	GLY	deletion	UNP P30822
C	?	-	SER	deletion	UNP P30822
C	?	-	GLN	deletion	UNP P30822
C	?	-	ALA	deletion	UNP P30822
C	?	-	ILE	deletion	UNP P30822
C	?	-	SER	deletion	UNP P30822
C	?	-	THR	deletion	UNP P30822
C	?	-	GLY	deletion	UNP P30822
C	?	-	SER	deletion	UNP P30822
C	?	-	GLY	deletion	UNP P30822
C	?	-	ALA	deletion	UNP P30822
C	?	-	LEU	deletion	UNP P30822
C	?	-	ASN	deletion	UNP P30822
C	?	-	PRO	deletion	UNP P30822
C	?	-	GLU	deletion	UNP P30822
C	?	-	TYR	deletion	UNP P30822
C	?	-	MET	deletion	UNP P30822
C	?	-	LYS	deletion	UNP P30822
C	?	-	ARG	deletion	UNP P30822
C	?	-	PHE	deletion	UNP P30822
C	537	GLY	ASP	engineered mutation	UNP P30822
C	539	CYS	THR	engineered mutation	UNP P30822
C	540	GLU	VAL	engineered mutation	UNP P30822
C	541	GLN	LYS	engineered mutation	UNP P30822
C	1022	CYS	TYR	conflict	UNP P30822

- Molecule 4 is PHOSPHOAMINOPHOSPHONIC ACID-GUANYLATE ESTER (CCD ID: GNP) (formula: $C_{10}H_{17}N_6O_{13}P_3$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	H	N	O	P	
			44	10	12	6	13	3	

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mg	0	0
			1	1		

- Molecule 6 is propan-2-yl 3-{3-[3-methoxy-5-(trifluoromethyl)phenyl]-1H-1,2,4-triazol-1-yl}propanoate (CCD ID: K85) (formula: C₁₆H₁₈F₃N₃O₃) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	C	1	Total	C	F	N	O	0	0
			25	16	3	3	3		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: GTP-binding nuclear protein Ran



- Molecule 2: Ran-specific GTPase-activating protein 1



- Molecule 3: Exportin-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	105.95Å 105.95Å 305.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.99 – 2.80 46.99 – 2.80	Depositor EDS
% Data completeness (in resolution range)	83.1 (46.99-2.80) 83.1 (46.99-2.80)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.26 (at 2.81Å)	Xtriage
Refinement program	PHENIX 1.11.1_2575	Depositor
R, R_{free}	0.215 , 0.260 0.209 , 0.247	Depositor DCC
R_{free} test set	2000 reflections (5.49%)	wwPDB-VP
Wilson B-factor (Å ²)	41.6	Xtriage
Anisotropy	0.154	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 35.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	22206	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GNP, MG, K85

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.14	0/1754	0.36	0/2379
2	B	0.13	0/1059	0.31	0/1413
3	C	0.14	0/8579	0.31	0/11618
All	All	0.14	0/11392	0.31	0/15410

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1694	1698	1677	6	0
2	B	1029	1031	1021	5	0
3	C	8283	8401	8233	33	0
4	A	32	12	12	0	0
5	A	1	0	0	0	0
6	C	25	0	17	3	0
All	All	11064	11142	10960	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:534:ASP:N	3:C:534:ASP:OD1	2.21	0.73
1:A:106:ARG:NH1	1:A:107:ASP:OD1	2.27	0.67
3:C:499:GLU:N	3:C:499:GLU:OE1	2.30	0.65
3:C:583:PHE:CE1	6:C:1101:K85:H15	2.33	0.62
3:C:453:ARG:O	3:C:453:ARG:NE	2.33	0.61
3:C:440:GLU:OE2	3:C:440:GLU:N	2.34	0.60
3:C:544:GLY:O	3:C:546:ASP:N	2.35	0.59
3:C:533:LYS:HG3	3:C:534:ASP:N	2.18	0.57
3:C:453:ARG:HE	3:C:453:ARG:C	2.14	0.55
2:B:143:GLU:OE2	2:B:143:GLU:N	2.39	0.55
3:C:542:LYS:O	3:C:548:LYS:HE2	2.08	0.53
3:C:583:PHE:CD1	6:C:1101:K85:H15	2.44	0.53
3:C:748:LEU:HD21	3:C:803:VAL:HG11	1.91	0.52
3:C:586:GLU:CD	6:C:1101:K85:H14	2.35	0.52
3:C:669:MET:HE2	3:C:669:MET:HA	1.93	0.51
3:C:205[A]:SER:OG	3:C:206[A]:SER:N	2.44	0.51
3:C:465:TYR:OH	3:C:557:TYR:OH	2.27	0.48
3:C:800:MET:HE2	3:C:836:SER:HA	1.96	0.47
3:C:79:TRP:NE1	3:C:129:GLU:OE2	2.37	0.47
1:A:23:LYS:HA	1:A:89:MET:CE	2.45	0.47
3:C:442:LEU:O	3:C:453:ARG:HD2	2.16	0.46
3:C:270:LEU:O	3:C:274:GLN:HG3	2.14	0.46
3:C:615:PRO:C	3:C:616:ARG:HG2	2.41	0.46
3:C:327:ARG:NH2	3:C:368:ASN:OD1	2.46	0.45
3:C:536:LEU:C	3:C:538:LEU:H	2.24	0.45
2:B:101:LYS:HG3	2:B:101:LYS:O	2.17	0.45
3:C:679:ILE:HD11	3:C:695:THR:HG23	1.99	0.45
3:C:453:ARG:NE	3:C:453:ARG:C	2.75	0.44
2:B:126:MET:HG2	2:B:135:CYS:SG	2.58	0.43
1:A:85:CYS:HB2	1:A:164:LEU:HD22	2.00	0.43
3:C:278:PHE:CD1	3:C:278:PHE:C	2.96	0.43
3:C:439:GLU:HG2	3:C:462:ILE:HG12	2.01	0.43
3:C:55:LEU:HD22	3:C:97:MET:HE1	2.01	0.43
1:A:158:GLU:HG2	1:A:159:LYS:HG2	2.01	0.42
3:C:500:TRP:CE2	3:C:542:LYS:HE2	2.55	0.42
3:C:240:THR:OG1	3:C:241:LYS:N	2.53	0.42
3:C:542:LYS:C	3:C:543:ARG:HG3	2.44	0.42
3:C:99:ILE:HG13	3:C:100:SER:N	2.35	0.41
1:A:185:PRO:HG2	2:B:162:ALA:HB1	2.02	0.41
1:A:181:ALA:H	2:B:79:MET:HB2	1.86	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:452:VAL:HG22	3:C:453:ARG:N	2.35	0.40
3:C:51:ALA:O	3:C:55:LEU:HG	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	211/216 (98%)	203 (96%)	8 (4%)	0	100	100
2	B	124/140 (89%)	116 (94%)	8 (6%)	0	100	100
3	C	1042/1024 (102%)	993 (95%)	42 (4%)	7 (1%)	18	47
All	All	1377/1380 (100%)	1312 (95%)	58 (4%)	7 (0%)	24	55

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	9	ASN
3	C	205[A]	SER
3	C	205[B]	SER
3	C	545	LYS
3	C	684	THR
3	C	982	PRO
3	C	265	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/185 (100%)	183 (99%)	2 (1%)	65	87
2	B	108/121 (89%)	107 (99%)	1 (1%)	70	89
3	C	957/933 (103%)	933 (98%)	24 (2%)	42	76
All	All	1250/1239 (101%)	1223 (98%)	27 (2%)	48	78

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	24	THR
1	A	197	TYR
2	B	125	LEU
3	C	9	ASN
3	C	147[A]	SER
3	C	147[B]	SER
3	C	182	LEU
3	C	215[A]	GLU
3	C	215[B]	GLU
3	C	264	ILE
3	C	270	LEU
3	C	428	ARG
3	C	499	GLU
3	C	530	THR
3	C	533	LYS
3	C	534	ASP
3	C	538	LEU
3	C	539	CYS
3	C	541	GLN
3	C	543	ARG
3	C	629	ILE
3	C	658	VAL
3	C	695	THR
3	C	798[A]	ASP
3	C	798[B]	ASP
3	C	803	VAL
3	C	953	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	30	HIS
3	C	274	GLN
3	C	494	GLN
3	C	506	ASN
3	C	742	GLN
3	C	813	ASN
3	C	928	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GNP	A	301	5	34,34,34	3.70	16 (47%)	47,54,54	1.66	8 (17%)
6	K85	C	1101	3	26,26,26	1.34	3 (11%)	36,37,37	2.17	8 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GNP	A	301	5	-	7/18/38/38	0/3/3/3
6	K85	C	1101	3	-	3/21/21/21	0/2/2/2

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	301	GNP	C2-N2	8.77	1.54	1.34
4	A	301	GNP	O4'-C1'	8.53	1.61	1.42
4	A	301	GNP	PA-O3A	7.41	1.67	1.59
4	A	301	GNP	PB-O3A	7.02	1.67	1.59
4	A	301	GNP	C2'-C1'	-7.01	1.31	1.53
4	A	301	GNP	PB-O1B	5.79	1.54	1.46
4	A	301	GNP	O4'-C4'	-5.36	1.33	1.45
4	A	301	GNP	PG-O1G	4.02	1.52	1.46
4	A	301	GNP	O2'-C2'	3.97	1.52	1.43
6	C	1101	K85	C9-C6	3.53	1.53	1.47
4	A	301	GNP	C6-N1	-3.25	1.32	1.38
6	C	1101	K85	O1'-C2'	-3.03	1.40	1.47
4	A	301	GNP	O3'-C3'	-2.89	1.35	1.43
4	A	301	GNP	PB-N3B	2.67	1.70	1.63
4	A	301	GNP	O6-C6	-2.59	1.18	1.23
4	A	301	GNP	PA-O5'	2.49	1.69	1.59
4	A	301	GNP	PG-N3B	2.37	1.69	1.63
6	C	1101	K85	O1'-C1	2.12	1.40	1.34
4	A	301	GNP	PB-O2B	-2.03	1.51	1.56

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	C	1101	K85	C3-N4-N5	6.24	125.87	119.56
4	A	301	GNP	C5-C4-N3	-5.31	119.94	128.39
6	C	1101	K85	C8-N7-C6	5.07	107.07	102.71
6	C	1101	K85	N4-C8-N7	-4.96	108.17	110.82
4	A	301	GNP	C2-N3-C4	4.46	119.99	112.30
6	C	1101	K85	O1'-C1-C2	4.13	120.41	111.48
6	C	1101	K85	C8-N4-N5	-3.89	108.04	109.97
6	C	1101	K85	N5-C6-N7	-3.44	107.62	114.62
4	A	301	GNP	N9-C4-N3	2.94	131.84	125.95
4	A	301	GNP	N9-C8-N7	-2.80	108.21	113.40
4	A	301	GNP	C2-N1-C6	-2.78	120.06	125.11
4	A	301	GNP	C5-C6-N1	2.61	119.89	113.25
4	A	301	GNP	O6-C6-C5	-2.49	119.97	126.53
4	A	301	GNP	C8-N7-C5	2.04	107.89	104.26

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	C	1101	K85	C9-C6-N7	2.01	126.43	122.92
6	C	1101	K85	C2'-O1'-C1	-2.00	113.68	117.45

There are no chirality outliers.

All (10) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	301	GNP	PG-N3B-PB-O1B
4	A	301	GNP	PG-N3B-PB-O3A
6	C	1101	K85	C1-C2-C3-N4
6	C	1101	K85	C2-C1-O1'-C2'
4	A	301	GNP	O4'-C4'-C5'-O5'
6	C	1101	K85	O-C1-O1'-C2'
4	A	301	GNP	C3'-C4'-C5'-O5'
4	A	301	GNP	C5'-O5'-PA-O3A
4	A	301	GNP	C5'-O5'-PA-O1A
4	A	301	GNP	C5'-O5'-PA-O2A

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	C	1101	K85	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å²)	Q<0.9
1	A	208/216 (96%)	0.28	18 (8%)	16	11	17, 41, 140, 185	3 (1%)
2	B	123/140 (87%)	0.42	6 (4%)	35	27	17, 62, 102, 166	2 (1%)
3	C	1008/1024 (98%)	0.31	52 (5%)	33	25	11, 50, 99, 154	26 (2%)
All	All	1339/1380 (97%)	0.32	76 (5%)	29	22	11, 50, 104, 185	31 (2%)

All (76) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	446	ASN	9.8
3	C	206[A]	SER	8.8
3	C	205[A]	SER	7.5
3	C	39[A]	THR	6.8
3	C	-1	GLY	6.7
3	C	450	GLU	5.8
3	C	690	LEU	5.5
3	C	451	ILE	5.3
1	A	191	PRO	5.3
1	A	189	MET	5.2
3	C	270	LEU	5.2
3	C	691	LEU	4.3
2	B	78	THR	4.2
3	C	688	THR	4.1
1	A	188	VAL	4.1
3	C	445	GLU	4.0
3	C	687	PRO	4.0
1	A	197	TYR	3.9
3	C	981	VAL	3.9
1	A	193	LEU	3.9
3	C	0	SER	3.6
1	A	187	VAL	3.4
3	C	262	LEU	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	C	689	LEU	3.4
3	C	207[A]	SER	3.4
3	C	3	GLY	3.4
2	B	200	LYS	3.4
3	C	192[A]	GLN	3.3
1	A	194	ALA	3.3
3	C	452	VAL	3.2
1	A	179	MET	3.2
3	C	197[A]	CYS	3.1
1	A	199	HIS	3.1
1	A	195	ALA	3.1
3	C	204[A]	GLY	3.1
3	C	1052	ALA	3.0
3	C	542	LYS	2.9
3	C	265	PRO	2.9
3	C	674	MET	2.8
3	C	133	ASN	2.8
3	C	72[A]	ASP	2.8
3	C	1	MET	2.8
3	C	453	ARG	2.7
1	A	9	VAL	2.7
1	A	192	ALA	2.7
2	B	199	LYS	2.6
3	C	5	LEU	2.6
3	C	266	GLN	2.6
3	C	530	THR	2.6
1	A	180	PRO	2.6
3	C	498	SER	2.5
1	A	113	GLU	2.5
3	C	549	ALA	2.5
3	C	10	ASP	2.4
1	A	182	LEU	2.4
3	C	611	VAL	2.4
3	C	572	PHE	2.3
3	C	291[A]	MET	2.3
1	A	186	GLU	2.3
3	C	271	ILE	2.3
3	C	264	ILE	2.3
3	C	684	THR	2.2
3	C	982	PRO	2.2
1	A	216	LEU	2.2
2	B	79	MET	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	129	ASP	2.1
3	C	654	GLU	2.1
1	A	134	LYS	2.1
3	C	570	TRP	2.1
2	B	80	GLU	2.1
3	C	83	PRO	2.0
3	C	93	PHE	2.0
3	C	457	LYS	2.0
3	C	967	TYR	2.0
3	C	58	SER	2.0
3	C	238	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	MG	A	302	1/1	0.77	0.10	15,15,15,15	0
6	K85	C	1101	25/25	0.83	0.20	91,92,93,96	0
4	GNP	A	301	32/32	0.90	0.10	0,5,23,24	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around K85 C 1101:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around GNP A 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.