



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 06:09 AM UTC

PDB ID : 8XJJ / pdb_00008xjj
Title : Co-crystal structure of SOS-1 and a potent, selective and orally bioavailable SOS1 inhibitor RGT-018
Authors : Ren, X.
Deposited on : 2023-12-21
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	590	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	A	1213	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4109 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Son of sevenless homolog 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	473	3925	2515	675	721	14	0	1	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	560	GLY	-	engineered mutation	UNP Q07889
A	561	ALA	-	engineered mutation	UNP Q07889
A	562	MET	-	engineered mutation	UNP Q07889
A	563	ALA	-	engineered mutation	UNP Q07889

- Molecule 2 is 5-[4-[(1 {R})-1-[3-[bis(fluoranyl)methyl]-2-fluoranyl-phenyl]ethyl]amino]-2-methyl-6-morpholin-4-yl-7-oxidanylidene-pyrido[4,3-d]pyrimidin-8-yl]pyridine-2-carbonitrile (CCD ID: A1LVJ) (formula: C₂₇H₂₄F₃N₇O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	0	0
			39	27	3	7	2		

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	97	Total	O	0	0
			97	97		

- Molecule 1: Son of sevenless homolog 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	40.76Å 83.85Å 179.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.80 – 2.10 44.80 – 2.10	Depositor EDS
% Data completeness (in resolution range)	100.0 (44.80-2.10) 100.0 (44.80-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.39 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.220 , 0.264 0.228 , 0.271	Depositor DCC
R_{free} test set	1855 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	52.7	Xtriage
Anisotropy	0.123	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 38.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4109	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A1LVJ, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.01	0/4020	1.42	2/5436 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	629	THR	CA-CB-OG1	-5.82	100.87	109.60
1	A	882	PRO	CB-CA-C	-5.12	104.63	112.11

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3925	0	3941	32	0
2	A	39	0	0	0	0
3	A	48	0	72	9	0
4	A	97	0	0	3	0
All	All	4109	0	4013	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:605:THR:HB	3:A:1212:EDO:H22	1.67	0.75
1:A:656:THR:H	3:A:1213:EDO:H22	1.52	0.74
1:A:671:SER:HA	3:A:1213:EDO:H12	1.72	0.70
1:A:605:THR:HB	3:A:1212:EDO:C2	2.23	0.68
1:A:683:GLN:HB3	1:A:684:PRO:HD3	1.82	0.62
1:A:1007:ASP:HB2	4:A:1337:HOH:O	2.00	0.61
1:A:1020:ASN:HB3	1:A:1021:PRO:HD3	1.86	0.58
1:A:884:TYR:O	3:A:1207:EDO:C2	2.54	0.55
1:A:656:THR:HG23	3:A:1213:EDO:H22	1.89	0.54
1:A:741:LYS:O	1:A:744:ARG:NH1	2.41	0.54
1:A:1037:SER:HB2	3:A:1203:EDO:H11	1.93	0.51
1:A:856:ILE:HG12	1:A:901:LEU:HD13	1.93	0.51
1:A:561:ALA:O	1:A:564:GLU:HB2	2.12	0.49
1:A:671:SER:HA	3:A:1213:EDO:C1	2.44	0.47
1:A:726:MET:O	1:A:730:VAL:HG23	2.15	0.47
1:A:859:LEU:C	1:A:859:LEU:HD23	2.41	0.45
1:A:636:LYS:HB2	1:A:639:GLU:HG3	1.98	0.45
1:A:1042:PRO:O	1:A:1043:SER:O	2.35	0.44
1:A:767:ARG:O	1:A:768:PRO:C	2.60	0.44
1:A:881:SER:OG	1:A:882:PRO:HD3	2.18	0.44
1:A:764:HIS:CE1	1:A:1038:PRO:HD3	2.53	0.43
1:A:606:VAL:O	1:A:610:ILE:HG12	2.19	0.43
1:A:589:GLU:HG2	1:A:602:LYS:HB2	2.00	0.43
1:A:622:ASN:OD1	1:A:625:ARG:NH2	2.51	0.42
1:A:1037:SER:CB	3:A:1203:EDO:H11	2.50	0.42
1:A:695:HIS:HE1	4:A:1353:HOH:O	2.02	0.41
1:A:821:LEU:CD1	1:A:930:PHE:HE2	2.33	0.41
1:A:1020:ASN:O	1:A:1021:PRO:C	2.63	0.41
1:A:615:TYR:O	1:A:688:ARG:NH1	2.51	0.41
1:A:620:ASP:OD1	4:A:1301:HOH:O	2.22	0.41
1:A:655:PRO:HG2	1:A:660:ARG:HG2	2.02	0.40
1:A:941:GLU:OE2	1:A:964:VAL:HG21	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	468/590 (79%)	445 (95%)	20 (4%)	3 (1%)	21	18

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1043	SER
1	A	743	ALA
1	A	924	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	440/544 (81%)	424 (96%)	16 (4%)	31	34

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	562	MET
1	A	579	GLU
1	A	660	ARG
1	A	724	LYS
1	A	726	MET
1	A	742	ILE
1	A	745	ASP
1	A	772	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	816	ILE
1	A	895	SER
1	A	919	LEU
1	A	1000	SER
1	A	1003	LYS
1	A	1004	GLU
1	A	1012	LYS
1	A	1029	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	665	ASN
1	A	691	ASN
1	A	764	HIS
1	A	860	GLN
1	A	973	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	A	1204	-	3,3,3	0.14	0	2,2,2	0.28	0
3	EDO	A	1212	-	3,3,3	0.26	0	2,2,2	0.55	0
3	EDO	A	1210	-	3,3,3	0.13	0	2,2,2	0.30	0
3	EDO	A	1207	-	3,3,3	0.47	0	2,2,2	0.47	0
3	EDO	A	1211	-	3,3,3	0.23	0	2,2,2	0.23	0
3	EDO	A	1202	-	3,3,3	0.10	0	2,2,2	0.23	0
3	EDO	A	1206	-	3,3,3	0.13	0	2,2,2	0.21	0
3	EDO	A	1205	-	3,3,3	0.06	0	2,2,2	0.29	0
3	EDO	A	1208	-	3,3,3	0.16	0	2,2,2	0.19	0
3	EDO	A	1209	-	3,3,3	0.10	0	2,2,2	0.09	0
3	EDO	A	1213	-	3,3,3	0.40	0	2,2,2	0.17	0
2	A1LVJ	A	1201	-	39,43,43	3.48	13 (33%)	45,62,62	2.68	20 (44%)
3	EDO	A	1203	-	3,3,3	0.17	0	2,2,2	0.16	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	1204	-	-	1/1/1/1	-
3	EDO	A	1212	-	-	1/1/1/1	-
3	EDO	A	1210	-	-	1/1/1/1	-
3	EDO	A	1207	-	-	1/1/1/1	-
3	EDO	A	1211	-	-	1/1/1/1	-
3	EDO	A	1202	-	-	1/1/1/1	-
3	EDO	A	1206	-	-	1/1/1/1	-
3	EDO	A	1205	-	-	0/1/1/1	-
3	EDO	A	1208	-	-	0/1/1/1	-
3	EDO	A	1209	-	-	1/1/1/1	-
3	EDO	A	1213	-	-	0/1/1/1	-
2	A1LVJ	A	1201	-	-	3/16/30/30	0/5/5/5
3	EDO	A	1203	-	-	1/1/1/1	-

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1201	A1LVJ	N13-N16	-14.20	1.28	1.41
2	A	1201	A1LVJ	C10-C09	9.52	1.53	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1201	A1LVJ	C21-N16	-6.71	1.38	1.47
2	A	1201	A1LVJ	C17-N16	-6.49	1.38	1.47
2	A	1201	A1LVJ	C11-N13	4.06	1.45	1.39
2	A	1201	A1LVJ	C06-N05	3.26	1.39	1.34
2	A	1201	A1LVJ	C14-N13	2.80	1.45	1.37
2	A	1201	A1LVJ	C25-C28	2.67	1.52	1.45
2	A	1201	A1LVJ	C34-C38	2.65	1.41	1.38
2	A	1201	A1LVJ	C07-C06	2.49	1.56	1.49
2	A	1201	A1LVJ	O12-C11	-2.43	1.18	1.23
2	A	1201	A1LVJ	F39-C38	2.17	1.38	1.34
2	A	1201	A1LVJ	C06-N08	2.08	1.37	1.34

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1201	A1LVJ	F36-C35-C34	6.78	117.83	110.64
2	A	1201	A1LVJ	C18-C17-N16	5.57	113.58	107.95
2	A	1201	A1LVJ	C20-C21-N16	5.48	113.49	107.95
2	A	1201	A1LVJ	F39-C38-C34	4.72	124.36	119.67
2	A	1201	A1LVJ	O12-C11-C10	-4.70	115.11	125.16
2	A	1201	A1LVJ	C31-C30-C38	4.60	121.83	116.69
2	A	1201	A1LVJ	C24-C23-C22	-4.29	116.22	120.80
2	A	1201	A1LVJ	C23-C22-C27	3.83	121.88	117.61
2	A	1201	A1LVJ	C33-C34-C38	3.76	120.90	116.69
2	A	1201	A1LVJ	N05-C06-N08	-3.60	119.41	125.77
2	A	1201	A1LVJ	C22-C10-C09	-3.59	118.96	123.45
2	A	1201	A1LVJ	C04-C15-C09	3.51	118.22	115.87
2	A	1201	A1LVJ	C21-N16-N13	2.71	118.30	113.20
2	A	1201	A1LVJ	O12-C11-N13	2.41	124.80	120.59
2	A	1201	A1LVJ	C22-C27-N26	-2.29	120.22	123.59
2	A	1201	A1LVJ	C38-C34-C35	2.27	127.49	121.30
2	A	1201	A1LVJ	C07-C06-N05	2.15	120.34	117.13
2	A	1201	A1LVJ	C27-N26-C25	2.09	120.03	116.73
2	A	1201	A1LVJ	F37-C35-F36	2.08	115.66	104.78
2	A	1201	A1LVJ	C07-C06-N08	2.08	120.24	117.13

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1201	A1LVJ	C38-C34-C35-F36
3	A	1204	EDO	O1-C1-C2-O2

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	A	1211	EDO	O1-C1-C2-O2
3	A	1207	EDO	O1-C1-C2-O2
3	A	1212	EDO	O1-C1-C2-O2
2	A	1201	A1LVJ	C30-C02-N03-C04
3	A	1202	EDO	O1-C1-C2-O2
3	A	1206	EDO	O1-C1-C2-O2
3	A	1209	EDO	O1-C1-C2-O2
3	A	1203	EDO	O1-C1-C2-O2
3	A	1210	EDO	O1-C1-C2-O2
2	A	1201	A1LVJ	C33-C34-C35-F37

There are no ring outliers.

4 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1212	EDO	2	0
3	A	1207	EDO	1	0
3	A	1213	EDO	4	0
3	A	1203	EDO	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	473/590 (80%)	0.42	20 (4%)	40 42	42, 59, 98, 158	1 (0%)

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	567	MET	3.9
1	A	563	ALA	3.4
1	A	561	ALA	3.4
1	A	753	THR	3.3
1	A	754	PHE	3.2
1	A	763	TRP	3.1
1	A	1001	MET	3.1
1	A	1023	PRO	2.8
1	A	595	LYS	2.7
1	A	566	GLN	2.5
1	A	598	ILE	2.5
1	A	596	ALA	2.4
1	A	742	ILE	2.3
1	A	1020	ASN	2.2
1	A	761	VAL	2.2
1	A	729	TRP	2.2
1	A	1003	LYS	2.1
1	A	1021	PRO	2.1
1	A	579	GLU	2.1
1	A	590	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	EDO	A	1212	4/4	0.75	0.16	68,73,74,74	0
3	EDO	A	1210	4/4	0.78	0.18	78,91,92,93	0
3	EDO	A	1206	4/4	0.78	0.13	78,80,84,85	0
3	EDO	A	1207	4/4	0.83	0.17	56,60,67,74	0
3	EDO	A	1204	4/4	0.85	0.14	68,79,80,81	0
3	EDO	A	1203	4/4	0.86	0.11	66,73,79,81	0
3	EDO	A	1213	4/4	0.87	0.18	52,54,56,66	0
3	EDO	A	1209	4/4	0.88	0.16	74,75,88,90	0
3	EDO	A	1211	4/4	0.88	0.16	60,72,73,74	0
3	EDO	A	1208	4/4	0.89	0.14	71,71,76,78	0
3	EDO	A	1202	4/4	0.91	0.13	59,59,60,64	0
3	EDO	A	1205	4/4	0.91	0.14	77,78,79,93	0
2	A1LVJ	A	1201	39/39	0.95	0.07	43,50,68,69	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around A1LVJ A 1201:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.