



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:20 PM UTC

PDB ID : 1XKP / pdb_00001xkp
Title : Crystal structure of the virulence factor YopN in complex with its heterodimeric chaperone SycN-YscB
Authors : Schubot, F.D.; Jackson, M.W.; Penrose, K.J.; Cherry, S.; Tropea, J.E.; Plano, G.V.; Waugh, D.S.
Deposited on : 2004-09-29
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	246	17% 78% 14% 6%
2	B	124	10% 88% 9% 2%
3	C	143	18% 79% 8% 12%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3952 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called putative membrane-bound Yop targeting protein YopN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	231	Total	C	N	O	S	0	0	0
			1823	1146	309	363	5			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	32	DMG	GLY	modified residue	GB 3822072
A	64	MLY	LYS	modified residue	GB 3822072
A	70	MLY	LYS	modified residue	GB 3822072
A	72	MLY	LYS	modified residue	GB 3822072
A	93	MLY	LYS	modified residue	GB 3822072
A	100	MLY	LYS	modified residue	GB 3822072
A	122	MLY	LYS	modified residue	GB 3822072
A	128	MLY	LYS	modified residue	GB 3822072
A	137	MLY	LYS	modified residue	GB 3822072
A	147	MLY	LYS	modified residue	GB 3822072
A	221	MLY	LYS	modified residue	GB 3822072
A	237	MLY	LYS	modified residue	GB 3822072
A	254	MLY	LYS	modified residue	GB 3822072
A	264	MLY	LYS	modified residue	GB 3822072
A	266	MLY	LYS	modified residue	GB 3822072
A	276	MLY	LYS	modified residue	GB 3822072

- Molecule 2 is a protein called Chaperone protein sycN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	121	Total	C	N	O	S	0	0	0
			945	608	161	172	4			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	65	MLY	LYS	modified residue	UNP P61380
B	74	MLY	LYS	modified residue	UNP P61380
B	123	PRO	SER	engineered mutation	UNP P61380

- Molecule 3 is a protein called Chaperone protein yscB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	126	Total	C	N	O	S	0	0	0
			1005	643	181	177	4			

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	6	MLY	LYS	modified residue	UNP Q56973
C	15	MLY	LYS	modified residue	UNP Q56973
C	21	MLY	LYS	modified residue	UNP Q56973
C	31	MLY	LYS	modified residue	UNP Q56973
C	113	MLY	LYS	modified residue	UNP Q56973
C	138	HIS	-	expression tag	UNP Q56973
C	139	HIS	-	expression tag	UNP Q56973
C	140	HIS	-	expression tag	UNP Q56973
C	141	HIS	-	expression tag	UNP Q56973
C	142	HIS	-	expression tag	UNP Q56973
C	143	HIS	-	expression tag	UNP Q56973


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	104	Total	O	0	0
			104	104		
4	B	35	Total	O	0	0
			35	35		
4	C	40	Total	O	0	0
			40	40		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: putative membrane-bound Yop targeting protein YopN



- Molecule 2: Chaperone protein sycN



- Molecule 3: Chaperone protein yscB



4 Data and refinement statistics

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, α , β , γ	60.72Å 60.72Å 140.42Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	60.86 – 1.70 60.72 – 1.70	Depositor EDS
% Data completeness (in resolution range)	98.0 (60.86-1.70) 89.9 (60.72-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.82 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.198 , 0.242 0.206 , 0.249	Depositor DCC
R_{free} test set	2574 reflections (4.62%)	wwPDB-VP
Wilson B-factor (Å ²)	30.0	Xtriage
Anisotropy	0.029	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 43.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.063 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3952	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: DMG, MLY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.77	0/1695	0.88	0/2297
2	B	0.77	0/943	0.80	0/1288
3	C	0.78	0/965	0.85	0/1316
All	All	0.77	0/3603	0.85	0/4901

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1823	0	1844	32	0
2	B	945	0	966	10	0
3	C	1005	0	1064	9	0
4	A	104	0	0	6	0
4	B	35	0	0	1	0
4	C	40	0	0	0	0
All	All	3952	0	3874	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:100:MLY:HH21	4:A:342:HOH:O	1.63	0.96
1:A:100:MLY:CH2	4:A:342:HOH:O	2.22	0.81
1:A:47:GLN:HG2	1:A:52:MET:HE3	1.66	0.76
1:A:100:MLY:NZ	4:A:342:HOH:O	2.20	0.74
1:A:100:MLY:HH11	4:A:342:HOH:O	1.90	0.70
1:A:100:MLY:CH1	4:A:342:HOH:O	2.42	0.67
1:A:157:HIS:CD2	1:A:161:GLN:HE21	2.15	0.65
1:A:264:MLY:HH11	4:A:331:HOH:O	1.97	0.65
1:A:148:GLY:O	1:A:149:ARG:HG3	1.99	0.62
1:A:196:GLN:NE2	1:A:199:ARG:HH21	1.98	0.62
1:A:166:MET:HE2	1:A:174:ILE:HD11	1.83	0.60
2:B:14:ASP:OD2	2:B:107:HIS:HE1	1.85	0.60
1:A:173:THR:HG23	1:A:203:ARG:HD2	1.85	0.58
3:C:122:ILE:O	3:C:126:THR:OG1	2.21	0.57
1:A:116:ILE:N	1:A:116:ILE:HD12	2.18	0.57
2:B:4:ILE:HD11	2:B:41:GLN:HG2	1.88	0.55
1:A:89:GLN:C	1:A:89:GLN:NE2	2.64	0.55
1:A:117:SER:H	1:A:120:GLN:HE21	1.55	0.54
2:B:67:LEU:CD1	3:C:65:LEU:HD11	2.38	0.54
1:A:94:VAL:HG13	1:A:94:VAL:O	2.09	0.53
1:A:52:MET:HA	1:A:52:MET:HE2	1.91	0.52
1:A:97:LEU:HD13	1:A:99:GLN:NE2	2.24	0.52
1:A:118:LEU:HG	1:A:122:MLY:HE2	1.93	0.51
3:C:2:GLN:HG2	3:C:5:LEU:H	1.75	0.51
1:A:89:GLN:C	1:A:89:GLN:HE21	2.19	0.51
2:B:97:ASP:OD2	2:B:99:ARG:NE	2.44	0.50
2:B:87:GLU:H	2:B:87:GLU:CD	2.21	0.49
1:A:46:LEU:HD22	2:B:28:GLN:OE1	2.13	0.49
1:A:94:VAL:O	1:A:96:GLU:N	2.46	0.49
1:A:111:SER:O	1:A:113:SER:N	2.46	0.48
1:A:196:GLN:HE22	1:A:199:ARG:HH21	1.60	0.48
1:A:72:MLY:HH22	3:C:30:ASP:OD1	2.14	0.47
3:C:5:LEU:HD13	3:C:25:TYR:CD2	2.50	0.47
1:A:114:PRO:HG3	3:C:58:ASN:HD21	1.79	0.46
2:B:87:GLU:CD	2:B:87:GLU:N	2.74	0.45
1:A:113:SER:HB2	1:A:116:ILE:HG13	1.99	0.45
1:A:194:PRO:HG2	1:A:197:PRO:HG2	1.99	0.45
3:C:80:PRO:HD2	3:C:81:GLN:OE1	2.17	0.44
3:C:15:MLY:HH23	3:C:15:MLY:HD3	1.90	0.44
2:B:108:GLN:HG2	4:B:142:HOH:O	2.17	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:148:GLY:O	1:A:149:ARG:CB	2.66	0.43
1:A:101:GLN:HA	1:A:101:GLN:OE1	2.19	0.43
1:A:116:ILE:N	1:A:116:ILE:CD1	2.82	0.43
1:A:148:GLY:O	1:A:149:ARG:CG	2.65	0.42
2:B:4:ILE:HD11	2:B:41:GLN:CG	2.50	0.42
2:B:67:LEU:HD11	3:C:65:LEU:HD11	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	214/246 (87%)	206 (96%)	4 (2%)	4 (2%)	6	1
2	B	117/124 (94%)	116 (99%)	1 (1%)	0	100	100
3	C	119/143 (83%)	118 (99%)	1 (1%)	0	100	100
All	All	450/513 (88%)	440 (98%)	6 (1%)	4 (1%)	14	4

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	112	ASN
1	A	149	ARG
1	A	109	LEU
1	A	95	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	188/200 (94%)	181 (96%)	7 (4%)	30	14
2	B	101/104 (97%)	99 (98%)	2 (2%)	48	32
3	C	105/120 (88%)	102 (97%)	3 (3%)	37	20
All	All	394/424 (93%)	382 (97%)	12 (3%)	36	19

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	45	THR
1	A	65	GLU
1	A	89	GLN
1	A	106	LEU
1	A	109	LEU
1	A	116	ILE
1	A	155	LEU
2	B	58	ARG
2	B	87	GLU
3	C	50	ASP
3	C	91	LEU
3	C	126	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	33	GLN
1	A	77	GLN
1	A	102	ASN
1	A	112	ASN
1	A	120	GLN
1	A	154	HIS
1	A	157	HIS
1	A	161	GLN
1	A	189	GLN
1	A	196	GLN
2	B	33	GLN
2	B	73	GLN
2	B	89	GLN
2	B	107	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	C	32	HIS
3	C	58	ASN
3	C	90	GLN
3	C	105	HIS
3	C	112	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

20 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MLY	B	65	2	9,10,11	0.48	0	6,11,13	1.06	1 (16%)
1	MLY	A	122	1	9,10,11	0.63	0	6,11,13	0.64	0
3	MLY	C	113	3	9,10,11	0.68	0	6,11,13	0.32	0
3	MLY	C	15	3	9,10,11	0.72	0	6,11,13	0.81	0
1	MLY	A	93	1	9,10,11	0.64	0	6,11,13	0.77	0
1	MLY	A	221	1	9,10,11	0.70	0	6,11,13	0.81	0
1	MLY	A	100	1	9,10,11	0.89	0	6,11,13	1.53	1 (16%)
3	MLY	C	6	3	9,10,11	0.67	0	6,11,13	0.85	0
1	MLY	A	266	1	9,10,11	0.60	0	6,11,13	0.79	0
1	MLY	A	128	1	9,10,11	0.50	0	6,11,13	0.71	0
3	MLY	C	21	3	9,10,11	0.64	0	6,11,13	0.73	0
3	MLY	C	31	3	9,10,11	0.59	0	6,11,13	0.66	0
1	MLY	A	70	1	9,10,11	0.71	0	6,11,13	0.75	0
1	MLY	A	147	1	9,10,11	0.64	0	6,11,13	0.76	0
2	MLY	B	74	2	9,10,11	0.71	0	6,11,13	0.36	0
1	MLY	A	237	1	9,10,11	0.60	0	6,11,13	1.11	0
1	MLY	A	264	1	9,10,11	0.66	0	6,11,13	0.82	0
1	MLY	A	254	1	9,10,11	0.64	0	6,11,13	0.60	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	MLY	A	72	1	9,10,11	0.79	0	6,11,13	1.08	0
1	MLY	A	137	1	9,10,11	0.70	0	6,11,13	0.92	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MLY	B	65	2	-	3/8/9/11	-
1	MLY	A	122	1	-	3/8/9/11	-
3	MLY	C	113	3	-	1/8/9/11	-
3	MLY	C	15	3	-	4/8/9/11	-
1	MLY	A	93	1	-	3/8/9/11	-
1	MLY	A	221	1	-	2/8/9/11	-
1	MLY	A	100	1	-	2/8/9/11	-
3	MLY	C	6	3	-	2/8/9/11	-
1	MLY	A	266	1	-	2/8/9/11	-
1	MLY	A	128	1	-	0/8/9/11	-
3	MLY	C	21	3	-	3/8/9/11	-
3	MLY	C	31	3	-	2/8/9/11	-
1	MLY	A	70	1	-	2/8/9/11	-
1	MLY	A	147	1	-	5/8/9/11	-
2	MLY	B	74	2	-	2/8/9/11	-
1	MLY	A	237	1	-	0/8/9/11	-
1	MLY	A	264	1	-	1/8/9/11	-
1	MLY	A	254	1	-	1/8/9/11	-
1	MLY	A	72	1	-	4/8/9/11	-
1	MLY	A	137	1	-	1/8/9/11	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	100	MLY	CH2-NZ-CH1	2.75	116.76	109.72
2	B	65	MLY	CD-CE-NZ	-2.07	108.37	113.71

There are no chirality outliers.

All (43) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	147	MLY	N-CA-CB-CG
1	A	147	MLY	C-CA-CB-CG
1	A	266	MLY	O-C-CA-CB
2	B	65	MLY	N-CA-CB-CG
2	B	65	MLY	C-CA-CB-CG
3	C	21	MLY	O-C-CA-CB
1	A	72	MLY	CD-CE-NZ-CH1
1	A	72	MLY	CD-CE-NZ-CH2
1	A	93	MLY	CD-CE-NZ-CH1
1	A	93	MLY	CD-CE-NZ-CH2
1	A	147	MLY	CD-CE-NZ-CH2
1	A	221	MLY	CD-CE-NZ-CH1
1	A	221	MLY	CD-CE-NZ-CH2
2	B	74	MLY	CG-CD-CE-NZ
1	A	70	MLY	CD-CE-NZ-CH1
3	C	6	MLY	CD-CE-NZ-CH2
3	C	15	MLY	CD-CE-NZ-CH1
1	A	72	MLY	CG-CD-CE-NZ
3	C	15	MLY	CG-CD-CE-NZ
1	A	147	MLY	CD-CE-NZ-CH1
3	C	6	MLY	CD-CE-NZ-CH1
3	C	21	MLY	CG-CD-CE-NZ
3	C	15	MLY	CD-CE-NZ-CH2
1	A	137	MLY	CG-CD-CE-NZ
1	A	264	MLY	CA-CB-CG-CD
2	B	65	MLY	CG-CD-CE-NZ
1	A	100	MLY	CE-CD-CG-CB
1	A	72	MLY	CE-CD-CG-CB
3	C	31	MLY	CE-CD-CG-CB
1	A	122	MLY	CG-CD-CE-NZ
1	A	147	MLY	CA-CB-CG-CD
3	C	31	MLY	CD-CE-NZ-CH1
3	C	21	MLY	CE-CD-CG-CB
1	A	93	MLY	CA-CB-CG-CD
3	C	15	MLY	CE-CD-CG-CB
1	A	254	MLY	CG-CD-CE-NZ
1	A	70	MLY	CE-CD-CG-CB
1	A	100	MLY	CG-CD-CE-NZ
1	A	266	MLY	N-CA-CB-CG
1	A	122	MLY	CD-CE-NZ-CH2
2	B	74	MLY	CD-CE-NZ-CH1
1	A	122	MLY	C-CA-CB-CG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	C	113	MLY	C-CA-CB-CG

There are no ring outliers.

5 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	122	MLY	1	0
3	C	15	MLY	1	0
1	A	100	MLY	5	0
1	A	264	MLY	1	0
1	A	72	MLY	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	217/246 (88%)	1.35	42 (19%) 3 2	23, 31, 66, 70	0
2	B	119/124 (95%)	1.03	13 (10%) 10 10	28, 32, 42, 53	0
3	C	121/143 (84%)	1.37	26 (21%) 2 2	26, 34, 50, 51	0
All	All	457/513 (89%)	1.27	81 (17%) 4 3	23, 32, 52, 70	0

All (81) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	269	GLY	8.3
1	A	107	LEU	8.1
1	A	110	LEU	6.8
1	A	97	LEU	6.4
1	A	112	ASN	6.3
1	A	114	PRO	6.1
1	A	104	SER	6.0
2	B	56	TRP	5.5
3	C	127	PRO	5.2
1	A	109	LEU	5.0
3	C	126	THR	4.9
1	A	150	PRO	4.8
2	B	3	TRP	4.8
2	B	122	ALA	4.7
1	A	103	VAL	4.6
1	A	113	SER	4.5
1	A	124	TYR	4.3
1	A	106	LEU	4.3
3	C	50	ASP	4.3
1	A	44	GLY	4.2
1	A	102	ASN	4.0
3	C	4	LEU	3.9
1	A	89	GLN	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	98	GLU	3.7
3	C	3	ASN	3.7
3	C	60	VAL	3.7
1	A	96	GLU	3.6
3	C	58	ASN	3.4
1	A	46	LEU	3.4
1	A	111	SER	3.2
2	B	4	ILE	3.2
1	A	108	SER	3.2
1	A	268	PHE	3.2
1	A	152	LEU	3.1
1	A	153	ALA	3.0
3	C	2	GLN	2.9
3	C	59	ASN	2.9
3	C	54	LEU	2.9
1	A	95	PRO	2.9
1	A	116	ILE	2.9
1	A	94	VAL	2.8
2	B	98	GLU	2.8
1	A	248	SER	2.8
1	A	99	GLN	2.8
1	A	52	MET	2.8
1	A	118	LEU	2.8
1	A	251	GLY	2.7
1	A	149	ARG	2.6
3	C	82	THR	2.6
1	A	157	HIS	2.6
2	B	2	SER	2.6
3	C	57	GLY	2.6
1	A	66	LEU	2.6
2	B	25	PRO	2.5
3	C	53	MET	2.5
1	A	247	GLN	2.5
3	C	5	LEU	2.5
1	A	101	GLN	2.4
3	C	61	ASN	2.4
3	C	125	LEU	2.4
3	C	10	ALA	2.4
1	A	249	GLY	2.3
2	B	55	ALA	2.3
3	C	52	PRO	2.3
3	C	97	LEU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	91	LEU	2.2
2	B	7	ILE	2.2
3	C	66	ARG	2.2
2	B	32	ALA	2.1
2	B	10	HIS	2.1
1	A	151	GLU	2.1
1	A	126	GLU	2.1
2	B	73	GLN	2.1
3	C	18	VAL	2.1
3	C	64	LEU	2.1
3	C	99	LEU	2.1
3	C	9	ALA	2.0
1	A	56	VAL	2.0
3	C	49	ILE	2.0
3	C	87	ASP	2.0
2	B	21	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	MLY	A	100	11/12	0.48	0.30	64,65,70,70	0
1	MLY	A	70	11/12	0.74	0.16	33,34,48,48	0
1	MLY	A	147	11/12	0.75	0.20	45,47,54,54	0
1	MLY	A	72	11/12	0.79	0.16	30,32,44,46	0
1	MLY	A	93	11/12	0.80	0.17	48,52,60,61	0
3	MLY	C	6	11/12	0.80	0.19	33,36,49,49	0
3	MLY	C	21	11/12	0.80	0.15	34,36,46,46	0
2	MLY	B	65	11/12	0.82	0.15	29,30,40,42	0
1	MLY	A	122	11/12	0.82	0.14	34,35,49,49	0
1	MLY	A	266	11/12	0.82	0.16	39,44,48,51	0
3	MLY	C	31	11/12	0.82	0.17	30,31,47,48	0
1	MLY	A	128	11/12	0.84	0.12	33,33,37,39	0
1	MLY	A	137	11/12	0.86	0.14	28,31,45,45	0
1	MLY	A	254	11/12	0.86	0.13	31,36,52,54	0
2	MLY	B	74	11/12	0.86	0.14	32,34,45,45	0
3	MLY	C	15	11/12	0.87	0.11	38,39,48,49	0
1	MLY	A	264	11/12	0.89	0.15	32,37,44,45	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	MLY	A	221	11/12	0.89	0.15	27,29,48,51	0
1	MLY	A	237	11/12	0.94	0.08	21,23,30,32	0
3	MLY	C	113	11/12	0.94	0.10	28,30,41,43	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.