



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 02:35 PM UTC

PDB ID : 1XLZ / pdb_00001xlz
Title : Catalytic Domain Of Human Phosphodiesterase 4B In Complex With Filaminast
Authors : Card, G.L.; England, B.P.; Suzuki, Y.; Fong, D.; Powell, B.; Lee, B.; Luu, C.; Tabrizizad, M.; Gillette, S.; Ibrahim, P.N.; Artis, D.R.; Bollag, G.; Milburn, M.V.; Kim, S.-H.; Schlessinger, J.; Zhang, K.Y.J.
Deposited on : 2004-09-30
Resolution : 2.06 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtrriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.06 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3774 (2.08-2.04)
Clashscore	190562	3883 (2.08-2.04)
Ramachandran outliers	187476	3860 (2.08-2.04)
Sidechain outliers	187428	3860 (2.08-2.04)
RSRZ outliers	180081	3775 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	398	
1	B	398	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5424 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cAMP-specific 3',5'-cyclic phosphodiesterase 4B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	323	Total	C	N	O	S	8	0	0
			2613	1651	441	502	19			
1	B	323	Total	C	N	O	S	4	0	0
			2613	1651	441	502	19			

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	131	MET	-	initiating methionine	UNP Q07343
A	132	GLY	-	cloning artifact	UNP Q07343
A	133	SER	-	cloning artifact	UNP Q07343
A	134	SER	-	cloning artifact	UNP Q07343
A	135	HIS	-	expression tag	UNP Q07343
A	136	HIS	-	expression tag	UNP Q07343
A	137	HIS	-	expression tag	UNP Q07343
A	138	HIS	-	expression tag	UNP Q07343
A	139	HIS	-	expression tag	UNP Q07343
A	140	HIS	-	expression tag	UNP Q07343
A	141	SER	-	cloning artifact	UNP Q07343
A	142	SER	-	cloning artifact	UNP Q07343
A	143	GLY	-	cloning artifact	UNP Q07343
A	144	LEU	-	cloning artifact	UNP Q07343
A	145	VAL	-	cloning artifact	UNP Q07343
A	146	PRO	-	cloning artifact	UNP Q07343
A	147	ARG	-	cloning artifact	UNP Q07343
A	148	GLY	-	cloning artifact	UNP Q07343
A	149	SER	-	cloning artifact	UNP Q07343
A	150	HIS	-	cloning artifact	UNP Q07343
A	151	MET	-	cloning artifact	UNP Q07343
A	194	CME	CYS	modified residue	UNP Q07343
A	320	CME	CYS	modified residue	UNP Q07343
B	131	MET	-	initiating methionine	UNP Q07343
B	132	GLY	-	cloning artifact	UNP Q07343

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	133	SER	-	cloning artifact	UNP Q07343
B	134	SER	-	cloning artifact	UNP Q07343
B	135	HIS	-	expression tag	UNP Q07343
B	136	HIS	-	expression tag	UNP Q07343
B	137	HIS	-	expression tag	UNP Q07343
B	138	HIS	-	expression tag	UNP Q07343
B	139	HIS	-	expression tag	UNP Q07343
B	140	HIS	-	expression tag	UNP Q07343
B	141	SER	-	cloning artifact	UNP Q07343
B	142	SER	-	cloning artifact	UNP Q07343
B	143	GLY	-	cloning artifact	UNP Q07343
B	144	LEU	-	cloning artifact	UNP Q07343
B	145	VAL	-	cloning artifact	UNP Q07343
B	146	PRO	-	cloning artifact	UNP Q07343
B	147	ARG	-	cloning artifact	UNP Q07343
B	148	GLY	-	cloning artifact	UNP Q07343
B	149	SER	-	cloning artifact	UNP Q07343
B	150	HIS	-	cloning artifact	UNP Q07343
B	151	MET	-	cloning artifact	UNP Q07343
B	194	CME	CYS	modified residue	UNP Q07343
B	320	CME	CYS	modified residue	UNP Q07343


- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Zn 1 1	0	0
2	B	1	Total Zn 1 1	0	0

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mg 1 1	0	0
3	B	1	Total Mg 1 1	0	0

- Molecule 4 is (1E)-1-[3-(CYCLOPENTYLOXY)-4-METHOXYPHENYL]ETHANONE O-(AMINOCARBONYL)OXIME (CCD ID: FIL) (formula: C₁₅H₂₀N₂O₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			21	15	2	4		
4	B	1	Total	C	N	O	0	0
			21	15	2	4		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	83	Total	O	0	0
			83	83		
5	B	69	Total	O	0	0
			69	69		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: cAMP-specific 3',5'-cyclic phosphodiesterase 4B



- Molecule 1: cAMP-specific 3',5'-cyclic phosphodiesterase 4B



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	89.54Å 94.17Å 106.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	70.71 – 2.06 70.49 – 2.06	Depositor EDS
% Data completeness (in resolution range)	99.2 (70.71-2.06) 99.2 (70.49-2.06)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.92 (at 2.07Å)	Xtriage
Refinement program	REFMAC 5.1.25	Depositor
R, R_{free}	0.209 , 0.237 0.215 , 0.241	Depositor DCC
R_{free} test set	2830 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	39.4	Xtriage
Anisotropy	0.348	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 36.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.012 for k,h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5424	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 42.59 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.9948e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN, CME, FIL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.85	6/2654 (0.2%)	0.99	4/3597 (0.1%)
1	B	0.83	3/2654 (0.1%)	0.97	4/3597 (0.1%)
All	All	0.84	9/5308 (0.2%)	0.98	8/7194 (0.1%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	411	MET	SD-CE	-10.96	1.52	1.79
1	A	426	MET	SD-CE	-9.52	1.55	1.79
1	A	387	MET	SD-CE	-8.37	1.58	1.79
1	B	411	MET	SD-CE	-7.52	1.60	1.79
1	B	221	MET	SD-CE	-6.93	1.62	1.79
1	B	387	MET	SD-CE	-6.80	1.62	1.79
1	A	318	GLU	CB-CG	-6.77	1.32	1.52
1	A	347	MET	SD-CE	-6.11	1.64	1.79
1	A	328	LYS	CB-CG	5.37	1.68	1.52

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	326	LEU	N-CA-C	-6.09	102.92	110.41
1	B	164	ASP	N-CA-C	-6.03	104.78	111.36
1	A	164	ASP	N-CA-C	-5.99	104.88	111.82
1	B	232	ALA	N-CA-C	5.40	117.17	111.28
1	A	232	ALA	N-CA-C	5.39	117.16	111.28
1	B	316	GLN	N-CA-C	5.36	118.04	111.82
1	A	385	ARG	CB-CA-C	-5.29	102.34	110.81
1	A	183	VAL	N-CA-CB	5.01	118.10	110.58

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2613	0	2541	25	0
1	B	2613	0	2541	22	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	21	0	20	3	0
4	B	21	0	20	4	0
5	A	83	0	0	1	0
5	B	69	0	0	1	0
All	All	5424	0	5122	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:217:ILE:HG22	1:B:221:MET:HE2	1.58	0.85
1:B:340:ASP:OD1	1:B:385:ARG:HD3	1.81	0.81
1:A:424:ARG:HD2	1:A:426:MET:CE	2.20	0.71
1:A:163:GLU:CD	1:A:163:GLU:N	2.50	0.69
1:B:410:ILE:HG23	1:B:411:MET:HE2	1.77	0.66
1:A:340:ASP:OD1	1:A:385:ARG:HD3	1.99	0.62
1:A:424:ARG:HD2	1:A:426:MET:HE2	1.83	0.61
1:B:289:THR:HG22	1:B:426:MET:HE1	1.84	0.60
1:A:258:VAL:HG11	1:A:374:LEU:HD12	1.84	0.60
1:B:258:VAL:HG11	1:B:374:LEU:HD12	1.86	0.58
1:A:246:THR:HG23	1:A:387:MET:HE3	1.86	0.57
1:B:376:ASN:CG	1:B:377:TYR:H	2.13	0.56
1:A:252:THR:HG22	1:A:465:ASP:OD2	2.06	0.56
1:A:163:GLU:N	1:A:163:GLU:OE1	2.38	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:410:ILE:HG23	1:B:411:MET:CE	2.36	0.55
1:B:411:MET:HE1	4:B:1003:FIL:C15	2.38	0.54
1:B:274:HIS:O	1:B:307:HIS:CD2	2.61	0.53
1:B:411:MET:HE1	4:B:1003:FIL:H151	1.91	0.53
1:A:327:THR:HG22	1:A:330:GLN:H	1.73	0.52
1:A:250:LEU:CD2	1:A:387:MET:HE1	2.39	0.52
1:A:309:ALA:O	1:A:313:LYS:HB2	2.11	0.51
1:B:397:THR:HB	1:B:469:ILE:HG23	1.91	0.51
1:A:356:ASP:HB3	1:A:382:GLN:NE2	2.26	0.51
1:B:424:ARG:CB	1:B:426:MET:HE3	2.43	0.48
4:B:1003:FIL:H4	5:B:2028:HOH:O	2.13	0.48
1:A:411:MET:CE	4:A:1003:FIL:H152	2.45	0.46
1:A:397:THR:HB	1:A:469:ILE:HG23	1.97	0.46
1:B:356:ASP:HB3	1:B:382:GLN:NE2	2.31	0.46
1:B:255:LEU:HD23	1:B:372:LEU:HD12	1.98	0.45
1:A:411:MET:HE1	4:A:1003:FIL:C15	2.47	0.45
1:A:258:VAL:CG1	1:A:374:LEU:HD12	2.45	0.45
1:A:180:ILE:HA	1:A:183:VAL:HG13	1.99	0.44
1:A:270:ALA:O	1:A:274:HIS:HB3	2.17	0.44
1:B:300:GLU:O	1:B:301:SER:C	2.61	0.44
1:B:255:LEU:CD2	1:B:372:LEU:HD12	2.47	0.44
1:B:442:SER:HB3	4:B:1003:FIL:H142	2.00	0.44
1:B:459:ALA:HB2	1:B:470:LEU:HD13	2.00	0.43
1:B:180:ILE:HA	1:B:183:VAL:HG13	2.00	0.43
1:A:275:ASP:O	1:A:278:HIS:HB2	2.19	0.43
4:A:1003:FIL:H16	4:A:1003:FIL:H11	1.87	0.42
1:A:250:LEU:HG	1:A:387:MET:HE1	2.02	0.41
1:B:275:ASP:HA	1:B:307:HIS:CD2	2.55	0.41
1:A:443:GLN:HG2	5:A:1019:HOH:O	2.19	0.41
1:A:250:LEU:HD21	1:A:387:MET:HE1	2.01	0.41
1:B:327:THR:HG22	1:B:329:LYS:N	2.35	0.41
1:A:275:ASP:HA	1:A:307:HIS:CD2	2.56	0.41
1:B:424:ARG:HB3	1:B:426:MET:HE3	2.02	0.41
1:A:190:ARG:CD	1:A:221:MET:HE3	2.51	0.41
1:A:390:CYS:HB3	1:A:455:TRP:CZ2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	319/398 (80%)	313 (98%)	6 (2%)	0	100	100
1	B	319/398 (80%)	312 (98%)	7 (2%)	0	100	100
All	All	638/796 (80%)	625 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	291/358 (81%)	278 (96%)	13 (4%)	24	18
1	B	291/358 (81%)	279 (96%)	12 (4%)	27	21
All	All	582/716 (81%)	557 (96%)	25 (4%)	26	20

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	163	GLU
1	A	168	LYS
1	A	178	LEU
1	A	183	VAL
1	A	204	ASP
1	A	212	SER
1	A	313	LYS
1	A	326	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	327	THR
1	A	352	SER
1	A	373	LEU
1	A	416	GLN
1	A	463	GLN
1	B	163	GLU
1	B	168	LYS
1	B	183	VAL
1	B	189	ASN
1	B	317	GLU
1	B	327	THR
1	B	352	SER
1	B	360	MET
1	B	364	LYS
1	B	365	LYS
1	B	366	VAL
1	B	470	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	284	GLN
1	A	382	GLN
1	A	386	ASN
1	A	443	GLN
1	A	463	GLN
1	B	284	GLN
1	B	376	ASN
1	B	382	GLN
1	B	443	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CME	A	194	1	4,5,10	1.46	1 (25%)	1,5,11	2.16	1 (100%)
1	CME	A	320	1	4,5,10	1.23	1 (25%)	1,5,11	1.65	0
1	CME	B	194	1	4,5,10	1.51	1 (25%)	1,5,11	0.04	0
1	CME	B	320	1	4,5,10	1.59	1 (25%)	1,5,11	0.75	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CME	A	194	1	-	1/1/4/10	-
1	CME	A	320	1	-	1/1/4/10	-
1	CME	B	194	1	-	1/1/4/10	-
1	CME	B	320	1	-	0/1/4/10	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	194	CME	CB-SG	-2.56	1.76	1.81
1	B	194	CME	CB-SG	-2.50	1.76	1.81
1	B	320	CME	CB-CA	2.34	1.55	1.53
1	A	320	CME	CB-SG	-2.25	1.76	1.81

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	194	CME	CA-CB-SG	-2.16	109.79	114.40

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	320	CME	N-CA-CB-SG
1	A	194	CME	N-CA-CB-SG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
1	B	194	CME	N-CA-CB-SG

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	FIL	B	1003	-	22,22,22	3.90	2 (9%)	26,29,29	2.39	5 (19%)
4	FIL	A	1003	-	22,22,22	3.73	2 (9%)	26,29,29	2.25	6 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FIL	B	1003	-	-	5/13/22/22	0/2/2/2
4	FIL	A	1003	-	-	8/13/22/22	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	1003	FIL	C1-N2	16.45	1.47	1.28
4	A	1003	FIL	C1-N2	15.84	1.46	1.28

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	1003	FIL	O21-C18	7.41	1.47	1.38
4	A	1003	FIL	O21-C18	7.14	1.46	1.38

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	1003	FIL	C17-C1-N2	-8.91	106.66	123.95
4	A	1003	FIL	C17-C1-N2	-6.09	112.14	123.95
4	A	1003	FIL	C17-C1-C3	-5.73	110.21	119.48
4	B	1003	FIL	C17-C1-C3	-5.53	110.53	119.48
4	A	1003	FIL	C8-O7-C6	5.15	125.07	117.51
4	B	1003	FIL	C8-O7-C6	3.81	123.10	117.51
4	A	1003	FIL	O7-C6-C9	3.79	120.54	115.40
4	A	1003	FIL	C3-C1-N2	-2.47	109.79	115.30
4	A	1003	FIL	O7-C6-C5	-2.36	120.33	124.30
4	B	1003	FIL	O19-C18-N20	-2.27	122.01	125.58
4	B	1003	FIL	O7-C6-C9	2.24	118.44	115.40

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1003	FIL	O19-C18-O21-N2
4	A	1003	FIL	C3-C1-N2-O21
4	A	1003	FIL	N2-C1-C3-C4
4	A	1003	FIL	N2-C1-C3-C16
4	A	1003	FIL	C12-C11-O10-C9
4	A	1003	FIL	C15-C11-O10-C9
4	B	1003	FIL	O19-C18-O21-N2
4	B	1003	FIL	C17-C1-N2-O21
4	B	1003	FIL	C3-C1-N2-O21
4	B	1003	FIL	C17-C1-C3-C4
4	B	1003	FIL	C17-C1-C3-C16
4	A	1003	FIL	C17-C1-C3-C16
4	A	1003	FIL	C17-C1-C3-C4

There are no ring outliers.

2 monomers are involved in 7 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	1003	FIL	4	0


Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1003	FIL	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	321/398 (80%)	0.30	8 (2%) 58 60	17, 27, 39, 49	2 (0%)
1	B	321/398 (80%)	0.18	5 (1%) 70 73	20, 27, 38, 42	1 (0%)
All	All	642/796 (80%)	0.24	13 (2%) 65 67	17, 27, 39, 49	3 (0%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	316	GLN	3.2
1	B	375	ASP	3.0
1	A	318	GLU	2.8
1	A	204	ASP	2.6
1	A	322	ILE	2.5
1	B	163	GLU	2.4
1	A	360	MET	2.3
1	B	214	ASP	2.3
1	A	423	GLU	2.2
1	A	197	TYR	2.2
1	A	331	ARG	2.2
1	A	296	MET	2.2
1	B	327	THR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CME	A	320	6/11	0.83	0.12	43,46,46,54	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	CME	B	320	6/11	0.89	0.12	37,39,39,44	0
1	CME	B	194	6/11	0.95	0.08	28,29,31,38	0
1	CME	A	194	6/11	0.96	0.07	29,30,32,39	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	FIL	A	1003	21/21	0.86	0.19	51,55,74,78	0
4	FIL	B	1003	21/21	0.88	0.18	56,58,74,76	0
3	MG	A	1002	1/1	0.99	0.03	21,21,21,21	0
3	MG	B	1002	1/1	0.99	0.05	23,23,23,23	0
2	ZN	A	1001	1/1	1.00	0.01	33,33,33,33	0
2	ZN	B	1001	1/1	1.00	0.03	34,34,34,34	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FIL A 1003:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.