



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 07:21 AM UTC

PDB ID : 4XL3 / pdb_00004xl3
Title : Crystal structure of reduced form of thiolase from Clostridium acetobutylicum
Authors : Kim, S.; Ha, S.C.; Ahn, J.W.; Kim, E.J.; Lim, J.H.; Kim, K.J.
Deposited on : 2015-01-13
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	400	2% 87% 10% ..
1	B	400	8% 84% 14% ..

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6761 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acetyl-CoA acetyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	393	Total	C	N	O	S	0	7	0
			2974	1873	521	566	14			
1	B	393	Total	C	N	O	S	0	7	0
			2965	1868	516	567	14			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	378	SER	CYS	engineered mutation	UNP F0K5D8
A	393	LEU	-	expression tag	UNP F0K5D8
A	394	GLU	-	expression tag	UNP F0K5D8
A	395	HIS	-	expression tag	UNP F0K5D8
A	396	HIS	-	expression tag	UNP F0K5D8
A	397	HIS	-	expression tag	UNP F0K5D8
A	398	HIS	-	expression tag	UNP F0K5D8
A	399	HIS	-	expression tag	UNP F0K5D8
A	400	HIS	-	expression tag	UNP F0K5D8
B	378	SER	CYS	engineered mutation	UNP F0K5D8
B	393	LEU	-	expression tag	UNP F0K5D8
B	394	GLU	-	expression tag	UNP F0K5D8
B	395	HIS	-	expression tag	UNP F0K5D8
B	396	HIS	-	expression tag	UNP F0K5D8
B	397	HIS	-	expression tag	UNP F0K5D8
B	398	HIS	-	expression tag	UNP F0K5D8
B	399	HIS	-	expression tag	UNP F0K5D8
B	400	HIS	-	expression tag	UNP F0K5D8

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	427	Total	O	0	0
			427	427		
3	B	359	Total	O	0	0
			359	359		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Acetyl-CoA acetyltransferase



- Molecule 1: Acetyl-CoA acetyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	204.42Å 54.29Å 73.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.70 50.00 – 1.70	Depositor EDS
% Data completeness (in resolution range)	97.7 (50.00-1.70) 97.7 (50.00-1.70)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	11.38 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.153 , 0.192 (Not available) , 0.211	Depositor DCC
R_{free} test set	4449 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	17.2	Xtriage
Anisotropy	0.330	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 47.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6761	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.41	8/3021 (0.3%)	1.14	4/4072 (0.1%)
1	B	1.36	8/3018 (0.3%)	1.21	11/4070 (0.3%)
All	All	1.39	16/6039 (0.3%)	1.18	15/8142 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	177[A]	GLU	CA-C	8.26	1.63	1.52
1	B	177[B]	GLU	CA-C	8.26	1.63	1.52
1	B	114	GLY	N-CA	6.37	1.51	1.45
1	A	165	ALA	N-CA	6.20	1.53	1.46
1	A	365	MET	N-CA	6.07	1.53	1.46
1	B	254	ALA	C-O	5.80	1.30	1.23
1	B	241	VAL	CA-C	-5.68	1.45	1.52
1	A	182	SER	CA-C	-5.67	1.45	1.52
1	A	92	LEU	C-O	5.34	1.30	1.24
1	A	276	VAL	C-O	5.34	1.29	1.24
1	A	19	LYS	C-O	-5.33	1.19	1.23
1	B	373	GLY	N-CA	5.28	1.50	1.44
1	B	85	ASN	C-O	-5.14	1.18	1.24
1	A	295	ALA	C-O	5.13	1.30	1.24
1	B	70	SER	N-CA	5.10	1.52	1.46
1	A	367	LYS	N-CA	5.01	1.52	1.46

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	177[A]	GLU	CA-C-O	12.28	133.56	120.55
1	B	177[B]	GLU	CA-C-O	12.28	133.56	120.55
1	B	7	ALA	N-CA-C	7.87	119.94	111.36
1	B	177[A]	GLU	N-CA-C	7.47	119.42	111.28
1	B	177[B]	GLU	N-CA-C	7.47	119.42	111.28
1	B	350	ILE	CB-CA-C	-6.73	103.27	111.88
1	A	19	LYS	N-CA-C	6.38	117.72	108.34
1	B	165	ALA	N-CA-C	-6.27	104.45	111.28
1	A	7	ALA	N-CA-C	6.24	118.16	111.36
1	B	371	LYS	N-CA-C	6.18	119.29	111.69
1	A	350	ILE	CB-CA-C	-6.16	104.00	111.88
1	A	299	ALA	N-CA-C	-5.31	105.58	111.36
1	B	377	LEU	N-CA-C	5.12	116.84	108.76
1	B	218	HIS	CA-C-N	-5.07	114.55	119.77
1	B	218	HIS	C-N-CA	-5.07	114.55	119.77

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	349	PRO	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2974	0	3052	28	0
1	B	2965	0	3040	40	0
2	A	24	0	32	1	0
2	B	12	0	16	4	0
3	A	427	0	0	3	0
3	B	359	0	0	6	0
All	All	6761	0	6140	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1:MET:HB2	3:B:704:HOH:O	1.78	0.82
1:A:84:ILE:HD12	1:B:84:ILE:HD12	1.60	0.81
1:B:133[A]:ARG:HG3	1:B:133[A]:ARG:HH11	1.48	0.78
1:A:366:GLN:HG2	3:A:890:HOH:O	1.93	0.67
1:A:362:VAL:HG12	1:A:389:LEU:CD1	2.25	0.66
1:B:385:THR:CG2	3:B:693:HOH:O	2.44	0.65
1:B:40:LYS:HZ1	2:B:502:GOL:C3	2.10	0.65
1:A:167[B]:ARG:CD	1:A:167[B]:ARG:C	2.70	0.63
1:B:385:THR:HG23	3:B:693:HOH:O	1.99	0.63
1:B:133[A]:ARG:HG3	1:B:133[A]:ARG:NH1	2.11	0.61
1:A:157:MET:HA	1:A:157:MET:HE2	1.84	0.60
1:B:348:HIS:CD2	2:B:501:GOL:H12	2.37	0.59
1:A:167[B]:ARG:C	1:A:167[B]:ARG:HD2	2.28	0.58
1:A:358:LEU:O	1:A:362:VAL:HG13	2.05	0.57
1:B:297:LYS:HA	1:B:297:LYS:CE	2.35	0.57
1:A:167[B]:ARG:O	1:A:167[B]:ARG:HD3	2.05	0.56
1:A:383:GLN:HE22	1:B:81:ALA:H	1.54	0.54
1:A:81:ALA:H	1:B:383:GLN:HE22	1.56	0.54
1:B:297:LYS:HA	1:B:297:LYS:HE2	1.90	0.54
1:B:22:LYS:HE2	3:B:661:HOH:O	2.08	0.54
1:B:252:CYS:HB3	1:B:349:PRO:HG3	1.91	0.53
1:A:250:ASN:HD22	1:A:348:HIS:H	1.57	0.52
1:B:40:LYS:HZ1	2:B:502:GOL:H31	1.75	0.52
1:B:348:HIS:HD2	2:B:501:GOL:H12	1.76	0.50
1:A:85:ASN:HD22	1:B:58:GLN:NE2	2.11	0.48
1:A:250:ASN:ND2	1:A:348:HIS:H	2.13	0.47
1:B:55:ASN:HD22	1:B:58:GLN:HG2	1.79	0.47
1:B:250:ASN:HD22	1:B:348:HIS:H	1.61	0.46
1:A:369:ASP:HA	1:A:391:LYS:HE2	1.98	0.46
1:A:167[B]:ARG:CD	1:A:167[B]:ARG:O	2.65	0.45
1:A:383:GLN:NE2	1:B:81:ALA:H	2.15	0.45
1:A:277:SER:OG	1:A:388:LEU:HD12	2.17	0.45
1:B:250:ASN:ND2	1:B:348:HIS:H	2.16	0.44
1:B:72:LYS:HE3	3:B:620:HOH:O	2.17	0.44
1:A:218:HIS:N	1:A:219:PRO:CD	2.81	0.44
1:B:88:CYS:HB2	1:B:379:ILE:HA	2.00	0.44
1:B:55:ASN:HD21	1:B:61:LEU:HD12	1.82	0.43
1:B:311:ASP:HB2	1:B:370:ALA:HB1	2.00	0.43
1:B:367:LYS:HE2	3:B:794:HOH:O	2.19	0.43
1:A:55:ASN:HD22	1:A:58:GLN:HG2	1.84	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:501:GOL:H11	3:A:889:HOH:O	2.18	0.42
1:B:291:GLY:N	1:B:292:PRO:CD	2.82	0.42
1:A:81:ALA:H	1:B:383:GLN:NE2	2.17	0.42
1:B:157:MET:HE2	1:B:157:MET:HA	2.02	0.41
1:A:124:LEU:CD2	1:B:124:LEU:CD2	2.99	0.41
1:B:55:ASN:ND2	1:B:58:GLN:HG2	2.35	0.41
1:A:55:ASN:HD21	1:A:61:LEU:HD12	1.84	0.41
1:B:306:THR:O	1:B:309:GLU:HB2	2.21	0.41
1:A:317:GLU:CD	1:A:342:GLY:HA3	2.46	0.41
1:B:39:LYS:HB3	1:B:39:LYS:HE2	1.77	0.41
1:B:305:TRP:CZ2	1:B:372:LYS:HG2	2.56	0.41
1:A:157:MET:HE2	1:A:157:MET:CA	2.51	0.41
1:B:111:ILE:HD13	1:B:257:VAL:HG22	2.04	0.40
1:B:263:LYS:HA	1:B:263:LYS:HD3	1.93	0.40
1:A:124:LEU:CD2	1:B:124:LEU:HD22	2.51	0.40
1:A:141:ASP:OD1	1:A:141:ASP:C	2.64	0.40
1:B:111:ILE:CD1	1:B:257:VAL:HG22	2.52	0.40
1:B:301:GLU:OE1	1:B:301:GLU:HA	2.21	0.40
1:A:22:LYS:HE2	3:A:636:HOH:O	2.20	0.40
1:A:55:ASN:ND2	1:A:58:GLN:HG2	2.36	0.40
1:B:44:LYS:O	1:B:47[A]:ASP:HB2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	399/400 (100%)	385 (96%)	13 (3%)	1 (0%)	36	22
1	B	399/400 (100%)	387 (97%)	11 (3%)	1 (0%)	36	22
All	All	798/800 (100%)	772 (97%)	24 (3%)	2 (0%)	36	22

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	87	VAL
1	B	87	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	303/302 (100%)	294 (97%)	9 (3%)	36	19
1	B	303/302 (100%)	290 (96%)	13 (4%)	26	10
All	All	606/604 (100%)	584 (96%)	22 (4%)	34	14

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	87	VAL
1	A	188	GLU
1	A	207	ARG
1	A	270	LYS
1	A	362	VAL
1	A	366	GLN
1	A	367	LYS
1	A	371	LYS
1	B	1	MET
1	B	87	VAL
1	B	133[A]	ARG
1	B	133[B]	ARG
1	B	133[C]	ARG
1	B	210[A]	GLU
1	B	210[B]	GLU
1	B	230	LYS
1	B	270	LYS
1	B	300	ILE
1	B	322	GLN
1	B	329	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	393	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	55	ASN
1	A	58	GLN
1	A	153	ASN
1	A	194	GLN
1	A	250	ASN
1	A	366	GLN
1	A	383	GLN
1	B	55	ASN
1	B	58	GLN
1	B	153	ASN
1	B	250	ASN
1	B	316	ASN
1	B	348	HIS
1	B	383	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	A	502	-	5,5,5	0.45	0	5,5,5	0.72	0
2	GOL	B	502	-	5,5,5	0.41	0	5,5,5	0.67	0
2	GOL	A	503	-	5,5,5	0.45	0	5,5,5	0.53	0
2	GOL	B	501	-	5,5,5	0.36	0	5,5,5	1.17	0
2	GOL	A	504	-	5,5,5	0.61	0	5,5,5	1.14	0
2	GOL	A	501	-	5,5,5	0.51	0	5,5,5	1.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	502	-	-	0/4/4/4	-
2	GOL	B	502	-	-	0/4/4/4	-
2	GOL	A	503	-	-	0/4/4/4	-
2	GOL	B	501	-	-	2/4/4/4	-
2	GOL	A	504	-	-	2/4/4/4	-
2	GOL	A	501	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	504	GOL	C1-C2-C3-O3
2	B	501	GOL	O1-C1-C2-O2
2	A	501	GOL	C1-C2-C3-O3
2	B	501	GOL	O1-C1-C2-C3
2	A	504	GOL	O2-C2-C3-O3
2	A	501	GOL	O1-C1-C2-C3

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	502	GOL	2	0
2	B	501	GOL	2	0
2	A	501	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	393/400 (98%)	-0.23	6 (1%) 72 76	7, 16, 32, 60	7 (1%)
1	B	393/400 (98%)	0.17	33 (8%) 17 18	6, 19, 42, 97	7 (1%)
All	All	786/800 (98%)	-0.03	39 (4%) 34 38	6, 17, 37, 97	14 (1%)

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	290	TYR	6.0
1	B	393	LEU	5.6
1	A	133[A]	ARG	5.1
1	B	206	GLY	4.9
1	B	298	ALA	4.6
1	A	393	LEU	4.4
1	B	303	ALA	4.0
1	B	304	GLY	4.0
1	B	133[A]	ARG	4.0
1	B	300	ILE	3.9
1	B	209	GLY	3.7
1	A	206	GLY	3.6
1	B	168	TRP	3.4
1	B	379	ILE	3.3
1	B	378	SER	3.2
1	B	294	TYR	3.2
1	B	291	GLY	3.2
1	B	392	CYS	2.9
1	B	305	TRP	2.9
1	B	282	GLY	2.9
1	B	292	PRO	2.7
1	B	287	ILE	2.7
1	B	382	GLY	2.7
1	B	307	VAL	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	285	PRO	2.6
1	B	284	ASP	2.6
1	B	302	LYS	2.6
1	B	293	PHE	2.5
1	A	392	CYS	2.5
1	B	208	LYS	2.4
1	B	299	ALA	2.4
1	B	286	ALA	2.4
1	A	207	ARG	2.1
1	B	381	GLY	2.1
1	B	308	ASP	2.1
1	B	283	VAL	2.1
1	A	308	ASP	2.1
1	B	210[A]	GLU	2.0
1	B	281	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GOL	A	504	6/6	0.77	0.17	27,40,45,53	0
2	GOL	B	501	6/6	0.80	0.16	26,34,36,41	0
2	GOL	B	502	6/6	0.81	0.15	29,41,44,45	0
2	GOL	A	501	6/6	0.82	0.14	17,25,28,28	0
2	GOL	A	503	6/6	0.88	0.12	19,32,35,38	0
2	GOL	A	502	6/6	0.91	0.11	16,25,32,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.