



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 10:21 PM UTC

PDB ID : 4XLN / pdb_00004xln
Title : Crystal structure of T. aquaticus transcription initiation complex containing bubble promoter and RNA
Authors : Bae, B.; Darst, S.A.
Deposited on : 2015-01-13
Resolution : 4.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 4.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1082 (4.20-3.80)
Clashscore	190562	1129 (4.20-3.80)
Ramachandran outliers	187476	1064 (4.20-3.80)
Sidechain outliers	187428	1055 (4.20-3.80)
RSRZ outliers	180081	1082 (4.20-3.80)
RNA backbone	3983	1017 (4.84-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	314	
1	B	314	
1	G	314	
1	H	314	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	C	1119	
2	I	1119	
3	D	1524	
3	J	1524	
4	E	99	
4	K	99	
5	F	347	
5	L	347	
6	O	48	
6	R	48	
7	P	48	
7	S	48	
8	Q	4	
8	T	4	

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 58255 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	227	Total	C	N	O	S	0	0	0
			1770	1130	303	334	3			
1	B	227	Total	C	N	O	S	0	0	0
			1770	1130	303	334	3			
1	G	227	Total	C	N	O	S	0	0	0
			1770	1130	303	334	3			
1	H	227	Total	C	N	O	S	0	0	0
			1770	1130	303	334	3			

- Molecule 2 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	1112	Total	C	N	O	S	0	0	0
			8739	5531	1553	1632	23			
2	I	1112	Total	C	N	O	S	0	0	0
			8739	5531	1553	1632	23			

- Molecule 3 is a protein called DNA-directed RNA polymerase subunit beta'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	1490	Total	C	N	O	S	0	0	0
			11761	7439	2088	2196	38			
3	J	1367	Total	C	N	O	S	0	0	0
			10779	6810	1923	2010	36			

- Molecule 4 is a protein called DNA-directed RNA polymerase subunit omega.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	E	93	Total	C	N	O	S	0	0	0
			768	490	136	138	4			
4	K	93	Total	C	N	O	S	0	0	0
			768	490	136	138	4			

- Molecule 5 is a protein called RNA polymerase sigma factor SigA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	F	347	Total	C	N	O	S	0	0	0
			2801	1767	505	525	4			
5	L	347	Total	C	N	O	S	0	0	0
			2801	1767	505	525	4			

- Molecule 6 is a DNA chain called DNA (48-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	O	48	Total	C	N	O	P	0	0	0
			988	472	182	287	47			
6	R	48	Total	C	N	O	P	0	0	0
			988	472	182	287	47			

- Molecule 7 is a DNA chain called DNA (48-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	P	48	Total	C	N	O	P	0	0	0
			985	471	183	284	47			
7	S	43	Total	C	N	O	P	0	0	0
			882	421	164	255	42			

- Molecule 8 is a RNA chain called RNA (5'-D(P*UP*CP*GP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	Q	4	Total	C	N	O	P	0	0	0
			85	38	15	28	4			
8	T	4	Total	C	N	O	P	0	0	0
			85	38	15	28	4			

- Molecule 9 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	D	2	Total	Zn	0	0
			2	2		
9	J	2	Total	Zn	0	0
			2	2		


- Molecule 10 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	D	1	Total 1	Mg 1	0	0
10	J	1	Total 1	Mg 1	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DNA-directed RNA polymerase subunit alpha





• Molecule 1: DNA-directed RNA polymerase subunit alpha




• Molecule 1: DNA-directed RNA polymerase subunit alpha










• Molecule 2: DNA-directed RNA polymerase subunit beta






• Molecule 3: DNA-directed RNA polymerase subunit beta'






• Molecule 3: DNA-directed RNA polymerase subunit beta'







- Molecule 4: DNA-directed RNA polymerase subunit omega



- Molecule 5: RNA polymerase sigma factor SigA



- Molecule 5: RNA polymerase sigma factor SigA



- Molecule 6: DNA (48-MER)







- Molecule 6: DNA (48-MER)




- Molecule 7: DNA (48-MER)



- Molecule 7: DNA (48-MER)



- Molecule 8: RNA (5'-D(P*UP*CP*GP*A)-3')



- Molecule 8: RNA (5'-D(P*UP*CP*GP*A)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	289.26Å 289.26Å 536.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.96 – 4.00 34.96 – 4.00	Depositor EDS
% Data completeness (in resolution range)	98.9 (34.96-4.00) 98.7 (34.96-4.00)	Depositor EDS
R_{merge}	0.26	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.07 (at 3.99Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1839)	Depositor
R, R_{free}	0.280 , 0.315 0.282 , 0.316	Depositor DCC
R_{free} test set	9497 reflections (4.25%)	wwPDB-VP
Wilson B-factor (Å ²)	99.6	Xtriage
Anisotropy	0.022	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 128.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.38$, $\langle L^2 \rangle = 0.21$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.84	EDS
Total number of atoms	58255	wwPDB-VP
Average B, all atoms (Å ²)	127.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.34	0/1804	0.86	3/2455 (0.1%)
1	B	0.34	0/1804	0.86	3/2455 (0.1%)
1	G	0.34	0/1804	0.85	2/2455 (0.1%)
1	H	0.34	0/1804	0.85	3/2455 (0.1%)
2	C	0.33	0/8905	0.84	24/12040 (0.2%)
2	I	0.33	1/8905 (0.0%)	0.84	24/12040 (0.2%)
3	D	0.31	0/11963	0.81	12/16165 (0.1%)
3	J	0.32	0/10959	0.81	10/14802 (0.1%)
4	E	0.26	0/783	0.67	0/1054
4	K	0.26	0/783	0.67	0/1054
5	F	0.29	0/2843	0.81	4/3822 (0.1%)
5	L	0.30	0/2843	0.80	4/3822 (0.1%)
6	O	0.11	0/1109	0.54	0/1712
6	R	0.11	0/1109	0.54	0/1712
7	P	0.15	0/1106	0.63	0/1706
7	S	0.11	0/989	0.51	0/1523
8	Q	0.07	0/94	0.15	0/144
8	T	0.07	0/94	0.15	0/144
All	All	0.31	1/59701 (0.0%)	0.80	89/81560 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2
2	C	0	2
2	I	0	1
All	All	0	5

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	I	764	GLU	CD-OE2	5.03	1.34	1.25

The worst 5 of 89 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	I	764	GLU	CA-C-N	11.37	142.16	121.70
2	I	764	GLU	C-N-CA	11.37	142.16	121.70
1	G	227	ASN	CA-C-N	8.78	128.92	120.31
1	G	227	ASN	C-N-CA	8.78	128.92	120.31
1	A	227	ASN	CA-C-N	8.75	128.88	120.31

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	177	VAL	Peptide,Mainchain
2	C	178	ALA	Peptide
2	C	764	GLU	Peptide
2	I	178	ALA	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1770	0	1799	57	0
1	B	1770	0	1799	56	0
1	G	1770	0	1799	59	0
1	H	1770	0	1799	56	0
2	C	8739	0	8841	300	0
2	I	8739	0	8841	285	0
3	D	11761	0	11976	354	0
3	J	10779	0	10993	325	0
4	E	768	0	784	19	0
4	K	768	0	784	21	0
5	F	2801	0	2881	98	0
5	L	2801	0	2881	88	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	O	988	0	544	17	0
6	R	988	0	544	16	0
7	P	985	0	543	17	0
7	S	882	0	487	14	0
8	Q	85	0	43	0	0
8	T	85	0	43	1	0
9	D	2	0	0	0	0
9	J	2	0	0	0	0
10	D	1	0	0	0	0
10	J	1	0	0	0	0
All	All	58255	0	57381	1598	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 1598 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:14:THR:HB	1:G:22:GLU:HB2	1.54	0.90
1:A:14:THR:HB	1:A:22:GLU:HB2	1.54	0.89
2:C:630:ARG:HG3	2:C:705:ILE:HB	1.54	0.88
3:D:977:ALA:HB2	3:J:831:GLY:HA3	1.56	0.88
2:C:502:PRO:HG3	2:C:510:THR:HG22	1.56	0.85

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	225/314 (72%)	206 (92%)	18 (8%)	1 (0%)	30	65
1	B	225/314 (72%)	205 (91%)	17 (8%)	3 (1%)	9	41

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	G	225/314 (72%)	205 (91%)	19 (8%)	1 (0%)	30	65
1	H	225/314 (72%)	205 (91%)	19 (8%)	1 (0%)	30	65
2	C	1108/1119 (99%)	1011 (91%)	94 (8%)	3 (0%)	36	70
2	I	1108/1119 (99%)	1012 (91%)	94 (8%)	2 (0%)	43	75
3	D	1486/1524 (98%)	1340 (90%)	141 (10%)	5 (0%)	36	70
3	J	1361/1524 (89%)	1235 (91%)	123 (9%)	3 (0%)	43	75
4	E	91/99 (92%)	83 (91%)	8 (9%)	0	100	100
4	K	91/99 (92%)	82 (90%)	9 (10%)	0	100	100
5	F	345/347 (99%)	316 (92%)	29 (8%)	0	100	100
5	L	345/347 (99%)	314 (91%)	31 (9%)	0	100	100
All	All	6835/7434 (92%)	6214 (91%)	602 (9%)	19 (0%)	36	70

5 of 19 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	765	GLN
3	D	1128	VAL
3	J	1128	VAL
1	A	53	VAL
1	B	178	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/270 (72%)	180 (93%)	14 (7%)	13	37
1	B	194/270 (72%)	178 (92%)	16 (8%)	10	34
1	G	194/270 (72%)	180 (93%)	14 (7%)	13	37
1	H	194/270 (72%)	178 (92%)	16 (8%)	10	34
2	C	931/936 (100%)	829 (89%)	102 (11%)	6	23
2	I	931/936 (100%)	829 (89%)	102 (11%)	6	23

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	D	1252/1281 (98%)	1123 (90%)	129 (10%)	7	25
3	J	1150/1281 (90%)	1042 (91%)	108 (9%)	8	29
4	E	83/88 (94%)	75 (90%)	8 (10%)	8	28
4	K	83/88 (94%)	75 (90%)	8 (10%)	8	28
5	F	297/299 (99%)	270 (91%)	27 (9%)	9	30
5	L	297/299 (99%)	270 (91%)	27 (9%)	9	30
All	All	5800/6288 (92%)	5229 (90%)	571 (10%)	7	27

5 of 571 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	J	534	ARG
3	J	685	ASP
3	J	480	GLU
3	J	1278	ASP
3	D	636	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 79 such sidechains are listed below:

Mol	Chain	Res	Type
2	I	1068	GLN
5	L	98	GLN
3	J	66	GLN
3	J	973	GLN
5	L	260	GLN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
8	Q	3/4 (75%)	1 (33%)	0
8	T	3/4 (75%)	1 (33%)	0
All	All	6/8 (75%)	2 (33%)	0

All (2) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
8	Q	2	C
8	T	2	C

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	227/314 (72%)	0.26	1 (0%) 88 75	68, 121, 176, 228	0
1	B	227/314 (72%)	0.13	4 (1%) 67 50	42, 95, 150, 180	0
1	G	227/314 (72%)	0.65	14 (6%) 26 24	96, 156, 194, 223	0
1	H	227/314 (72%)	0.40	5 (2%) 62 46	80, 131, 171, 192	0
2	C	1112/1119 (99%)	0.30	18 (1%) 70 52	30, 123, 203, 260	0
2	I	1112/1119 (99%)	0.51	40 (3%) 46 35	60, 146, 214, 259	0
3	D	1490/1524 (97%)	0.06	9 (0%) 85 71	22, 88, 155, 231	0
3	J	1367/1524 (89%)	0.24	16 (1%) 76 59	29, 112, 177, 240	0
4	E	93/99 (93%)	0.09	1 (1%) 78 60	23, 89, 154, 187	0
4	K	93/99 (93%)	0.39	1 (1%) 78 60	67, 126, 181, 236	0
5	F	347/347 (100%)	0.40	13 (3%) 45 34	52, 131, 221, 275	0
5	L	347/347 (100%)	0.58	23 (6%) 24 22	81, 151, 225, 270	0
6	O	48/48 (100%)	0.51	0 100 100	86, 193, 267, 293	0
6	R	48/48 (100%)	0.56	0 100 100	109, 184, 277, 300	0
7	P	48/48 (100%)	0.61	0 100 100	77, 199, 251, 283	0
7	S	43/48 (89%)	0.55	1 (2%) 61 45	100, 195, 234, 248	0
8	Q	4/4 (100%)	-0.23	0 100 100	116, 124, 128, 150	0
8	T	4/4 (100%)	-0.08	0 100 100	122, 130, 143, 154	0
All	All	7064/7634 (92%)	0.30	146 (2%) 63 47	22, 120, 200, 300	0

The worst 5 of 146 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
5	F	159	ILE	4.7
4	E	94	PRO	4.4
5	L	159	ILE	4.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
5	F	416	GLU	4.1
2	I	62	GLY	4.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
10	MG	D	2003	1/1	0.96	0.07	180,180,180,180	0
10	MG	J	2003	1/1	0.96	0.06	188,188,188,188	0
9	ZN	J	2002	1/1	0.97	0.06	166,166,166,166	0
9	ZN	D	2002	1/1	0.98	0.04	79,79,79,79	0
9	ZN	J	2001	1/1	0.99	0.05	97,97,97,97	0
9	ZN	D	2001	1/1	1.00	0.05	79,79,79,79	0

6.5 Other polymers [i](#)

There are no such residues in this entry.