



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2026 – 12:50 PM UTC

PDB ID : 5XL1 / pdb_00005xl1
Title : The structure of hemagglutinin from an avian-origin H4N6 influenza virus
Authors : Song, H.; Qi, J.; Gao, F.G.
Deposited on : 2017-05-10
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	327	0% 86% 10% ..
1	B	327	5% 84% 12% ..
2	C	176	2% 88% 8% ...
2	D	176	2% 89% 9% ..

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 8220 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Hemagglutinin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	319	Total	C	N	O	S	0	0	0
			2457	1538	437	470	12			
1	A	319	Total	C	N	O	S	0	0	0
			2457	1538	437	470	12			

- Molecule 2 is a protein called Hemagglutinin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	172	Total	C	N	O	S	0	0	0
			1404	871	250	279	4			
2	D	172	Total	C	N	O	S	0	0	0
			1404	871	250	279	4			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	78	Total	O	0	0
			78	78		
4	C	97	Total	O	0	0
			97	97		
4	A	161	Total	O	0	0
			161	161		
4	D	106	Total	O	0	0
			106	106		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hemagglutinin




- Molecule 1: Hemagglutinin



- Molecule 2: Hemagglutinin



- Molecule 2: Hemagglutinin



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	100.48Å 100.48Å 685.13Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.42 – 2.30 43.42 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.8 (43.42-2.30) 99.9 (43.42-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.46 (at 2.29Å)	Xtrriage
Refinement program	PHENIX 1.8.4_1496	Depositor
R, R_{free}	0.189 , 0.230 0.198 , 0.230	Depositor DCC
R_{free} test set	3038 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	35.8	Xtrriage
Anisotropy	0.207	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 48.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8220	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 45.84 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.2477e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/2511	0.76	0/3418
1	B	0.37	0/2511	0.76	0/3418
2	C	0.46	0/1428	0.77	1/1922 (0.1%)
2	D	0.46	0/1428	0.76	1/1922 (0.1%)
All	All	0.41	0/7878	0.76	2/10680 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
2	C	0	2
All	All	0	3

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	386	THR	N-CA-C	10.32	122.22	110.97
2	C	454	ARG	CB-CA-C	-6.26	109.34	116.54

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	138	ALA	Peptide
2	C	384	GLU	Peptide

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
2	C	386	THR	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2457	0	2402	23	0
1	B	2457	0	2402	34	0
2	C	1404	0	1320	14	0
2	D	1404	0	1320	14	0
3	A	28	0	26	1	0
3	B	28	0	26	0	0
4	A	161	0	0	6	0
4	B	78	0	0	8	0
4	C	97	0	0	1	0
4	D	106	0	0	6	0
All	All	8220	0	7496	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:18:ASN:OD1	4:B:701:HOH:O	1.88	0.90
1:A:28:ASP:O	4:A:701:HOH:O	1.90	0.89
1:A:244:THR:HG22	1:A:246:GLY:H	1.36	0.89
1:B:158:TYR:H	1:B:193:LYS:HE2	1.42	0.82
1:B:76:ASN:OD1	1:B:146:ARG:NH2	2.13	0.82
1:A:311:SER:O	4:A:702:HOH:O	1.99	0.79
1:B:190:ASN:O	4:B:702:HOH:O	2.01	0.77
1:B:5:GLY:HA2	2:C:468:PHE:HA	1.66	0.76
2:D:384:GLU:OE2	4:D:601:HOH:O	2.03	0.76
2:D:385:LYS:HD2	2:D:386:THR:N	2.05	0.72
1:B:319:ARG:NE	4:B:704:HOH:O	2.15	0.72
1:B:167:ASN:OD1	4:B:703:HOH:O	2.10	0.69

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:158:TYR:N	1:B:193:LYS:HE2	2.09	0.68
1:A:202:SER:OG	4:A:704:HOH:O	2.13	0.66
2:C:386:THR:OG1	2:C:387:ASN:N	2.29	0.65
1:B:11:MET:HE1	2:C:351:PHE:CE2	2.34	0.63
1:B:256:LYS:NZ	4:B:711:HOH:O	2.32	0.62
2:D:385:LYS:HD2	2:D:386:THR:H	1.66	0.61
2:D:369:GLN:OE1	4:D:603:HOH:O	2.16	0.61
2:D:450:ARG:NH2	4:D:606:HOH:O	2.34	0.61
2:C:388:ASP:OD1	4:C:601:HOH:O	2.17	0.60
1:A:322:PRO:O	1:A:323:GLU:HB2	2.01	0.60
1:A:147:LEU:HD12	1:A:249:ILE:HG22	1.84	0.58
1:B:319:ARG:HG2	1:B:319:ARG:HH11	1.69	0.58
1:A:319:ARG:HH11	1:A:319:ARG:HG2	1.69	0.57
1:A:11:MET:HE1	2:D:351:PHE:CE2	2.39	0.56
1:A:5:GLY:HA2	2:D:468:PHE:HA	1.86	0.56
2:C:380:ASN:O	2:C:384:GLU:HG2	2.06	0.55
1:B:137:ARG:O	1:B:138:ALA:HB3	2.07	0.54
1:A:76:ASN:OD1	1:A:146:ARG:NH2	2.39	0.54
1:B:46:GLU:OE1	1:B:273:GLY:HA3	2.08	0.53
1:B:6:ASN:ND2	4:B:715:HOH:O	2.41	0.53
1:B:11:MET:HE1	2:C:351:PHE:CD2	2.43	0.53
1:A:11:MET:HE1	2:D:351:PHE:CD2	2.45	0.52
2:D:354:GLN:HG3	2:D:359:THR:HG22	1.93	0.51
1:B:137:ARG:O	1:B:139:ASN:O	2.28	0.51
1:B:93:CYS:O	1:B:221:ARG:NH2	2.34	0.51
3:A:602:NAG:H83	4:A:731:HOH:O	2.12	0.50
1:B:289:THR:O	2:C:383:ILE:HG23	2.12	0.50
1:B:192:TYR:O	1:B:194:ASN:N	2.44	0.49
2:D:357:GLU:OE2	2:D:472:ASP:HB2	2.13	0.48
1:B:158:TYR:H	1:B:193:LYS:CE	2.19	0.47
1:B:52:LEU:HB2	4:B:736:HOH:O	2.15	0.46
1:A:319:ARG:HD3	4:D:652:HOH:O	2.16	0.46
1:A:217:ARG:HB2	1:A:224:SER:O	2.15	0.46
1:B:217:ARG:HB2	1:B:224:SER:O	2.16	0.46
1:B:161:GLN:HB2	1:B:244:THR:HG22	1.97	0.45
1:A:137:ARG:NE	1:A:142:ASP:OD2	2.46	0.45
1:A:14:HIS:CE1	4:A:784:HOH:O	2.69	0.45
1:A:164:THR:HG23	1:A:239:LEU:HD23	1.99	0.45
1:A:182:PRO:HG2	1:A:188:GLN:OE1	2.16	0.45
2:C:354:GLN:HG3	2:C:359:THR:HG22	1.99	0.45
2:D:451:ARG:HD2	4:D:607:HOH:O	2.17	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:6:ASN:H	1:A:7:PRO:CD	2.31	0.44
1:B:104:LEU:HB2	1:B:231:TRP:CZ3	2.54	0.43
1:A:138:ALA:N	4:A:716:HOH:O	2.51	0.43
1:A:64:ASN:HB3	1:A:69:SER:OG	2.18	0.43
2:D:490:ARG:O	2:D:494:ILE:HG12	2.19	0.43
1:B:303:CYS:O	2:C:387:ASN:HB3	2.19	0.42
1:B:5:GLY:N	4:B:722:HOH:O	2.53	0.42
1:B:175:TYR:CE2	1:B:254:HIS:HB3	2.55	0.42
1:B:182:PRO:HG2	1:B:188:GLN:OE1	2.20	0.42
1:B:139:ASN:HD22	1:B:140:VAL:H	1.66	0.41
2:C:468:PHE:O	2:C:493:ALA:HA	2.20	0.41
2:D:481:ASN:ND2	4:D:602:HOH:O	2.12	0.41
2:C:379:LEU:O	2:C:383:ILE:HG13	2.20	0.41
1:A:289:THR:O	2:D:383:ILE:HG23	2.20	0.41
2:C:462:ASN:OD1	2:C:462:ASN:N	2.53	0.41
1:B:11:MET:HB3	1:B:11:MET:HE3	1.72	0.41
1:B:139:ASN:O	1:B:140:VAL:CB	2.69	0.41
1:B:193:LYS:HB3	1:B:193:LYS:HE3	1.85	0.41
2:C:450:ARG:HB2	2:C:465:PHE:HZ	1.85	0.41
2:C:330:PHE:CE1	2:C:440:SER:HB2	2.55	0.40
1:B:94:TYR:CG	1:B:95:PRO:HD2	2.56	0.40
1:B:110:ASN:HA	1:B:263:SER:O	2.21	0.40
1:A:144:PHE:HB2	1:A:147:LEU:HB2	2.03	0.40
1:A:33:VAL:HG12	1:A:34:THR:HG23	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	317/327 (97%)	305 (96%)	10 (3%)	2 (1%)	21 27

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	317/327 (97%)	299 (94%)	15 (5%)	3 (1%)	14	17
2	C	170/176 (97%)	161 (95%)	8 (5%)	1 (1%)	21	27
2	D	170/176 (97%)	161 (95%)	9 (5%)	0	100	100
All	All	974/1006 (97%)	926 (95%)	42 (4%)	6 (1%)	21	27

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	6	ASN
2	C	386	THR
1	A	6	ASN
1	B	140	VAL
1	B	193	LYS
1	A	193	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	274/282 (97%)	266 (97%)	8 (3%)	37	55
1	B	274/282 (97%)	266 (97%)	8 (3%)	37	55
2	C	147/150 (98%)	145 (99%)	2 (1%)	59	76
2	D	147/150 (98%)	147 (100%)	0	100	100
All	All	842/864 (98%)	824 (98%)	18 (2%)	47	66

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	14	HIS
1	B	120	GLU
1	B	139	ASN
1	B	186	THR
1	B	193	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	239	LEU
1	B	243	ASN
1	B	319	ARG
2	C	379	LEU
2	C	440	SER
1	A	14	HIS
1	A	160	LEU
1	A	163	LEU
1	A	239	LEU
1	A	262	LYS
1	A	295	ILE
1	A	319	ARG
1	A	323	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	43	ASN
1	B	139	ASN
2	C	342	GLN
2	C	452	GLN
1	A	14	HIS
2	D	342	GLN
2	D	452	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	A	602	1	14,14,15	0.29	0	17,19,21	0.71	1 (5%)
3	NAG	B	602	1	14,14,15	0.38	0	17,19,21	0.49	0
3	NAG	B	601	1	14,14,15	0.43	0	17,19,21	0.41	0
3	NAG	A	601	1	14,14,15	0.28	0	17,19,21	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	602	1	-	2/6/23/26	0/1/1/1
3	NAG	B	602	1	-	2/6/23/26	0/1/1/1
3	NAG	B	601	1	-	2/6/23/26	0/1/1/1
3	NAG	A	601	1	-	1/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	602	NAG	C1-O5-C5	2.03	114.90	112.19

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	602	NAG	C8-C7-N2-C2
3	B	602	NAG	O7-C7-N2-C2
3	A	602	NAG	C8-C7-N2-C2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	602	NAG	O7-C7-N2-C2
3	B	601	NAG	O5-C5-C6-O6
3	A	601	NAG	O5-C5-C6-O6
3	B	601	NAG	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	602	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	319/327 (97%)	-0.14	4 (1%) 75 76	19, 37, 57, 81	0
1	B	319/327 (97%)	0.42	15 (4%) 36 38	22, 56, 80, 124	0
2	C	172/176 (97%)	-0.26	4 (2%) 61 63	22, 31, 51, 116	0
2	D	172/176 (97%)	-0.31	4 (2%) 61 63	17, 30, 46, 106	0
All	All	982/1006 (97%)	-0.01	27 (2%) 56 58	17, 38, 74, 124	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	386	THR	6.9
2	C	386	THR	5.1
2	C	385	LYS	4.8
2	D	383	ILE	4.6
2	D	385	LYS	4.3
1	B	163	LEU	4.1
2	C	383	ILE	4.0
1	B	138	ALA	3.8
1	B	5	GLY	3.6
1	B	157	ALA	2.8
1	B	123	TRP	2.8
1	A	323	GLU	2.8
1	A	5	GLY	2.7
1	B	137	ARG	2.7
1	B	221	ARG	2.7
1	B	197	GLY	2.7
1	B	222	GLY	2.7
2	D	499	GLN	2.6
1	A	157	ALA	2.4
1	A	26	ALA	2.3
1	B	323	GLU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	192	TYR	2.3
1	B	140	VAL	2.2
1	B	60	CYS	2.1
1	B	213	ASN	2.1
2	C	404	ILE	2.1
1	B	218	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	B	601	14/15	0.69	0.20	101,106,110,110	0
3	NAG	A	601	14/15	0.73	0.17	80,86,92,93	0
3	NAG	B	602	14/15	0.87	0.10	36,42,44,51	0
3	NAG	A	602	14/15	0.88	0.09	25,29,35,40	0

6.5 Other polymers [i](#)

There are no such residues in this entry.