



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 04:58 AM UTC

PDB ID : 1XM4 / pdb_00001xm4
Title : Catalytic Domain Of Human Phosphodiesterase 4B In Complex With Piclami-
last
Authors : Card, G.L.; England, B.P.; Suzuki, Y.; Fong, D.; Powell, B.; Lee, B.; Luu, C.;
Tabrizizad, M.; Gillette, S.; Ibrahim, P.N.; Artis, D.R.; Bollag, G.; Milburn,
M.V.; Kim, S.-H.; Schlessinger, J.; Zhang, K.Y.J.
Deposited on : 2004-10-01
Resolution : 2.31 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7754 (2.34-2.30)
Clashscore	190562	8383 (2.34-2.30)
Ramachandran outliers	187476	8303 (2.34-2.30)
Sidechain outliers	187428	8303 (2.34-2.30)
RSRZ outliers	180081	7760 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	398	 2% 65% 15% 19%
2	B	398	 7% 67% 12% 19%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5417 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cAMP-specific 3',5'-cyclic phosphodiesterase 4B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	323	Total	C	N	O	S	0	0	0
			2621	1655	441	504	21			

There are 23 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	131	MET	-	initiating methionine	UNP Q07343
A	132	GLY	-	cloning artifact	UNP Q07343
A	133	SER	-	cloning artifact	UNP Q07343
A	134	SER	-	cloning artifact	UNP Q07343
A	135	HIS	-	expression tag	UNP Q07343
A	136	HIS	-	expression tag	UNP Q07343
A	137	HIS	-	expression tag	UNP Q07343
A	138	HIS	-	expression tag	UNP Q07343
A	139	HIS	-	expression tag	UNP Q07343
A	140	HIS	-	expression tag	UNP Q07343
A	141	SER	-	cloning artifact	UNP Q07343
A	142	SER	-	cloning artifact	UNP Q07343
A	143	GLY	-	cloning artifact	UNP Q07343
A	144	LEU	-	cloning artifact	UNP Q07343
A	145	VAL	-	cloning artifact	UNP Q07343
A	146	PRO	-	cloning artifact	UNP Q07343
A	147	ARG	-	cloning artifact	UNP Q07343
A	148	GLY	-	cloning artifact	UNP Q07343
A	149	SER	-	cloning artifact	UNP Q07343
A	150	HIS	-	cloning artifact	UNP Q07343
A	151	MET	-	cloning artifact	UNP Q07343
A	194	CME	CYS	modified residue	UNP Q07343
A	432	CME	CYS	modified residue	UNP Q07343

- Molecule 2 is a protein called cAMP-specific 3',5'-cyclic phosphodiesterase 4B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	323	Total	C	N	O	S	0	0	0
			2621	1655	441	504	21			

There are 23 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	131	MET	-	initiating methionine	UNP Q07343
B	132	GLY	-	cloning artifact	UNP Q07343
B	133	SER	-	cloning artifact	UNP Q07343
B	134	SER	-	cloning artifact	UNP Q07343
B	135	HIS	-	expression tag	UNP Q07343
B	136	HIS	-	expression tag	UNP Q07343
B	137	HIS	-	expression tag	UNP Q07343
B	138	HIS	-	expression tag	UNP Q07343
B	139	HIS	-	expression tag	UNP Q07343
B	140	HIS	-	expression tag	UNP Q07343
B	141	SER	-	cloning artifact	UNP Q07343
B	142	SER	-	cloning artifact	UNP Q07343
B	143	GLY	-	cloning artifact	UNP Q07343
B	144	LEU	-	cloning artifact	UNP Q07343
B	145	VAL	-	cloning artifact	UNP Q07343
B	146	PRO	-	cloning artifact	UNP Q07343
B	147	ARG	-	cloning artifact	UNP Q07343
B	148	GLY	-	cloning artifact	UNP Q07343
B	149	SER	-	cloning artifact	UNP Q07343
B	150	HIS	-	cloning artifact	UNP Q07343
B	151	MET	-	cloning artifact	UNP Q07343
B	194	CME	CYS	modified residue	UNP Q07343
B	320	CME	CYS	modified residue	UNP Q07343

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		
3	B	1	Total	Zn	0	0
			1	1		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Mg	0	0
			1	1		

- Molecule 5 is 3-(CYCLOPENTYLOXY)-N-(3,5-DICHLOROPYRIDIN-4-YL)-4-METHOXYBENZAMIDE (CCD ID: PIL) (formula: C₁₈H₁₈Cl₂N₂O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	Cl	N	O	
			25	18	2	2	3	0
5	B	1	Total	C	Cl	N	O	
			25	18	2	2	3	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	66	Total	O	0	0
			66	66		
6	B	55	Total	O	0	0
			55	55		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: cAMP-specific 3',5'-cyclic phosphodiesterase 4B



- Molecule 2: cAMP-specific 3',5'-cyclic phosphodiesterase 4B



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	89.57Å 94.32Å 106.68Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	70.71 – 2.31 70.66 – 2.31	Depositor EDS
% Data completeness (in resolution range)	97.6 (70.71-2.31) 97.6 (70.66-2.31)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.06 (at 2.32Å)	Xtriage
Refinement program	REFMAC 5.1.25	Depositor
R, R_{free}	0.212 , 0.263 (Not available) , 0.288	Depositor DCC
R_{free} test set	1971 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	40.6	Xtriage
Anisotropy	0.422	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 45.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.009 for k,h,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5417	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 30.21 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.3587e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN, PIL, CME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.05	4/2654 (0.2%)	1.12	5/3597 (0.1%)
2	B	1.07	5/2654 (0.2%)	1.09	4/3597 (0.1%)
All	All	1.06	9/5308 (0.2%)	1.11	9/7194 (0.1%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	324	MET	SD-CE	7.30	1.97	1.79
2	B	296	MET	SD-CE	6.23	1.95	1.79
2	B	351	MET	SD-CE	5.91	1.94	1.79
1	A	385	ARG	C-O	-5.74	1.17	1.24
1	A	431	MET	SD-CE	5.59	1.93	1.79
1	A	351	MET	SD-CE	5.46	1.93	1.79
1	A	243	ALA	CA-CB	-5.44	1.44	1.53
2	B	366	VAL	CA-CB	5.40	1.60	1.54
2	B	483	MET	SD-CE	5.18	1.92	1.79

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	484	ILE	CB-CA-C	-8.34	103.19	110.94
1	A	227	TYR	N-CA-C	-6.54	100.80	110.48
2	B	274	HIS	N-CA-C	6.53	120.35	112.38
1	A	183	VAL	N-CA-CB	5.49	118.00	110.54
1	A	274	HIS	N-CA-C	5.42	119.38	112.34
2	B	326	LEU	N-CA-C	-5.38	103.43	110.53
1	A	232	ALA	N-CA-C	5.29	117.12	111.36
2	B	465	ASP	N-CA-C	5.06	117.45	111.33
2	B	484	ILE	CB-CA-C	-5.04	105.31	110.71

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2621	0	2551	25	0
2	B	2621	0	2551	27	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	25	0	18	1	0
5	B	25	0	18	0	0
6	A	66	0	0	0	0
6	B	55	0	0	1	0
All	All	5417	0	5138	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:328:LYS:HZ3	2:B:331:ARG:NH2	1.86	0.72
1:A:340:ASP:OD1	1:A:385:ARG:HD3	1.92	0.69
1:A:252:THR:HG22	1:A:465:ASP:OD2	1.93	0.69
2:B:255:LEU:HD23	2:B:372:LEU:HD12	1.81	0.62
1:A:327:THR:HG22	1:A:330:GLN:H	1.67	0.60
2:B:340:ASP:OD1	2:B:385:ARG:HD3	2.02	0.60
2:B:356:ASP:HB3	2:B:382:GLN:HE22	1.69	0.57
2:B:255:LEU:CD2	2:B:372:LEU:HD12	2.34	0.57
1:A:296:MET:HE3	2:B:313:LYS:NZ	2.21	0.56
2:B:328:LYS:NZ	2:B:331:ARG:NH2	2.52	0.56
1:A:397:THR:HB	1:A:469:ILE:HG23	1.89	0.54
1:A:381:ILE:O	1:A:385:ARG:HG3	2.08	0.53
2:B:397:THR:HB	2:B:469:ILE:HG23	1.89	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:327:THR:HG22	2:B:328:LYS:N	2.23	0.53
1:A:356:ASP:HB3	1:A:382:GLN:NE2	2.24	0.52
2:B:270:ALA:O	2:B:274:HIS:HB3	2.12	0.49
1:A:180:ILE:HA	1:A:183:VAL:HG13	1.93	0.49
1:A:390:CYS:HB3	1:A:455:TRP:CZ2	2.47	0.49
1:A:270:ALA:O	1:A:274:HIS:HB3	2.13	0.49
2:B:300:GLU:O	2:B:301:SER:C	2.55	0.48
1:A:376:ASN:CG	1:A:377:TYR:H	2.22	0.47
2:B:328:LYS:HZ3	2:B:331:ARG:HH21	1.59	0.47
1:A:296:MET:HE3	2:B:313:LYS:HZ1	1.79	0.46
1:A:340:ASP:OD1	1:A:385:ARG:CD	2.63	0.46
2:B:173:LEU:HA	2:B:178:LEU:HD13	1.97	0.45
2:B:397:THR:HG22	2:B:473:LEU:HD13	1.99	0.45
2:B:303:LEU:HD12	6:B:119:HOH:O	2.17	0.45
1:A:443:GLN:NE2	5:A:1003:PIL:H81	2.32	0.44
2:B:376:ASN:CG	2:B:377:TYR:H	2.25	0.44
1:A:275:ASP:O	1:A:278:HIS:HB2	2.18	0.44
2:B:213:SER:O	2:B:217:ILE:HG12	2.19	0.43
2:B:258:VAL:HG11	2:B:374:LEU:HD12	2.00	0.43
1:A:296:MET:HE3	1:A:296:MET:HB2	1.82	0.43
2:B:481:GLN:HA	2:B:484:ILE:HD12	2.01	0.43
1:A:452:HIS:HB3	1:A:453:PRO:HD3	2.01	0.43
2:B:219:TYR:CD2	2:B:219:TYR:C	2.96	0.42
1:A:340:ASP:OD1	1:A:340:ASP:C	2.63	0.42
2:B:180:ILE:HA	2:B:183:VAL:HG13	2.01	0.41
2:B:275:ASP:HA	2:B:307:HIS:CD2	2.55	0.41
1:A:258:VAL:HG11	1:A:374:LEU:HD12	2.01	0.41
1:A:234:HIS:ND1	1:A:413:GLU:OE2	2.38	0.41
1:A:459:ALA:O	1:A:463:GLN:HB3	2.20	0.41
2:B:252:THR:HG22	2:B:255:LEU:HD12	2.01	0.41
1:A:326:LEU:HD23	1:A:326:LEU:N	2.35	0.41
2:B:293:LEU:HD23	2:B:293:LEU:HA	1.92	0.41
1:A:364:LYS:NZ	1:A:460:ASP:O	2.52	0.41
1:A:323:PHE:HB3	1:A:326:LEU:HG	2.02	0.41
2:B:381:ILE:O	2:B:385:ARG:HG3	2.21	0.41
1:A:263:GLU:HG2	1:A:337:MET:SD	2.61	0.41
2:B:327:THR:HG22	2:B:329:LYS:H	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	319/398 (80%)	309 (97%)	9 (3%)	1 (0%)	36	45
2	B	319/398 (80%)	309 (97%)	10 (3%)	0	100	100
All	All	638/796 (80%)	618 (97%)	19 (3%)	1 (0%)	43	53

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	375	ASP

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	291/358 (81%)	270 (93%)	21 (7%)	13	17
2	B	291/358 (81%)	276 (95%)	15 (5%)	21	30
All	All	582/716 (81%)	546 (94%)	36 (6%)	16	23

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	178	LEU
1	A	183	VAL
1	A	207	LYS
1	A	231	VAL
1	A	295	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	296	MET
1	A	313	LYS
1	A	316	GLN
1	A	318	GLU
1	A	319	HIS
1	A	327	THR
1	A	328	LYS
1	A	329	LYS
1	A	331	ARG
1	A	353	LEU
1	A	366	VAL
1	A	368	SER
1	A	373	LEU
1	A	416	GLN
1	A	424	ARG
1	A	429	SER
2	B	183	VAL
2	B	319	HIS
2	B	324	MET
2	B	326	LEU
2	B	328	LYS
2	B	353	LEU
2	B	364	LYS
2	B	365	LYS
2	B	366	VAL
2	B	368	SER
2	B	371	VAL
2	B	401	GLU
2	B	428	ILE
2	B	429	SER
2	B	436	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	443	GLN
2	B	376	ASN
2	B	382	GLN
2	B	481	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CME	A	194	1	8,9,10	0.99	0	6,9,11	1.63	2 (33%)
2	CME	B	194	2	8,9,10	0.86	0	6,9,11	1.46	1 (16%)
1	CME	A	432	1	8,9,10	0.77	0	6,9,11	1.23	0
2	CME	B	320	2	8,9,10	0.92	0	6,9,11	1.03	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CME	A	194	1	-	1/5/8/10	-
2	CME	B	194	2	-	0/5/8/10	-
1	CME	A	432	1	-	2/5/8/10	-
2	CME	B	320	2	-	3/5/8/10	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	194	CME	CE-SD-SG	3.09	117.02	103.46
2	B	194	CME	CE-SD-SG	2.61	114.91	103.46
1	A	194	CME	CZ-CE-SD	-2.00	106.69	113.39

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	194	CME	CE-SD-SG-CB
1	A	432	CME	CE-SD-SG-CB
1	A	432	CME	SD-CE-CZ-OH
2	B	320	CME	SD-CE-CZ-OH
2	B	320	CME	CZ-CE-SD-SG
2	B	320	CME	CA-CB-SG-SD

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	PIL	B	1003	-	27,27,27	0.84	1 (3%)	37,37,37	1.68	6 (16%)
5	PIL	A	1003	-	27,27,27	0.88	1 (3%)	37,37,37	1.81	5 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PIL	B	1003	-	-	1/14/21/21	0/3/3/3
5	PIL	A	1003	-	-	0/14/21/21	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	1003	PIL	C19-CL20	2.27	1.79	1.73
5	A	1003	PIL	C18-C19	-2.27	1.36	1.40

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1003	PIL	C23-C24-C18	6.67	123.82	120.16
5	A	1003	PIL	C24-C23-N22	-5.74	117.69	122.80
5	B	1003	PIL	C24-C23-N22	-4.80	118.53	122.80
5	B	1003	PIL	C11-O10-C9	4.50	124.11	117.51
5	B	1003	PIL	C23-C24-C18	3.90	122.30	120.16
5	A	1003	PIL	C11-O10-C9	3.07	122.02	117.51
5	B	1003	PIL	C21-C19-C18	-2.92	118.56	120.16
5	A	1003	PIL	C21-N22-C23	2.72	121.21	117.51
5	B	1003	PIL	C21-N22-C23	2.64	121.10	117.51
5	A	1003	PIL	C19-C21-N22	-2.39	120.67	122.80
5	B	1003	PIL	C18-C19-CL20	2.28	121.87	119.28

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	1003	PIL	C9-C2-O3-C4


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1003	PIL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	321/398 (80%)	0.25	6 (1%) 66 68	16, 22, 30, 39	0
2	B	321/398 (80%)	0.84	29 (9%) 15 17	16, 22, 30, 39	0
All	All	642/796 (80%)	0.55	35 (5%) 30 33	16, 22, 30, 39	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	315	LEU	4.5
2	B	316	GLN	3.3
2	B	319	HIS	3.3
2	B	376	ASN	3.3
1	A	369	SER	3.1
2	B	287	ILE	3.1
2	B	286	LEU	3.0
2	B	468	ASP	2.9
2	B	204	ASP	2.7
2	B	471	ASP	2.7
2	B	326	LEU	2.6
2	B	371	VAL	2.5
1	A	376	ASN	2.5
2	B	366	VAL	2.5
1	A	164	ASP	2.5
2	B	294	ALA	2.5
1	A	429	SER	2.4
1	A	356	ASP	2.4
1	A	319	HIS	2.4
2	B	327	THR	2.3
2	B	465	ASP	2.3
2	B	212	SER	2.2
2	B	251	SER	2.2
2	B	177	GLY	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	447	ILE	2.1
2	B	462	VAL	2.1
2	B	474	GLU	2.1
2	B	292	GLU	2.1
2	B	324	MET	2.1
2	B	223	LEU	2.1
2	B	164	ASP	2.1
2	B	231	VAL	2.1
2	B	363	THR	2.0
2	B	305	ASN	2.0
2	B	276	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CME	A	432	10/11	0.88	0.16	38,43,62,63	0
1	CME	A	194	10/11	0.93	0.12	30,35,57,57	0
2	CME	B	320	10/11	0.93	0.14	40,43,63,63	0
2	CME	B	194	10/11	0.94	0.11	29,31,54,54	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	PIL	A	1003	25/25	0.91	0.12	35,43,54,55	0
5	PIL	B	1003	25/25	0.93	0.11	41,48,57,58	0
3	ZN	B	1001	1/1	0.99	0.05	38,38,38,38	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	MG	B	1002	1/1	0.99	0.07	30,30,30,30	0
4	MG	A	1002	1/1	1.00	0.05	26,26,26,26	0
3	ZN	A	1001	1/1	1.00	0.03	35,35,35,35	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.