



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 30, 2026 – 03:52 AM UTC

PDB ID : 4XMP / pdb_00004xmp
Title : Crystal structure of broadly and potently neutralizing antibody VRC08 in complex with HIV-1 clade A strain Q842.d12 gp120
Authors : Zhou, T.; Srivatsan, S.; Kwong, P.D.
Deposited on : 2015-01-15
Resolution : 1.78 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1365 (1.78-1.78)
Clashscore	190562	1395 (1.78-1.78)
Ramachandran outliers	187476	1382 (1.78-1.78)
Sidechain outliers	187428	1382 (1.78-1.78)
RSRZ outliers	180081	1365 (1.78-1.78)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	G	354	11% 91% 5%
2	H	235	11% 95% • •
3	L	211	6% 99% •

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 12888 atoms, of which 6046 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Envelope glycoprotein gp160,Envelope glycoprotein gp160,Envelope glycoprotein gp160.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	G	337	Total	C	H	N	O	S	0	2	0
			5269	1680	2587	476	505	21			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	124	GLY	-	linker	UNP Q8JDI3
G	198	GLY	-	linker	UNP Q8JDI3
G	318	GLY	-	linker	UNP Q8JDI3
G	319	GLY	-	linker	UNP Q8JDI3
G	320	SER	-	linker	UNP Q8JDI3
G	321	GLY	-	linker	UNP Q8JDI3
G	322	SER	-	linker	UNP Q8JDI3
G	323	GLY	-	linker	UNP Q8JDI3


- Molecule 2 is a protein called HEAVY CHAIN OF ANTIBODY VRC08.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	H	232	Total	C	H	N	O	S	0	0	0
			3460	1103	1708	306	331	12			

- Molecule 3 is a protein called LIGHT CHAIN OF ANTIBODY VRC08.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	L	211	Total	C	H	N	O	S	0	0	0
			3202	1008	1588	277	322	7			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	G	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
4	G	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
4	G	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
4	G	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
4	G	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
4	G	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
4	G	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
4	G	1	Total	C	H	N	O	0	0
			27	8	13	1	5		

- Molecule 5 is (R,R)-2,3-BUTANEDIOL (CCD ID: BU3) (formula: C₄H₁₀O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	G	1	Total	C	H	O	0	0
			16	4	10	2		
5	G	1	Total	C	H	O	0	0
			16	4	10	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	G	247	Total	O	0	0
			247	247		
6	H	197	Total	O	0	0
			197	197		
6	L	184	Total	O	0	0
			184	184		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Envelope glycoprotein gp160,Envelope glycoprotein gp160,Envelope glycoprotein gp160



- Molecule 2: HEAVY CHAIN OF ANTIBODY VRC08



- Molecule 3: LIGHT CHAIN OF ANTIBODY VRC08



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	69.71Å 82.50Å 163.07Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.50 – 1.78 35.50 – 1.78	Depositor EDS
% Data completeness (in resolution range)	97.6 (35.50-1.78) 97.5 (35.50-1.78)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$	-	Xtrriage
Refinement program	PHENIX dev_1702	Depositor
R, R_{free}	0.165 , 0.201 0.169 , 0.204	Depositor DCC
R_{free} test set	4414 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å ²)	(Not available)	Xtrriage
Anisotropy	(Not available)	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 51.7	EDS
L-test for twinning ¹	$\langle L \rangle =$ (Not available), $\langle L^2 \rangle =$ (Not available)	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	12888	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *(Not available)*

¹Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BU3, NAG, PCA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	G	0.47	0/2746	0.66	0/3729
2	H	0.48	0/1788	0.67	0/2433
3	L	0.47	0/1645	0.70	0/2230
All	All	0.47	0/6179	0.67	0/8392

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	2682	2587	2608	10	0
2	H	1752	1708	1710	3	0
3	L	1614	1588	1589	1	0
4	G	154	143	143	2	0
5	G	12	20	20	0	0
6	G	247	0	0	5	1
6	H	197	0	0	0	0
6	L	184	0	0	0	1
All	All	6842	6046	6070	13	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 1.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:211[A]:GLU:OE2	6:G:660:HOH:O	2.07	0.72
1:G:344:GLN:NE2	6:G:791:HOH:O	2.26	0.66
1:G:62:GLU:OE1	6:G:601:HOH:O	2.18	0.57
1:G:357:LYS:NZ	6:G:839:HOH:O	2.41	0.53
1:G:350:ARG:NH2	1:G:395:TRP:O	2.43	0.52
1:G:460:ASN:OD1	2:H:61:ARG:NH2	2.45	0.49
1:G:295:ASN:HD22	4:G:504:NAG:H83	1.78	0.48
1:G:446:GLU:OE1	4:G:504:NAG:O6	2.31	0.47
3:L:4:VAL:CG1	3:L:88:CYS:SG	3.06	0.44
1:G:211[B]:GLU:OE2	6:G:602:HOH:O	2.21	0.43
2:H:178:LEU:C	2:H:178:LEU:HD12	2.45	0.41
2:H:48:MET:HE2	2:H:67:VAL:HG11	2.02	0.40
1:G:83:GLU:HG2	1:G:245:VAL:HG12	2.02	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:G:604:HOH:O	6:L:333:HOH:O[2_775]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	G	333/354 (94%)	322 (97%)	11 (3%)	0	100	100
2	H	228/235 (97%)	224 (98%)	4 (2%)	0	100	100
3	L	209/211 (99%)	208 (100%)	1 (0%)	0	100	100
All	All	770/800 (96%)	754 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	302/312 (97%)	298 (99%)	4 (1%)	61	46
2	H	194/197 (98%)	190 (98%)	4 (2%)	47	27
3	L	182/182 (100%)	182 (100%)	0	100	100
All	All	678/691 (98%)	670 (99%)	8 (1%)	63	49

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	371	ILE
1	G	398	THR
1	G	460	ASN
1	G	462	ASN
2	H	11	MET
2	H	13	GLU
2	H	129	LYS
2	H	140	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	G	377	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PCA	H	1	2	7,8,9	2.28	2 (28%)	9,10,12	2.54	5 (55%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PCA	H	1	2	-	0/0/11/13	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	1	PCA	CD-N	4.90	1.46	1.34
2	H	1	PCA	CA-N	3.30	1.50	1.46

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	1	PCA	CB-CA-C	-4.65	106.27	112.66
2	H	1	PCA	CA-N-CD	-3.21	102.59	113.58
2	H	1	PCA	OE-CD-CG	-2.83	121.67	126.72
2	H	1	PCA	CB-CA-N	2.75	110.80	103.24
2	H	1	PCA	CG-CD-N	2.50	114.50	108.39

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	G	504	1	14,14,15	0.44	0	17,19,21	0.62	0
4	NAG	G	510	1	14,14,15	0.33	0	17,19,21	0.46	0
4	NAG	G	503	1	14,14,15	0.33	0	17,19,21	0.39	0
4	NAG	G	502	1	14,14,15	0.55	0	17,19,21	0.68	0
5	BU3	G	512	-	4,5,5	1.32	0	6,6,6	0.26	0
4	NAG	G	507	1	14,14,15	0.24	0	17,19,21	0.41	0
4	NAG	G	509	1	14,14,15	0.42	0	17,19,21	0.51	0
4	NAG	G	511	1	14,14,15	0.29	0	17,19,21	0.43	0
5	BU3	G	513	-	4,5,5	1.48	1 (25%)	6,6,6	0.37	0
4	NAG	G	506	1	14,14,15	0.27	0	17,19,21	0.58	0
4	NAG	G	508	1	14,14,15	0.31	0	17,19,21	0.39	0
4	NAG	G	501	1	14,14,15	0.30	0	17,19,21	0.43	0
4	NAG	G	505	1	14,14,15	0.51	0	17,19,21	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	G	504	1	-	2/6/23/26	0/1/1/1
4	NAG	G	510	1	-	0/6/23/26	0/1/1/1
4	NAG	G	503	1	-	0/6/23/26	0/1/1/1
4	NAG	G	502	1	-	0/6/23/26	0/1/1/1
5	BU3	G	512	-	-	4/4/4/4	-
4	NAG	G	507	1	-	2/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	G	509	1	-	0/6/23/26	0/1/1/1
4	NAG	G	511	1	-	1/6/23/26	0/1/1/1
5	BU3	G	513	-	-	0/4/4/4	-
4	NAG	G	506	1	-	1/6/23/26	0/1/1/1
4	NAG	G	508	1	-	0/6/23/26	0/1/1/1
4	NAG	G	501	1	-	0/6/23/26	0/1/1/1
4	NAG	G	505	1	-	2/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	G	513	BU3	O6-C3	-2.03	1.37	1.43

There are no bond angle outliers.

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	G	512	BU3	C1-C2-C3-C4
4	G	504	NAG	C8-C7-N2-C2
4	G	504	NAG	O7-C7-N2-C2
4	G	506	NAG	O5-C5-C6-O6
4	G	505	NAG	O5-C5-C6-O6
4	G	505	NAG	C4-C5-C6-O6
5	G	512	BU3	C1-C2-C3-O6
5	G	512	BU3	O5-C2-C3-C4
4	G	507	NAG	C4-C5-C6-O6
4	G	507	NAG	O5-C5-C6-O6
5	G	512	BU3	O5-C2-C3-O6
4	G	511	NAG	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	G	504	NAG	2	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	G	337/354 (95%)	0.41	38 (11%)	10 11	24, 40, 117, 177	2 (0%)
2	H	231/235 (98%)	0.19	25 (10%)	11 12	18, 36, 113, 180	0
3	L	211/211 (100%)	0.12	13 (6%)	26 32	23, 40, 92, 158	0
All	All	779/800 (97%)	0.27	76 (9%)	13 15	18, 39, 110, 180	2 (0%)

All (76) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	445	CYS	6.6
2	H	132	SER	6.2
2	H	130	SER	6.0
1	G	399	ALA	5.8
2	H	133	GLY	5.6
1	G	45	TRP	5.4
2	H	134	GLY	5.0
1	G	122	LEU	4.8
1	G	397	HIS	4.8
1	G	461	THR	4.6
3	L	214	CYS	4.6
1	G	200	ALA	4.4
1	G	123	THR	4.4
1	G	460	ASN	4.2
1	G	419	ARG	4.0
1	G	205	CYS	4.0
1	G	398	THR	4.0
2	H	191	THR	4.0
1	G	72	HIS	3.5
1	G	463	SER	3.4
2	H	126	PRO	3.4
1	G	228	CYS	3.3
1	G	239	CYS	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	H	100(C)	ARG	3.3
2	H	189	LEU	3.3
2	H	187	SER	3.2
3	L	129	THR	3.2
1	G	491	ILE	3.2
1	G	124	GLY	3.2
1	G	79	PRO	3.1
3	L	37	ARG	3.1
2	H	215	SER	3.1
2	H	135	THR	3.1
3	L	7	SER	3.1
2	H	140	CYS	3.0
2	H	100(E)	CYS	2.9
2	H	74	TYR	2.9
1	G	201	ILE	2.9
1	G	462	ASN	2.8
3	L	154	LEU	2.8
1	G	231	GLU	2.8
3	L	53	GLU	2.8
2	H	99	CYS	2.7
2	H	188	SER	2.7
2	H	61	ARG	2.6
2	H	131	THR	2.6
1	G	80	ASN	2.6
1	G	339	ASN	2.6
1	G	464	THR	2.6
2	H	100(B)	ARG	2.6
1	G	202	THR	2.5
1	G	198	GLY	2.5
1	G	473	GLY	2.5
3	L	147	GLN	2.5
1	G	396	ASN	2.4
3	L	165	GLU	2.4
1	G	199	SER	2.4
3	L	42	GLN	2.4
3	L	126	LYS	2.4
1	G	48	ALA	2.4
2	H	208	ASP	2.4
3	L	185	ASP	2.4
2	H	129	LYS	2.4
1	G	83	GLU	2.3
1	G	97	LYS	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	G	121	LYS	2.3
2	H	193	THR	2.3
3	L	17	GLU	2.2
3	L	152	ASN	2.2
2	H	115	SER	2.2
1	G	412	ASP	2.2
1	G	81	PRO	2.2
2	H	205	THR	2.1
1	G	444	ARG	2.1
1	G	423	ILE	2.1
2	H	195	ILE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PCA	H	1	8/9	0.96	0.07	28,54,65,83	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	G	506	14/15	0.57	0.17	82,110,132,140	0
4	NAG	G	510	14/15	0.65	0.15	79,98,117,121	0
4	NAG	G	511	14/15	0.73	0.17	74,98,126,126	0
4	NAG	G	507	14/15	0.74	0.14	79,96,115,118	0
4	NAG	G	501	14/15	0.75	0.14	84,99,121,121	0
5	BU3	G	512	6/6	0.78	0.25	66,81,96,100	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	G	504	14/15	0.81	0.21	64,94,114,117	0
5	BU3	G	513	6/6	0.81	0.26	70,84,100,103	0
4	NAG	G	505	14/15	0.82	0.14	56,87,105,111	0
4	NAG	G	508	14/15	0.94	0.08	29,38,65,86	0
4	NAG	G	503	14/15	0.95	0.07	26,39,67,67	0
4	NAG	G	502	14/15	0.95	0.07	23,29,55,70	0
4	NAG	G	509	14/15	0.96	0.06	20,33,43,49	0

6.5 Other polymers [i](#)

There are no such residues in this entry.