



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 03:21 AM UTC

PDB ID : 5XM7 / pdb_00005xm7
Title : Crystal structure of Plasmodium falciparum aminopeptidase N in complex with (S)-2-((2S,3R)-3-amino-2-hydroxy-4-phenylbutanamido)-N-hydroxy-4-methylpentanamide
Authors : Marapaka, A.K.; Zhang, Y.; Addlagatta, A.
Deposited on : 2017-05-12
Resolution : 1.96 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.96 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	914	 90% 7%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7825 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called M1 family aminopeptidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	890	Total	C	N	O	S	0	8	0
			7345	4717	1196	1405	27			

There are 23 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	172	MET	-	expression tag	UNP O96935
A	173	GLY	-	expression tag	UNP O96935
A	174	SER	-	expression tag	UNP O96935
A	175	SER	-	expression tag	UNP O96935
A	176	HIS	-	expression tag	UNP O96935
A	177	HIS	-	expression tag	UNP O96935
A	178	HIS	-	expression tag	UNP O96935
A	179	HIS	-	expression tag	UNP O96935
A	180	HIS	-	expression tag	UNP O96935
A	181	HIS	-	expression tag	UNP O96935
A	182	SER	-	expression tag	UNP O96935
A	183	SER	-	expression tag	UNP O96935
A	184	GLY	-	expression tag	UNP O96935
A	185	LEU	-	expression tag	UNP O96935
A	186	VAL	-	expression tag	UNP O96935
A	187	PRO	-	expression tag	UNP O96935
A	188	ARG	-	expression tag	UNP O96935
A	189	GLY	-	expression tag	UNP O96935
A	190	SER	-	expression tag	UNP O96935
A	191	HIS	-	expression tag	UNP O96935
A	192	MET	-	expression tag	UNP O96935
A	193	ALA	-	expression tag	UNP O96935
A	194	SER	-	expression tag	UNP O96935


- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	Mg	0	0
			3	3		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is (2S)-4-methyl-N-[(1R)-2-(oxidanylamino)-2-oxidanylidene-1-phenyl-ethyl]-2-[(phenylmethyl)carbamoylamino]pentanamide (CCD ID: 89X) (formula: C₂₂H₂₈N₄O₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			30	22	4	4		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	380	Total	O	0	0
			380	380		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: M1 family aminopeptidase

Chain A:  90% 7% .



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	74.46Å 109.10Å 112.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.98 – 1.96 24.98 – 1.96	Depositor EDS
% Data completeness (in resolution range)	99.5 (24.98-1.96) 99.5 (24.98-1.96)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.35 (at 1.96Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.168 , 0.219 0.178 , 0.226	Depositor DCC
R_{free} test set	3360 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	29.8	Xtriage
Anisotropy	0.032	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 39.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.007 for -h,l,k	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7825	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, GOL, ZN, 89X

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.21	10/7533 (0.1%)	1.06	5/10183 (0.0%)

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	449	LEU	N-CA	8.60	1.56	1.46
1	A	237	LYS	C-O	-7.15	1.14	1.24
1	A	283	VAL	C-O	-6.58	1.16	1.24
1	A	859	LYS	C-O	-6.40	1.15	1.24
1	A	623	ALA	C-O	6.32	1.31	1.24
1	A	887	VAL	C-O	6.11	1.30	1.24
1	A	411	TYR	CA-C	5.63	1.59	1.52
1	A	674	TYR	C-O	-5.25	1.17	1.23
1	A	658	TYR	N-CA	5.23	1.52	1.45
1	A	1032	ASN	C-O	5.02	1.29	1.24

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1067	GLU	CA-C-N	-5.88	113.91	119.85
1	A	1067	GLU	C-N-CA	-5.88	113.91	119.85
1	A	574	PHE	CB-CA-C	5.61	120.05	110.56
1	A	986	VAL	N-CA-C	5.36	115.85	111.62
1	A	458	ASN	N-CA-C	5.08	117.71	111.82

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7345	0	7281	28	1
2	A	1	0	0	0	0
3	A	3	0	0	0	0
4	A	66	0	87	11	0
5	A	30	0	0	0	0
6	A	380	0	0	8	0
All	All	7825	0	7368	31	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:420:SER:O	6:A:1201:HOH:O	2.06	0.72
1:A:668:SER:OG	4:A:1107:GOL:H11	1.89	0.72
1:A:715:GLU:HG2	4:A:1112:GOL:O3	1.93	0.68
1:A:545:ASP:OD1	1:A:765:TYR:OH	2.16	0.64
1:A:811:LEU:HB3	1:A:867:MET:SD	2.42	0.60
1:A:834:ILE:HG12	1:A:840:LEU:HD22	1.88	0.54
1:A:659:ASN:HB3	4:A:1106:GOL:H11	1.91	0.51
1:A:306:GLY:O	1:A:316:SER:HA	2.11	0.51
1:A:918:ILE:HD12	1:A:941:LYS:HE2	1.93	0.51
1:A:658:TYR:OH	1:A:732:LEU:HD11	2.10	0.50
4:A:1109:GOL:C2	6:A:1214:HOH:O	2.60	0.49
1:A:423:GLN:NE2	6:A:1201:HOH:O	2.46	0.48
1:A:670:HIS:HB2	4:A:1107:GOL:H12	1.98	0.46
1:A:538:THR:HG21	1:A:819:GLY:HA3	1.97	0.45
1:A:620:PHE:CZ	1:A:624:MET:HE3	2.51	0.45
1:A:977:GLU:HG2	6:A:1530:HOH:O	2.14	0.45
4:A:1109:GOL:H2	6:A:1214:HOH:O	2.16	0.45
1:A:553:LEU:C	1:A:553:LEU:HD23	2.40	0.45
1:A:777:LEU:HD22	4:A:1114:GOL:O1	2.16	0.45
4:A:1110:GOL:H12	6:A:1223:HOH:O	2.16	0.44
1:A:778:MET:HE2	6:A:1557:HOH:O	2.17	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:969:ARG:HB3	1:A:1002:PRO:HB3	2.01	0.42
1:A:907:LYS:HG2	4:A:1105:GOL:C1	2.50	0.41
1:A:1046:TRP:O	1:A:1054:GLN:HG2	2.20	0.41
1:A:513:PHE:O	1:A:564:ARG:HG3	2.20	0.41
1:A:666:GLN:OE1	4:A:1106:GOL:H31	2.19	0.41
1:A:904:LEU:HD23	4:A:1105:GOL:H2	2.03	0.41
1:A:213:ASN:ND2	6:A:1205:HOH:O	2.45	0.41
1:A:550:VAL:HG12	1:A:577:THR:HG21	2.03	0.40
1:A:1046:TRP:CD1	1:A:1061:MET:HE1	2.56	0.40
1:A:1046:TRP:CG	1:A:1061:MET:HE1	2.57	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:723:ASN:ND2	1:A:792:GLU:OE1[4_545]	2.14	0.06

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	896/914 (98%)	880 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	830/842 (99%)	817 (98%)	13 (2%)	55	52

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	205	LYS
1	A	287	LYS
1	A	555	ASP
1	A	572	GLU
1	A	608	LYS
1	A	613	ASN
1	A	745	ASN
1	A	790	LYS
1	A	829	GLN
1	A	941	LYS
1	A	969	ARG
1	A	1045	LEU
1	A	1085	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	468	ASN
1	A	670	HIS
1	A	775	GLN
1	A	779	ASN
1	A	1005	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 4 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	A	1107	-	5,5,5	1.80	1 (20%)	5,5,5	1.62	1 (20%)
4	GOL	A	1112	-	5,5,5	0.49	0	5,5,5	0.67	0
4	GOL	A	1114	-	5,5,5	0.58	0	5,5,5	1.34	1 (20%)
4	GOL	A	1110	-	5,5,5	0.47	0	5,5,5	0.78	0
4	GOL	A	1108	-	5,5,5	0.91	0	5,5,5	1.17	0
4	GOL	A	1106	-	5,5,5	1.23	1 (20%)	5,5,5	2.59	1 (20%)
5	89X	A	1116	2	30,31,31	1.12	3 (10%)	38,40,40	1.40	6 (15%)
4	GOL	A	1111	-	5,5,5	0.30	0	5,5,5	0.38	0
4	GOL	A	1105	-	5,5,5	0.37	0	5,5,5	0.74	0
4	GOL	A	1109	-	5,5,5	0.38	0	5,5,5	0.55	0
4	GOL	A	1113	-	5,5,5	0.20	0	5,5,5	0.77	0
4	GOL	A	1115	-	5,5,5	0.31	0	5,5,5	0.75	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	1107	-	-	2/4/4/4	-
4	GOL	A	1112	-	-	2/4/4/4	-
4	GOL	A	1114	-	-	2/4/4/4	-
4	GOL	A	1110	-	-	4/4/4/4	-
4	GOL	A	1108	-	-	0/4/4/4	-
4	GOL	A	1106	-	-	2/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	89X	A	1116	2	-	3/31/31/31	0/2/2/2
4	GOL	A	1111	-	-	3/4/4/4	-
4	GOL	A	1105	-	-	0/4/4/4	-
4	GOL	A	1109	-	-	2/4/4/4	-
4	GOL	A	1113	-	-	1/4/4/4	-
4	GOL	A	1115	-	-	0/4/4/4	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1107	GOL	O2-C2	-3.71	1.32	1.43
5	A	1116	89X	CG-CB	-3.27	1.47	1.52
5	A	1116	89X	C9-N4	2.57	1.40	1.35
5	A	1116	89X	C3-N2	2.40	1.39	1.34
4	A	1106	GOL	O2-C2	-2.19	1.37	1.43

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1106	GOL	O2-C2-C3	-5.09	88.09	109.18
5	A	1116	89X	C3-C4-N3	3.75	121.25	111.11
5	A	1116	89X	O2-CA-CB	2.91	125.44	120.58
4	A	1107	GOL	O2-C2-C1	-2.88	97.24	109.18
5	A	1116	89X	O4-C9-N4	2.76	127.21	122.47
5	A	1116	89X	CZ-CE2-CD2	2.62	123.48	120.24
5	A	1116	89X	CB-CA-N	-2.53	110.38	116.84
5	A	1116	89X	CE2-CZ-CE1	-2.53	116.41	119.87
4	A	1114	GOL	O3-C3-C2	-2.00	101.36	110.38

There are no chirality outliers.

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1106	GOL	O1-C1-C2-C3
4	A	1107	GOL	O1-C1-C2-C3
4	A	1109	GOL	C1-C2-C3-O3
4	A	1110	GOL	O1-C1-C2-C3
4	A	1112	GOL	O1-C1-C2-C3
4	A	1114	GOL	C1-C2-C3-O3
4	A	1111	GOL	C1-C2-C3-O3

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	A	1107	GOL	O1-C1-C2-O2
4	A	1110	GOL	O1-C1-C2-O2
4	A	1112	GOL	O1-C1-C2-O2
4	A	1114	GOL	O2-C2-C3-O3
4	A	1106	GOL	O1-C1-C2-O2
4	A	1109	GOL	O2-C2-C3-O3
4	A	1110	GOL	O2-C2-C3-O3
4	A	1113	GOL	O2-C2-C3-O3
4	A	1111	GOL	O1-C1-C2-O2
5	A	1116	89X	N2-CB-CG-CD2
4	A	1110	GOL	C1-C2-C3-O3
4	A	1111	GOL	O2-C2-C3-O3
5	A	1116	89X	C4-C5-C6-C7
5	A	1116	89X	N2-CB-CG-CD1

There are no ring outliers.

7 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1107	GOL	2	0
4	A	1112	GOL	1	0
4	A	1114	GOL	1	0
4	A	1110	GOL	1	0
4	A	1106	GOL	2	0
4	A	1105	GOL	2	0
4	A	1109	GOL	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	890/914 (97%)	-0.29	2 (0%) 91 92	17, 29, 44, 73	8 (0%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	196	PRO	3.3
1	A	815	HIS	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	A	1113	6/6	0.80	0.18	40,44,45,51	6
4	GOL	A	1112	6/6	0.82	0.16	39,50,57,62	0
4	GOL	A	1111	6/6	0.83	0.14	58,60,68,71	0
4	GOL	A	1106	6/6	0.84	0.21	39,50,51,62	0
4	GOL	A	1109	6/6	0.84	0.12	48,61,64,77	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	A	1110	6/6	0.84	0.13	44,50,56,59	0
4	GOL	A	1108	6/6	0.85	0.13	52,56,59,61	0
4	GOL	A	1114	6/6	0.85	0.17	35,41,51,52	6
4	GOL	A	1105	6/6	0.87	0.12	42,47,52,53	0
4	GOL	A	1115	6/6	0.87	0.14	44,53,63,64	0
4	GOL	A	1107	6/6	0.90	0.20	31,37,44,46	0
5	89X	A	1116	30/30	0.92	0.10	27,35,55,63	0
3	MG	A	1104	1/1	0.93	0.05	65,65,65,65	0
3	MG	A	1103	1/1	0.95	0.12	46,46,46,46	0
3	MG	A	1102	1/1	0.98	0.02	33,33,33,33	0
2	ZN	A	1101	1/1	1.00	0.01	26,26,26,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.