



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 10, 2026 – 03:23 AM UTC

PDB ID : 1XN0 / pdb_00001xn0
Title : Catalytic Domain Of Human Phosphodiesterase 4B In Complex With (R,S)-Rolipram
Authors : Card, G.L.; England, B.P.; Suzuki, Y.; Fong, D.; Powell, B.; Lee, B.; Luu, C.; Tabrizizad, M.; Gillette, S.; Ibrahim, P.N.; Artis, D.R.; Bollag, G.; Milburn, M.V.; Kim, S.-H.; Schlessinger, J.; Zhang, K.Y.J.
Deposited on : 2004-10-04
Resolution : 2.31 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7754 (2.34-2.30)
Clashscore	190562	8383 (2.34-2.30)
Ramachandran outliers	187476	8303 (2.34-2.30)
Sidechain outliers	187428	8303 (2.34-2.30)
RSRZ outliers	180081	7760 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	398	% 66% 13% • 19%
1	B	398	% 68% 11% • 19%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	ROL	A	1003[A]	X	-	-	-
4	ROL	B	1003[A]	X	-	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5543 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cAMP-specific 3',5'-cyclic phosphodiesterase 4B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	323	Total	C	N	O	S	8	0	0
			2625	1657	441	505	22			
1	B	323	Total	C	N	O	S	4	0	0
			2625	1657	441	505	22			

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	131	MET	-	initiating methionine	UNP Q07343
A	132	GLY	-	cloning artifact	UNP Q07343
A	133	SER	-	cloning artifact	UNP Q07343
A	134	SER	-	cloning artifact	UNP Q07343
A	135	HIS	-	expression tag	UNP Q07343
A	136	HIS	-	expression tag	UNP Q07343
A	137	HIS	-	expression tag	UNP Q07343
A	138	HIS	-	expression tag	UNP Q07343
A	139	HIS	-	expression tag	UNP Q07343
A	140	HIS	-	expression tag	UNP Q07343
A	141	SER	-	cloning artifact	UNP Q07343
A	142	SER	-	cloning artifact	UNP Q07343
A	143	GLY	-	cloning artifact	UNP Q07343
A	144	LEU	-	cloning artifact	UNP Q07343
A	145	VAL	-	cloning artifact	UNP Q07343
A	146	PRO	-	cloning artifact	UNP Q07343
A	147	ARG	-	cloning artifact	UNP Q07343
A	148	GLY	-	cloning artifact	UNP Q07343
A	149	SER	-	cloning artifact	UNP Q07343
A	150	HIS	-	cloning artifact	UNP Q07343
A	151	MET	-	cloning artifact	UNP Q07343
B	131	MET	-	initiating methionine	UNP Q07343
B	132	GLY	-	cloning artifact	UNP Q07343
B	133	SER	-	cloning artifact	UNP Q07343
B	134	SER	-	cloning artifact	UNP Q07343

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	135	HIS	-	expression tag	UNP Q07343
B	136	HIS	-	expression tag	UNP Q07343
B	137	HIS	-	expression tag	UNP Q07343
B	138	HIS	-	expression tag	UNP Q07343
B	139	HIS	-	expression tag	UNP Q07343
B	140	HIS	-	expression tag	UNP Q07343
B	141	SER	-	cloning artifact	UNP Q07343
B	142	SER	-	cloning artifact	UNP Q07343
B	143	GLY	-	cloning artifact	UNP Q07343
B	144	LEU	-	cloning artifact	UNP Q07343
B	145	VAL	-	cloning artifact	UNP Q07343
B	146	PRO	-	cloning artifact	UNP Q07343
B	147	ARG	-	cloning artifact	UNP Q07343
B	148	GLY	-	cloning artifact	UNP Q07343
B	149	SER	-	cloning artifact	UNP Q07343
B	150	HIS	-	cloning artifact	UNP Q07343
B	151	MET	-	cloning artifact	UNP Q07343


- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		
3	B	1	Total	Mg	0	0
			1	1		

- Molecule 4 is ROLIPRAM (CCD ID: ROL) (formula: C₁₆H₂₁NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	1
			20	16	1	3		
4	A	1	Total	C	N	O	0	1
			20	16	1	3		
4	B	1	Total	C	N	O	0	1
			20	16	1	3		
4	B	1	Total	C	N	O	0	1
			20	16	1	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	105	Total	O	0	0
			105	105		
5	B	104	Total	O	0	0
			104	104		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	89.93Å 95.00Å 106.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	70.71 – 2.31 70.71 – 2.31	Depositor EDS
% Data completeness (in resolution range)	98.6 (70.71-2.31) 98.6 (70.71-2.31)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.88 (at 2.31Å)	Xtriage
Refinement program	REFMAC 5.1.25	Depositor
R, R_{free}	0.205 , 0.250 (Not available) , 0.286	Depositor DCC
R_{free} test set	2013 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	35.0	Xtriage
Anisotropy	0.367	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 42.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5543	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 43.55 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.7340e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG, ROL, CME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.02	6/2647 (0.2%)	1.09	9/3586 (0.3%)
1	B	0.96	3/2647 (0.1%)	1.04	5/3586 (0.1%)
All	All	0.99	9/5294 (0.2%)	1.06	14/7172 (0.2%)

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	411	MET	SD-CE	-12.71	1.47	1.79
1	A	411	MET	SD-CE	-12.54	1.48	1.79
1	A	328	LYS	CB-CG	-11.89	1.16	1.52
1	A	387	MET	SD-CE	-8.09	1.59	1.79
1	A	426	MET	SD-CE	-7.71	1.60	1.79

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	328	LYS	CA-CB-CG	11.65	137.40	114.10
1	A	164	ASP	N-CA-C	-6.74	103.74	112.23
1	B	326	LEU	N-CA-C	-6.49	101.78	110.55
1	A	318	GLU	CB-CG-CD	-6.47	101.61	112.60
1	A	318	GLU	CA-CB-CG	-6.38	101.35	114.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2625	0	2550	26	0
1	B	2625	0	2550	26	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	40	0	42	1	0
4	B	40	0	42	5	0
5	A	105	0	0	0	0
5	B	104	0	0	4	0
All	All	5543	0	5184	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 53 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:421:GLU:HA	1:A:426:MET:HE3	1.62	0.82
1:B:289:THR:HG22	1:B:426:MET:HE1	1.70	0.72
1:B:252:THR:HG21	5:B:2039:HOH:O	1.90	0.71
1:A:296:MET:SD	1:B:296:MET:SD	2.91	0.69
1:A:250:LEU:HG	1:A:387:MET:HE1	1.76	0.66

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	318/398 (80%)	307 (96%)	9 (3%)	2 (1%)	21 25

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	318/398 (80%)	307 (96%)	11 (4%)	0	100	100
All	All	636/796 (80%)	614 (96%)	20 (3%)	2 (0%)	36	45

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	317	GLU
1	A	319	HIS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	290/357 (81%)	275 (95%)	15 (5%)	21	30
1	B	290/357 (81%)	278 (96%)	12 (4%)	27	40
All	All	580/714 (81%)	553 (95%)	27 (5%)	23	34

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	469	ILE
1	B	204	ASP
1	B	371	VAL
1	B	189	ASN
1	B	207	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	481	GLN
1	B	416	GLN
1	B	376	ASN
1	B	332	GLN
1	B	395	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	CME	B	320	1	8,9,10	1.74	2 (25%)	6,9,11	2.10	2 (33%)
1	CME	A	432	1	8,9,10	1.69	1 (12%)	6,9,11	1.48	2 (33%)
1	CME	A	320	1	8,9,10	1.59	1 (12%)	6,9,11	2.11	1 (16%)
1	CME	B	194	1	8,9,10	1.69	2 (25%)	6,9,11	1.88	2 (33%)
1	CME	B	432	1	8,9,10	1.63	1 (12%)	6,9,11	2.83	3 (50%)
1	CME	A	194	1	8,9,10	1.53	1 (12%)	6,9,11	2.01	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CME	B	320	1	-	3/5/8/10	-
1	CME	A	432	1	-	3/5/8/10	-
1	CME	A	320	1	-	2/5/8/10	-
1	CME	B	194	1	-	0/5/8/10	-
1	CME	B	432	1	-	3/5/8/10	-
1	CME	A	194	1	-	2/5/8/10	-

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	432	CME	OH-CZ	-4.22	1.20	1.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	432	CME	OH-CZ	-3.97	1.21	1.42
1	A	320	CME	OH-CZ	-3.88	1.22	1.42
1	B	194	CME	OH-CZ	-3.73	1.23	1.42
1	A	194	CME	OH-CZ	-3.71	1.23	1.42

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	432	CME	CB-SG-SD	5.15	117.18	103.86
1	A	320	CME	CB-SG-SD	4.12	114.51	103.86
1	B	320	CME	CB-SG-SD	3.96	114.10	103.86
1	A	194	CME	CB-SG-SD	3.69	113.41	103.86
1	B	432	CME	CB-CA-C	3.27	119.68	110.80

There are no chirality outliers.

5 of 13 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	194	CME	CE-SD-SG-CB
1	B	320	CME	CE-SD-SG-CB
1	B	432	CME	SD-CE-CZ-OH
1	A	320	CME	CE-SD-SG-CB
1	B	432	CME	CE-SD-SG-CB

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	320	CME	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	ROL	B	1003[A]	-	22,22,22	1.32	3 (13%)	30,30,30	1.64	4 (13%)
4	ROL	A	1004[B]	-	22,22,22	1.00	1 (4%)	30,30,30	1.75	5 (16%)
4	ROL	A	1003[A]	-	22,22,22	1.24	2 (9%)	30,30,30	1.94	6 (20%)
4	ROL	B	1004[B]	-	22,22,22	1.21	2 (9%)	30,30,30	1.67	5 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ROL	B	1003[A]	-	1/1/3/4	2/10/26/26	0/3/3/3
4	ROL	A	1004[B]	-	-	1/10/26/26	0/3/3/3
4	ROL	A	1003[A]	-	1/1/3/4	4/10/26/26	0/3/3/3
4	ROL	B	1004[B]	-	-	2/10/26/26	0/3/3/3

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1003[A]	ROL	C2-C3	-3.56	1.48	1.54
4	B	1003[A]	ROL	C2-C3	-3.09	1.49	1.54
4	B	1004[B]	ROL	O3-C7	2.80	1.42	1.37
4	B	1004[B]	ROL	C12-C11	-2.53	1.39	1.52
4	A	1004[B]	ROL	C12-C11	-2.49	1.39	1.52

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1003[A]	ROL	O3-C11-C12	-5.26	90.61	108.71
4	A	1003[A]	ROL	C16-O2-C8	5.09	124.97	117.51
4	A	1004[B]	ROL	O3-C11-C12	-4.89	91.88	108.71
4	B	1004[B]	ROL	C16-O2-C8	4.85	124.63	117.51
4	B	1003[A]	ROL	O3-C11-C12	-4.78	92.25	108.71

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	A	1003[A]	ROL	C3
4	B	1003[A]	ROL	C3

5 of 9 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1003[A]	ROL	C15-C11-O3-C7
4	A	1003[A]	ROL	C12-C11-O3-C7
4	B	1004[B]	ROL	C12-C11-O3-C7
4	B	1004[B]	ROL	C15-C11-O3-C7
4	B	1003[A]	ROL	C7-C8-O2-C16


There are no ring outliers.

3 monomers are involved in 6 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	1003[A]	ROL	2	0
4	A	1003[A]	ROL	1	0
4	B	1004[B]	ROL	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand ROL B 1003 (A)



Ligand ROL A 1004 (B)





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	320/398 (80%)	0.08	5 (1%) 70 72	11, 19, 34, 57	2 (0%)
1	B	320/398 (80%)	-0.10	4 (1%) 75 76	12, 19, 34, 50	1 (0%)
All	All	640/796 (80%)	-0.01	9 (1%) 73 75	11, 19, 34, 57	3 (0%)

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	368	SER	2.5
1	A	319	HIS	2.4
1	A	318	GLU	2.2
1	B	430	PRO	2.2
1	B	485	PRO	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CME	B	320	10/11	0.89	0.14	27,28,38,39	0
1	CME	A	432	10/11	0.91	0.12	24,25,36,38	0
1	CME	B	432	10/11	0.92	0.12	19,22,42,43	0
1	CME	A	320	10/11	0.93	0.10	33,35,41,41	0
1	CME	A	194	10/11	0.95	0.08	14,15,26,29	0
1	CME	B	194	10/11	0.97	0.06	15,16,25,29	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	ROL	A	1003[A]	20/20	0.93	0.08	30,34,35,35	20
4	ROL	A	1004[B]	20/20	0.94	0.08	26,29,33,33	20
4	ROL	B	1004[B]	20/20	0.94	0.07	25,26,29,29	20
4	ROL	B	1003[A]	20/20	0.95	0.08	34,35,37,37	20
2	ZN	B	1001	1/1	0.99	0.02	28,28,28,28	0
3	MG	A	1002	1/1	1.00	0.01	12,12,12,12	0
3	MG	B	1002	1/1	1.00	0.02	17,17,17,17	0
2	ZN	A	1001	1/1	1.00	0.02	26,26,26,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ROL A 1003 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ROL A 1004 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ROL B 1004 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ROL B 1003 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.