



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 01:16 AM UTC

PDB ID : 4XN5 / pdb_00004xn5
Title : Crystal Structure of Met260Ala mutant of E. coli Aminopeptidase N in complex with L-Phenylalanine
Authors : Addlagatta, A.; Gumpena, R.; Kishor, C.
Deposited on : 2015-01-15
Resolution : 2.66 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.66 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1110 (2.66-2.66)
Clashscore	190562	1141 (2.66-2.66)
Ramachandran outliers	187476	1126 (2.66-2.66)
Sidechain outliers	187428	1126 (2.66-2.66)
RSRZ outliers	180081	1110 (2.66-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	866	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	A	906	-	X	X	-

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 7599 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aminopeptidase N.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	866	Total	C	N	O	S	0	7	0
			6991	4427	1219	1318	27			


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	260	ALA	MET	engineered mutation	UNP P04825

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is PHENYLALANINE (CCD ID: PHE) (formula: C₉H₁₁NO₂).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			12	9	1	2		

- Molecule 4 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	3	Total	Na	0	0
			3	3		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is MALONATE ION (CCD ID: MLI) (formula: C₃H₂O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			7	3	4		
6	A	1	Total	C	O	0	0
			7	3	4		
6	A	1	Total	C	O	0	0
			7	3	4		
6	A	1	Total	C	O	0	0
			7	3	4		
6	A	1	Total	C	O	0	0
			7	3	4		
6	A	1	Total	C	O	0	0
			7	3	4		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	519	Total	O	0	0
			519	519		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Aminopeptidase N



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	121.03Å 121.03Å 170.81Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	17.97 – 2.66 17.97 – 2.66	Depositor EDS
% Data completeness (in resolution range)	96.6 (17.97-2.66) 96.6 (17.97-2.66)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.17	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.35 (at 2.66Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.140 , 0.202 0.149 , 0.206	Depositor DCC
R_{free} test set	2067 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	22.3	Xtriage
Anisotropy	0.127	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 51.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.021 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7599	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN, MLI, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.02	1/7167 (0.0%)	1.04	8/9728 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	248	ASP	CG-OD2	5.19	1.35	1.25

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	78	GLU	N-CA-C	-14.50	79.92	110.80
1	A	64	VAL	CB-CA-C	-7.20	102.21	110.41
1	A	501	LYS	N-CA-C	-6.25	99.81	109.24
1	A	77	GLU	N-CA-C	-6.24	97.51	110.80
1	A	559	ARG	N-CA-C	5.13	116.95	111.36
1	A	781	ARG	NE-CZ-NH1	-5.10	116.40	121.50
1	A	279	ARG	N-CA-C	-5.07	101.03	108.99
1	A	838	ALA	N-CA-C	5.01	117.39	111.33

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	77	GLU	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6991	0	6857	57	0
2	A	1	0	0	0	0
3	A	12	0	8	1	0
4	A	3	0	0	0	0
5	A	24	0	32	5	0
6	A	49	0	14	4	0
7	A	519	0	0	12	1
All	All	7599	0	6911	60	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:781:ARG:HH11	1:A:781:ARG:HG3	0.99	1.15
1:A:781:ARG:HH11	1:A:781:ARG:CG	1.80	0.92
1:A:781:ARG:HG3	1:A:781:ARG:NH1	1.82	0.90
1:A:222:ARG:HD2	7:A:1211:HOH:O	1.78	0.83
1:A:346:ARG:NH1	1:A:615:GLU:OE1	2.17	0.77
1:A:204[B]:ARG:HD3	7:A:1379:HOH:O	1.88	0.74
1:A:78:GLU:HA	1:A:78:GLU:OE1	1.91	0.69
1:A:741:GLN:HB3	7:A:1474:HOH:O	1.93	0.67
1:A:423:GLN:HG3	7:A:1181:HOH:O	1.93	0.67
1:A:246:ASP:HB2	5:A:906:GOL:H31	1.79	0.64
1:A:845[B]:ARG:NH2	1:A:869:LEU:O	2.31	0.63
1:A:468:ARG:HD3	7:A:1164:HOH:O	2.00	0.61
1:A:222:ARG:CD	7:A:1211:HOH:O	2.44	0.61
1:A:6:GLN:NE2	7:A:1001:HOH:O	2.36	0.58
1:A:747:ASP:CG	1:A:781:ARG:HH12	2.14	0.56
1:A:30:ASP:HB3	1:A:33:LYS:O	2.07	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:246:ASP:CB	5:A:906:GOL:H31	2.36	0.54
1:A:810[A]:GLU:HG2	1:A:811:MET:HE2	1.90	0.52
1:A:390:MET:HE1	1:A:439:TRP:CE2	2.45	0.52
1:A:671:GLU:HG2	7:A:1410:HOH:O	2.12	0.50
1:A:265:ASN:HB2	1:A:268:LEU:O	2.12	0.49
6:A:910:MLI:O7	6:A:911:MLI:O7	2.29	0.49
1:A:727:CYS:O	1:A:731:LEU:HG	2.12	0.49
1:A:741:GLN:CD	1:A:741:GLN:H	2.20	0.49
1:A:224:PRO:HB2	6:A:914:MLI:O8	2.12	0.48
1:A:766:ARG:NH1	7:A:1006:HOH:O	2.46	0.48
1:A:286:LYS:HD2	6:A:916:MLI:H12	1.95	0.48
1:A:747:ASP:OD1	1:A:781:ARG:NH1	2.40	0.47
1:A:566:ASP:OD2	5:A:907:GOL:H2	2.14	0.47
6:A:913:MLI:O9	6:A:913:MLI:O7	2.31	0.47
1:A:300:PHE:HB3	1:A:326:ARG:NH2	2.30	0.47
1:A:500:GLN:NE2	1:A:523:ASN:HD22	2.12	0.47
1:A:108:GLY:O	1:A:118:THR:HA	2.15	0.46
1:A:307:ARG:HH22	5:A:906:GOL:H12	1.80	0.46
1:A:810[A]:GLU:HG2	1:A:811:MET:CE	2.45	0.46
3:A:902:PHE:HB3	7:A:1414:HOH:O	2.15	0.46
1:A:502:GLY:N	1:A:522:ASP:OD2	2.49	0.46
1:A:145:ILE:O	1:A:173:TRP:HA	2.16	0.46
1:A:593:VAL:HG23	7:A:1294:HOH:O	2.14	0.46
1:A:300:PHE:HB3	1:A:326:ARG:CZ	2.47	0.45
1:A:316:LEU:HD23	1:A:380:VAL:HG11	1.98	0.45
1:A:189:LEU:C	1:A:189:LEU:HD23	2.42	0.45
1:A:274:LYS:HE2	1:A:275:TYR:CZ	2.52	0.44
1:A:195:ASP:HB3	1:A:220:LEU:HD13	2.00	0.44
1:A:390:MET:HE1	1:A:439:TRP:CD2	2.54	0.43
1:A:579:ASN:OD1	1:A:589:LEU:HA	2.19	0.43
1:A:234:MET:HG2	1:A:299:TYR:CZ	2.54	0.42
1:A:236:TRP:CZ2	1:A:240:ARG:HG2	2.54	0.42
1:A:307:ARG:HH22	5:A:906:GOL:C1	2.32	0.42
1:A:776:MET:O	1:A:782:ILE:HD11	2.21	0.41
1:A:53[A]:ARG:NH1	1:A:53[A]:ARG:HG3	2.35	0.41
1:A:140:ARG:HD3	7:A:1060:HOH:O	2.20	0.41
1:A:189:LEU:HD23	1:A:190:VAL:N	2.36	0.41
1:A:831:ILE:HB	1:A:868:ALA:HB2	2.03	0.41
1:A:223:ALA:N	1:A:224:PRO:CD	2.84	0.41
1:A:378:LEU:O	1:A:382:GLU:HB2	2.20	0.41
1:A:106:LEU:HD22	1:A:110:TYR:CD2	2.56	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:312:ASP:OD1	1:A:314:PHE:HB2	2.21	0.40
1:A:501:LYS:CE	1:A:520:VAL:O	2.68	0.40
1:A:234:MET:HG2	1:A:299:TYR:CE1	2.57	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:1064:HOH:O	7:A:1245:HOH:O[5_554]	2.11	0.09

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	871/866 (101%)	845 (97%)	25 (3%)	1 (0%)	48 66

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	502	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	747/740 (101%)	723 (97%)	24 (3%)	34	55

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	48	SER
1	A	78	GLU
1	A	93	LYS
1	A	204[A]	ARG
1	A	204[B]	ARG
1	A	248	ASP
1	A	319	LYS
1	A	356	ASP
1	A	398	GLU
1	A	423	GLN
1	A	441	SER
1	A	451	LYS
1	A	500	GLN
1	A	505	PRO
1	A	517	GLN
1	A	572	LEU
1	A	639	GLU
1	A	775	THR
1	A	781	ARG
1	A	800	GLU
1	A	810[A]	GLU
1	A	810[B]	GLU
1	A	850	GLN
1	A	860	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	41	GLN
1	A	45	HIS
1	A	55	ASN
1	A	164	GLN
1	A	500	GLN
1	A	514	GLN
1	A	733	GLN
1	A	741	GLN
1	A	821	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 16 ligands modelled in this entry, 4 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	GOL	A	909	-	5,5,5	1.21	0	5,5,5	0.90	0
5	GOL	A	907	-	5,5,5	0.45	0	5,5,5	1.94	1 (20%)
3	PHE	A	902	2	11,12,12	0.91	1 (9%)	11,15,15	0.83	0
6	MLI	A	910	-	6,6,6	1.25	0	7,7,7	1.28	0
6	MLI	A	916	-	6,6,6	1.61	1 (16%)	7,7,7	0.91	0
6	MLI	A	911	-	6,6,6	1.21	0	7,7,7	2.49	3 (42%)
6	MLI	A	915	-	6,6,6	2.07	2 (33%)	7,7,7	1.71	2 (28%)
6	MLI	A	914	-	6,6,6	1.10	0	7,7,7	2.08	2 (28%)
5	GOL	A	906	-	5,5,5	1.86	1 (20%)	5,5,5	2.96	4 (80%)
6	MLI	A	912	-	6,6,6	1.77	2 (33%)	7,7,7	1.82	2 (28%)
5	GOL	A	908	-	5,5,5	0.94	0	5,5,5	1.09	0
6	MLI	A	913	-	6,6,6	1.31	1 (16%)	7,7,7	0.88	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	909	-	-	0/4/4/4	-
5	GOL	A	907	-	-	2/4/4/4	-
3	PHE	A	902	2	-	2/8/8/8	0/1/1/1
6	MLI	A	910	-	-	2/4/4/4	-
6	MLI	A	916	-	-	2/4/4/4	-
6	MLI	A	911	-	-	0/4/4/4	-
6	MLI	A	915	-	-	0/4/4/4	-
6	MLI	A	914	-	-	2/4/4/4	-
5	GOL	A	906	-	-	2/4/4/4	-
6	MLI	A	912	-	-	2/4/4/4	-
5	GOL	A	908	-	-	2/4/4/4	-
6	MLI	A	913	-	-	2/4/4/4	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	915	MLI	C1-C2	3.22	1.56	1.51
6	A	915	MLI	C1-C3	2.89	1.55	1.51
5	A	906	GOL	O2-C2	2.86	1.51	1.43
6	A	912	MLI	C1-C3	2.46	1.54	1.51
6	A	912	MLI	O8-C3	2.45	1.30	1.22
6	A	913	MLI	O9-C3	-2.28	1.23	1.30
3	A	902	PHE	OXT-C	-2.27	1.23	1.30
6	A	916	MLI	O6-C2	2.26	1.29	1.22

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	906	GOL	O3-C3-C2	4.25	129.50	110.38
6	A	911	MLI	O6-C2-C1	-4.09	110.45	122.11
6	A	914	MLI	C3-C1-C2	-4.04	98.70	112.95
5	A	906	GOL	O2-C2-C3	3.73	124.64	109.18
6	A	911	MLI	C3-C1-C2	-3.58	100.30	112.95
6	A	912	MLI	O6-C2-C1	-3.35	112.56	122.11
5	A	907	GOL	O3-C3-C2	-3.24	95.77	110.38
6	A	911	MLI	O7-C2-C1	3.03	123.89	114.51
6	A	914	MLI	O9-C3-C1	2.81	123.22	114.51
5	A	906	GOL	C3-C2-C1	-2.70	101.89	111.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	915	MLI	C3-C1-C2	2.69	122.43	112.95
6	A	915	MLI	O7-C2-C1	2.57	122.46	114.51
5	A	906	GOL	O1-C1-C2	2.11	119.90	110.38
6	A	912	MLI	O7-C2-C1	2.05	120.87	114.51

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	906	GOL	O1-C1-C2-C3
5	A	908	GOL	C1-C2-C3-O3
5	A	907	GOL	C1-C2-C3-O3
5	A	908	GOL	O2-C2-C3-O3
5	A	907	GOL	O2-C2-C3-O3
6	A	912	MLI	C3-C1-C2-O7
6	A	916	MLI	C2-C1-C3-O9
5	A	906	GOL	O1-C1-C2-O2
6	A	912	MLI	C3-C1-C2-O6
6	A	916	MLI	C2-C1-C3-O8
6	A	914	MLI	C3-C1-C2-O6
3	A	902	PHE	OXT-C-CA-N
6	A	913	MLI	C2-C1-C3-O8
6	A	914	MLI	C3-C1-C2-O7
6	A	910	MLI	C2-C1-C3-O9
6	A	910	MLI	C2-C1-C3-O8
6	A	913	MLI	C2-C1-C3-O9
3	A	902	PHE	O-C-CA-N

There are no ring outliers.

8 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	907	GOL	1	0
3	A	902	PHE	1	0
6	A	910	MLI	1	0
6	A	916	MLI	1	0
6	A	911	MLI	1	0
6	A	914	MLI	1	0
5	A	906	GOL	4	0
6	A	913	MLI	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	866/866 (100%)	-0.90	0 100 100	8, 19, 36, 67	7 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	MLI	A	916	7/7	0.67	0.25	46,64,78,81	0
5	GOL	A	909	6/6	0.80	0.19	43,43,53,54	0
6	MLI	A	914	7/7	0.86	0.13	28,29,33,35	7
6	MLI	A	912	7/7	0.86	0.14	39,44,49,53	0
5	GOL	A	906	6/6	0.91	0.09	22,23,24,26	0
6	MLI	A	915	7/7	0.92	0.09	44,45,54,55	0
5	GOL	A	908	6/6	0.92	0.10	38,47,47,48	0
6	MLI	A	910	7/7	0.93	0.09	31,34,38,46	0
5	GOL	A	907	6/6	0.94	0.09	29,33,35,37	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	PHE	A	902	12/12	0.95	0.12	25,36,46,47	0
4	NA	A	903	1/1	0.95	0.05	33,33,33,33	0
6	MLI	A	913	7/7	0.95	0.10	46,52,58,59	0
6	MLI	A	911	7/7	0.97	0.06	32,41,44,46	0
4	NA	A	904	1/1	0.97	0.03	28,28,28,28	0
4	NA	A	905	1/1	0.99	0.07	21,21,21,21	0
2	ZN	A	901	1/1	1.00	0.01	15,15,15,15	0

6.5 Other polymers [i](#)

There are no such residues in this entry.