



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 01:01 PM UTC

PDB ID : 7XNF / pdb_00007xnf
Title : Structure of SARS-CoV-2 antibody P2C-1F11 with GX/P2V/2017 RBD
Authors : Jia, Y.F.; Chai, Y.; Wang, Q.H.; Gao, G.F.
Deposited on : 2022-04-28
Resolution : 2.79 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	227	15% 76% 18% 6%
2	L	215	22% 87% 12%
3	A	215	47% 50% 21% 29%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4465 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called P2C-1F11 Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	H	213	Total	C	N	O	S	0	0	0
			1578	990	268	313	7			

- Molecule 2 is a protein called P2C-1F11 Lambda chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	214	Total	C	N	O	S	0	0	0
			1636	1021	277	333	5			

- Molecule 3 is a protein called Spike protein S1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	153	Total	C	N	O	S	0	0	0
			1232	800	203	223	6			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	528	HIS	-	expression tag	UNP A0A6G6A2Q2
A	529	HIS	-	expression tag	UNP A0A6G6A2Q2
A	530	HIS	-	expression tag	UNP A0A6G6A2Q2
A	531	HIS	-	expression tag	UNP A0A6G6A2Q2
A	532	HIS	-	expression tag	UNP A0A6G6A2Q2
A	533	HIS	-	expression tag	UNP A0A6G6A2Q2

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	H	8	Total	O	0	0
			8	8		
4	L	7	Total	O	0	0
			7	7		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	4	Total	O	0	0
			4	4		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: P2C-1F11 Heavy Chain



• Molecule 2: P2C-1F11 Lambda chain



• Molecule 3: Spike protein S1



Sequence logo for the 5' region of the 16S rRNA gene. The y-axis represents information content in bits (0.00 to 0.25). The x-axis shows positions 1 to 10. The sequence P521 A522 T523 V524 is highlighted in green. The background is gray.

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	55.72Å 85.44Å 183.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.61 – 2.79 41.61 – 2.79	Depositor EDS
% Data completeness (in resolution range)	95.1 (41.61-2.79) 95.1 (41.61-2.79)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.15 (at 2.81Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.320 , 0.380 0.318 , 0.376	Depositor DCC
R_{free} test set	1063 reflections (4.71%)	wwPDB-VP
Wilson B-factor (Å ²)	48.0	Xtriage
Anisotropy	0.141	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 22.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.82	EDS
Total number of atoms	4465	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	H	0.08	0/1612	0.23	0/2198
2	L	0.07	0/1671	0.23	0/2268
3	A	0.22	0/1263	0.40	0/1707
All	All	0.13	0/4546	0.29	0/6173

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1578	0	1549	25	0
2	L	1636	0	1589	13	0
3	A	1232	0	1187	28	0
4	A	4	0	0	2	0
4	H	8	0	0	0	0
4	L	7	0	0	1	0
All	All	4465	0	4325	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:82:MET:HB3	1:H:85:LEU:HD21	1.65	0.79
1:H:35:ASN:ND2	1:H:98:ASP:OD2	2.25	0.69
1:H:12:VAL:HG11	1:H:85:LEU:HD12	1.76	0.68
3:A:481:ASN:ND2	4:A:601:HOH:O	2.26	0.66
2:L:67:GLY:HA3	2:L:72:PHE:HA	1.78	0.66
3:A:426:PRO:HG3	3:A:464:PHE:HE2	1.60	0.66
2:L:78:ARG:NH2	4:L:302:HOH:O	2.28	0.65
2:L:145:LYS:HB3	2:L:197:THR:HB	1.81	0.63
3:A:461:LEU:HD22	3:A:465:GLU:HB3	1.82	0.62
2:L:38:GLN:HB2	2:L:48:LEU:HD11	1.86	0.57
1:H:214:ARG:NH1	1:H:216:GLU:OE1	2.37	0.57
2:L:61:ASP:N	2:L:61:ASP:OD1	2.39	0.56
3:A:393:THR:HA	3:A:522:ALA:HA	1.88	0.55
1:H:90:THR:HG23	1:H:114:THR:HA	1.89	0.54
2:L:6:GLN:H	2:L:100:GLN:HE22	1.54	0.54
1:H:36:TRP:HD1	1:H:69:ILE:HD12	1.72	0.54
3:A:392:PHE:HD1	3:A:517:LEU:HG	1.73	0.53
1:H:127:PRO:HD3	1:H:213:LYS:HE3	1.91	0.53
1:H:40:ALA:HB3	1:H:43:LYS:HB2	1.90	0.53
3:A:409:GLN:HE22	3:A:416:GLY:HA3	1.73	0.53
1:H:22:CYS:HB3	1:H:78:LEU:HB3	1.90	0.53
1:H:126:PHE:HD2	1:H:145:LEU:HD23	1.73	0.53
2:L:90:GLN:HE21	2:L:96:PRO:HB3	1.74	0.52
2:L:4:LEU:HD21	2:L:91:GLN:HB2	1.92	0.52
3:A:358:ILE:HB	3:A:395:VAL:HG13	1.92	0.52
1:H:62:SER:O	1:H:66:ARG:NH2	2.36	0.52
3:A:378:LYS:NZ	3:A:379:CYS:O	2.45	0.50
1:H:172:ALA:HB2	1:H:182:LEU:HD23	1.92	0.50
3:A:426:PRO:HG3	3:A:464:PHE:CE2	2.44	0.50
3:A:350:VAL:HG22	3:A:422:ASN:HB3	1.93	0.49
1:H:11:LEU:HD12	1:H:114:THR:HB	1.94	0.49
3:A:340:GLU:OE1	3:A:356:LYS:NZ	2.44	0.48
2:L:1:GLU:O	2:L:27:GLN:NE2	2.36	0.48
3:A:439:VAL:HB	3:A:507:PRO:HG2	1.94	0.48
2:L:13:LEU:O	2:L:107:LYS:N	2.41	0.47
3:A:409:GLN:O	3:A:414:GLN:NE2	2.47	0.47
3:A:453:TYR:CE1	3:A:493:GLU:HB2	2.49	0.47
1:H:4:LEU:HD21	1:H:27:ILE:HD11	1.96	0.47
3:A:490:TYR:HD1	3:A:492:LEU:H	1.61	0.47
1:H:35:ASN:OD1	1:H:47:TRP:NE1	2.41	0.46
1:H:36:TRP:CD1	1:H:69:ILE:HD12	2.49	0.46
3:A:418:ILE:HA	3:A:422:ASN:HB2	1.96	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:11:LEU:HD21	1:H:150:PHE:HE2	1.81	0.46
3:A:405:ASP:OD1	3:A:504:ASN:HB3	2.15	0.46
2:L:29:VAL:HG21	2:L:91:GLN:HG2	1.98	0.45
3:A:374:PHE:HD1	3:A:436:TRP:HB3	1.81	0.45
3:A:439:VAL:HG22	3:A:509:ARG:HD2	1.97	0.45
3:A:381:GLY:HA3	3:A:430:THR:HA	1.97	0.45
1:H:172:ALA:HA	1:H:182:LEU:HB3	1.99	0.45
1:H:38:ARG:NE	1:H:46:GLU:OE1	2.36	0.45
3:A:489:TYR:HA	4:A:604:HOH:O	2.18	0.43
2:L:21:LEU:HB2	2:L:74:LEU:HB3	2.00	0.43
1:H:38:ARG:NH1	1:H:93:TYR:OH	2.46	0.42
3:A:408:ARG:HD2	3:A:408:ARG:H	1.84	0.42
3:A:420:ASP:HB3	3:A:460:LYS:HG2	2.01	0.42
1:H:32:ASN:HB2	1:H:34:MET:HE3	2.01	0.41
1:H:72:ASP:OD1	1:H:74:SER:OG	2.35	0.41
1:H:68:THR:O	1:H:81:GLN:N	2.48	0.41
3:A:336:CYS:HA	3:A:337:PRO:HD3	1.85	0.41
1:H:99:LEU:O	1:H:103:GLY:N	2.54	0.41
3:A:420:ASP:O	3:A:461:LEU:N	2.39	0.41
2:L:158:ASN:N	2:L:158:ASN:OD1	2.53	0.41
3:A:394:ASN:HB2	3:A:516:GLU:HB3	2.03	0.40
3:A:436:TRP:CZ2	3:A:509:ARG:HD3	2.57	0.40
1:H:33:TYR:HB2	1:H:98:ASP:HB3	2.04	0.40
3:A:426:PRO:C	3:A:428:ASP:H	2.29	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	H	209/227 (92%)	203 (97%)	6 (3%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	L	212/215 (99%)	207 (98%)	5 (2%)	0	100	100
3	A	141/215 (66%)	134 (95%)	7 (5%)	0	100	100
All	All	562/657 (86%)	544 (97%)	18 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	178/192 (93%)	177 (99%)	1 (1%)	78	92
2	L	185/186 (100%)	181 (98%)	4 (2%)	45	78
3	A	134/188 (71%)	132 (98%)	2 (2%)	57	84
All	All	497/566 (88%)	490 (99%)	7 (1%)	59	85

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	2	VAL
2	L	30	SER
2	L	73	THR
2	L	94	SER
2	L	95	SER
3	A	361	CYS
3	A	518	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
2	L	91	GLN
2	L	100	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	H	213/227 (93%)	1.29	35 (16%) 4 3	20, 38, 57, 63	0
2	L	214/215 (99%)	1.28	48 (22%) 2 1	17, 37, 57, 74	0
3	A	153/215 (71%)	2.60	101 (66%) 0 0	24, 58, 82, 97	0
All	All	580/657 (88%)	1.63	184 (31%) 1 1	17, 41, 72, 97	0

All (184) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	A	434	ILE	7.2
2	L	25	ALA	6.2
3	A	451	TYR	6.0
3	A	441	GLN	5.9
3	A	414	GLN	5.8
3	A	508	PHE	5.7
3	A	432	CYS	5.7
3	A	357	ARG	5.6
3	A	361	CYS	5.5
3	A	505	TYR	5.4
3	A	524	VAL	5.4
3	A	395	VAL	5.1
2	L	4	LEU	5.1
2	L	67	GLY	5.0
3	A	377	PHE	5.0
3	A	380	TYR	5.0
3	A	401	VAL	4.8
3	A	499	PRO	4.8
3	A	376	THR	4.8
3	A	503	VAL	4.7
2	L	93	GLY	4.7
3	A	396	TYR	4.7
3	A	374	PHE	4.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	A	360	ASN	4.4
3	A	510	VAL	4.3
3	A	521	PRO	4.2
2	L	33	TYR	4.2
3	A	392	PHE	4.2
3	A	512	VAL	4.2
3	A	507	PRO	4.1
2	L	31	SER	4.1
3	A	430	THR	4.0
3	A	381	GLY	4.0
2	L	2	ILE	3.9
3	A	358	ILE	3.9
3	A	513	LEU	3.9
3	A	340	GLU	3.9
3	A	440	LYS	3.9
3	A	436	TRP	3.8
3	A	437	ASN	3.8
2	L	30	SER	3.8
2	L	28	SER	3.8
3	A	348	ALA	3.8
3	A	402	VAL	3.7
3	A	407	VAL	3.7
2	L	3	VAL	3.6
3	A	379	CYS	3.6
3	A	506	GLN	3.6
2	L	22	SER	3.6
2	L	152	ASN	3.5
1	H	12	VAL	3.5
3	A	514	SER	3.5
2	L	8	PRO	3.5
3	A	406	GLU	3.5
1	H	115	VAL	3.5
3	A	518	LEU	3.5
3	A	337	PRO	3.5
3	A	438	SER	3.5
3	A	349	SER	3.4
3	A	427	ASP	3.4
1	H	61	ASP	3.4
3	A	515	PHE	3.4
3	A	378	LYS	3.4
2	L	24	ARG	3.4
3	A	409	GLN	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	A	509	ARG	3.3
3	A	394	ASN	3.3
3	A	341	VAL	3.3
3	A	399	SER	3.3
3	A	495	TYR	3.3
1	H	60	ALA	3.3
3	A	356	LYS	3.3
3	A	342	PHE	3.3
3	A	425	LEU	3.3
3	A	439	VAL	3.3
1	H	16	GLY	3.2
2	L	23	CYS	3.2
2	L	96	PRO	3.2
1	H	63	VAL	3.2
1	H	113	VAL	3.2
1	H	65	GLY	3.2
3	A	345	SER	3.2
3	A	338	PHE	3.2
1	H	67	PHE	3.1
3	A	517	LEU	3.1
1	H	137	GLY	3.1
1	H	17	SER	3.1
3	A	431	GLY	3.0
1	H	11	LEU	3.0
3	A	516	GLU	3.0
2	L	5	THR	3.0
3	A	511	VAL	3.0
3	A	359	SER	3.0
1	H	208	ASN	2.9
1	H	18	LEU	2.9
2	L	53	SER	2.9
1	H	46	GLU	2.9
1	H	88	GLU	2.9
2	L	20	THR	2.9
3	A	400	PHE	2.9
3	A	433	VAL	2.9
1	H	14	PRO	2.8
3	A	398	ASP	2.8
3	A	413	GLY	2.8
2	L	52	ALA	2.8
3	A	403	LYS	2.8
2	L	99	GLY	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	A	429	PHE	2.8
3	A	504	ASN	2.7
2	L	72	PHE	2.7
2	L	214	CYS	2.7
3	A	410	ILE	2.7
2	L	212	GLY	2.7
2	L	29	VAL	2.7
3	A	500	THR	2.7
2	L	34	LEU	2.6
3	A	397	ALA	2.6
3	A	336	CYS	2.6
2	L	11	LEU	2.6
3	A	423	TYR	2.6
1	H	48	VAL	2.6
2	L	49	ILE	2.6
1	H	196	GLN	2.6
3	A	393	THR	2.5
2	L	69	GLY	2.5
3	A	476	GLY	2.5
2	L	68	SER	2.5
2	L	66	SER	2.5
2	L	92	TYR	2.5
2	L	100	GLN	2.5
2	L	70	THR	2.5
1	H	85	LEU	2.5
3	A	375	SER	2.5
2	L	7	SER	2.4
1	H	47	TRP	2.4
3	A	412	PRO	2.4
3	A	435	ALA	2.4
2	L	191	VAL	2.4
3	A	408	ARG	2.4
1	H	50	LEU	2.4
2	L	102	THR	2.4
2	L	32	SER	2.4
1	H	68	THR	2.3
2	L	81	PRO	2.3
2	L	78	ARG	2.3
1	H	69	ILE	2.3
2	L	192	TYR	2.3
3	A	354	ASN	2.3
1	H	30	SER	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	H	74	SER	2.3
3	A	405	ASP	2.3
3	A	501	THR	2.3
2	L	71	ASP	2.3
3	A	520	GLY	2.3
3	A	452	LEU	2.3
2	L	63	PHE	2.3
3	A	490	TYR	2.2
1	H	116	SER	2.2
2	L	94	SER	2.2
3	A	355	ARG	2.2
1	H	117	SER	2.2
1	H	211	VAL	2.2
3	A	523	THR	2.2
1	H	53	SER	2.2
3	A	415	THR	2.2
2	L	21	LEU	2.2
3	A	424	LYS	2.1
1	H	20	LEU	2.1
3	A	468	ILE	2.1
1	H	49	SER	2.1
3	A	493	GLU	2.1
1	H	151	PRO	2.1
1	H	82	MET	2.1
3	A	428	ASP	2.1
1	H	66	ARG	2.1
2	L	1	GLU	2.1
3	A	352	ALA	2.1
2	L	27	GLN	2.1
3	A	347	PHE	2.1
3	A	353	TRP	2.0
2	L	35	ALA	2.0
3	A	339	GLY	2.0
3	A	350	VAL	2.0
3	A	494	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.