



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 09:01 AM UTC

PDB ID : 1XQR / pdb_00001xqr
Title : Crystal structure of the HspBP1 core domain
Authors : Shomura, Y.; Dragovic, Z.; Chang, H.C.; Tzvetkov, N.; Young, J.C.; Brodsky, J.L.; Guerriero, V.; Hartl, F.U.; Bracher, A.
Deposited on : 2004-10-13
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	296	9% 74% 14% • 9%
1	B	296	6% 71% 16% • 9%

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 4262 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HspBP1 protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	268	Total	C	N	O	S	Se	0	0	0
			2058	1286	370	381	12	9			
1	B	268	Total	C	N	O	S	Se	0	0	0
			2060	1286	372	381	12	9			

There are 60 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	64	GLY	-	expression tag	UNP Q9NZL4
A	65	SER	-	expression tag	UNP Q9NZL4
A	66	SER	-	expression tag	UNP Q9NZL4
A	67	HIS	-	expression tag	UNP Q9NZL4
A	68	HIS	-	expression tag	UNP Q9NZL4
A	69	HIS	-	expression tag	UNP Q9NZL4
A	70	HIS	-	expression tag	UNP Q9NZL4
A	71	HIS	-	expression tag	UNP Q9NZL4
A	72	HIS	-	expression tag	UNP Q9NZL4
A	73	SER	-	expression tag	UNP Q9NZL4
A	74	SER	-	expression tag	UNP Q9NZL4
A	75	GLY	-	expression tag	UNP Q9NZL4
A	76	LEU	-	expression tag	UNP Q9NZL4
A	77	VAL	-	expression tag	UNP Q9NZL4
A	78	PRO	-	expression tag	UNP Q9NZL4
A	79	ARG	-	expression tag	UNP Q9NZL4
A	80	GLY	-	expression tag	UNP Q9NZL4
A	81	SER	-	expression tag	UNP Q9NZL4
A	82	HIS	-	expression tag	UNP Q9NZL4
A	83	MSE	-	expression tag	UNP Q9NZL4
A	88	GLY	GLU	engineered mutation	UNP Q9NZL4
A	93	MSE	MET	modified residue	UNP Q9NZL4
A	104	MSE	MET	modified residue	UNP Q9NZL4
A	134	MSE	MET	modified residue	UNP Q9NZL4
A	146	MSE	MET	modified residue	UNP Q9NZL4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	236	MSE	MET	modified residue	UNP Q9NZL4
A	239	MSE	MET	modified residue	UNP Q9NZL4
A	271	MSE	MET	modified residue	UNP Q9NZL4
A	273	MSE	MET	modified residue	UNP Q9NZL4
A	357	MSE	MET	modified residue	UNP Q9NZL4
B	64	GLY	-	expression tag	UNP Q9NZL4
B	65	SER	-	expression tag	UNP Q9NZL4
B	66	SER	-	expression tag	UNP Q9NZL4
B	67	HIS	-	expression tag	UNP Q9NZL4
B	68	HIS	-	expression tag	UNP Q9NZL4
B	69	HIS	-	expression tag	UNP Q9NZL4
B	70	HIS	-	expression tag	UNP Q9NZL4
B	71	HIS	-	expression tag	UNP Q9NZL4
B	72	HIS	-	expression tag	UNP Q9NZL4
B	73	SER	-	expression tag	UNP Q9NZL4
B	74	SER	-	expression tag	UNP Q9NZL4
B	75	GLY	-	expression tag	UNP Q9NZL4
B	76	LEU	-	expression tag	UNP Q9NZL4
B	77	VAL	-	expression tag	UNP Q9NZL4
B	78	PRO	-	expression tag	UNP Q9NZL4
B	79	ARG	-	expression tag	UNP Q9NZL4
B	80	GLY	-	expression tag	UNP Q9NZL4
B	81	SER	-	expression tag	UNP Q9NZL4
B	82	HIS	-	expression tag	UNP Q9NZL4
B	83	MSE	-	expression tag	UNP Q9NZL4
B	88	GLY	GLU	engineered mutation	UNP Q9NZL4
B	93	MSE	MET	modified residue	UNP Q9NZL4
B	104	MSE	MET	modified residue	UNP Q9NZL4
B	134	MSE	MET	modified residue	UNP Q9NZL4
B	146	MSE	MET	modified residue	UNP Q9NZL4
B	236	MSE	MET	modified residue	UNP Q9NZL4
B	239	MSE	MET	modified residue	UNP Q9NZL4
B	271	MSE	MET	modified residue	UNP Q9NZL4
B	273	MSE	MET	modified residue	UNP Q9NZL4
B	357	MSE	MET	modified residue	UNP Q9NZL4

- Molecule 2 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	76	Total O 76 76	0	0
2	B	68	Total O 68 68	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: HspBP1 protein

Chain A: 



• Molecule 1: HspBP1 protein

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	76.68Å 84.29Å 89.94Å 90.00° 96.00° 90.00°	Depositor
Resolution (Å)	38.07 – 2.10 38.07 – 2.10	Depositor EDS
% Data completeness (in resolution range)	(Not available) (38.07-2.10) 93.2 (38.07-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.36 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.0	Depositor
R, R_{free}	0.211 , 0.249 0.226 , 0.258	Depositor DCC
R_{free} test set	1568 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	26.4	Xtriage
Anisotropy	0.101	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 34.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4262	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.12	4/2077 (0.2%)	1.08	1/2786 (0.0%)
1	B	1.20	5/2079 (0.2%)	1.11	2/2789 (0.1%)
All	All	1.16	9/4156 (0.2%)	1.10	3/5575 (0.1%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	236	MSE	SE-CE	-10.47	1.64	1.95
1	A	236	MSE	SE-CE	-9.02	1.68	1.95
1	B	146	MSE	SE-CE	-7.82	1.72	1.95
1	A	83	MSE	SE-CE	6.13	2.13	1.95
1	B	134	MSE	SE-CE	6.08	2.13	1.95
1	B	273	MSE	SE-CE	5.95	2.13	1.95
1	A	194	ARG	C-O	-5.38	1.17	1.24
1	B	83	MSE	SE-CE	5.31	2.11	1.95
1	A	134	MSE	SE-CE	5.01	2.10	1.95

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	174	GLN	CA-C-N	-5.64	113.71	123.25
1	B	174	GLN	C-N-CA	-5.64	113.71	123.25
1	A	119	GLU	N-CA-C	-5.16	105.73	111.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2058	0	2060	33	0
1	B	2060	0	2060	37	0
2	A	76	0	0	2	0
2	B	68	0	0	1	0
All	All	4262	0	4120	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:149:LEU:HD21	1:A:168:LEU:HD13	1.56	0.88
1:B:286:SER:HB2	1:B:287:PRO:HD2	1.60	0.81
1:A:275:GLN:HE21	1:A:275:GLN:H	1.31	0.77
1:B:296:LEU:HD23	1:B:344:LEU:HD13	1.66	0.76
1:A:273:MSE:HE3	1:A:299:LEU:HD11	1.67	0.75
1:A:102:GLN:O	1:A:120:ARG:NH2	2.24	0.69
1:A:273:MSE:CE	1:A:299:LEU:HD11	2.22	0.68
1:B:197:ASP:OD1	1:B:237:ARG:NH1	2.28	0.67
1:B:296:LEU:HD23	1:B:344:LEU:CD1	2.24	0.67
1:B:286:SER:HB2	1:B:287:PRO:CD	2.24	0.66
1:B:278:VAL:HG11	1:B:315:LEU:HD22	1.78	0.65
1:A:116:ASP:OD2	1:A:120:ARG:NH1	2.30	0.64
1:A:293:LEU:HD12	1:A:337:GLU:HG3	1.80	0.64
1:B:209:LEU:HD23	1:B:209:LEU:O	1.99	0.62
1:A:268:LEU:HD23	1:A:273:MSE:HE1	1.82	0.61
1:B:346:GLN:HA	1:B:350:SER:HB2	1.84	0.60
1:B:234:VAL:HA	1:B:237:ARG:HH12	1.68	0.59
1:B:258:LEU:HD21	1:B:268:LEU:CD1	2.32	0.59
1:A:258:LEU:HD11	1:A:273:MSE:HE1	1.86	0.58
1:A:317:LEU:HD22	1:A:321:LEU:HG	1.84	0.58
1:A:275:GLN:HE21	1:A:275:GLN:N	2.02	0.56
1:B:133:ASN:HB3	1:B:136:ASN:HD22	1.71	0.56
1:B:245:LYS:HA	1:B:245:LYS:HE3	1.86	0.56
1:A:330:GLN:HA	2:B:600:HOH:O	2.06	0.55
1:B:87:ARG:O	1:B:91:GLU:HG2	2.07	0.55
1:A:258:LEU:HD11	1:A:273:MSE:CE	2.38	0.53
1:B:289:HIS:O	1:B:292:VAL:HG22	2.08	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:286:SER:CB	1:B:287:PRO:HD2	2.35	0.53
1:B:97:LEU:O	1:B:101:SER:HB2	2.09	0.53
1:A:133:ASN:HB3	1:A:136:ASN:HD22	1.73	0.52
1:A:268:LEU:HD23	1:A:273:MSE:CE	2.40	0.52
1:A:285:HIS:HE1	1:A:337:GLU:OE1	1.93	0.52
1:B:300:VAL:HG13	1:B:347:THR:HG21	1.93	0.50
1:B:237:ARG:HH11	1:B:237:ARG:HB2	1.77	0.50
1:B:236:MSE:CE	1:B:240:GLN:OE1	2.60	0.50
1:A:275:GLN:H	1:A:275:GLN:NE2	2.05	0.49
1:B:258:LEU:HD21	1:B:268:LEU:HD12	1.94	0.49
1:A:87:ARG:O	1:A:91:GLU:HG2	2.12	0.49
1:A:331:HIS:HB2	1:A:334:TYR:HD1	1.78	0.48
1:B:285:HIS:HE1	1:B:337:GLU:OE1	1.96	0.48
1:B:236:MSE:CE	1:B:240:GLN:CD	2.87	0.48
1:A:281:VAL:CG2	1:A:293:LEU:HD21	2.45	0.47
1:B:209:LEU:HD23	1:B:209:LEU:C	2.39	0.46
1:B:234:VAL:HA	1:B:237:ARG:NH1	2.29	0.46
1:B:83:MSE:HE1	1:B:94:LYS:NZ	2.30	0.46
1:B:282:ARG:CG	1:B:320:LEU:HD21	2.45	0.45
1:A:105:PRO:HB2	1:A:108:ALA:HB2	1.99	0.45
1:A:245:LYS:HB2	1:A:249:LYS:NZ	2.32	0.44
1:A:241:GLN:OE1	1:A:243:VAL:HB	2.18	0.44
1:B:105:PRO:HA	1:B:106:PRO:HD3	1.89	0.44
1:B:115:ALA:O	1:B:119:GLU:HG2	2.17	0.44
1:B:265:LYS:HG2	1:B:299:LEU:HD23	2.00	0.44
1:A:206:VAL:HG21	1:A:245:LYS:HD3	2.00	0.43
1:A:293:LEU:HD12	1:A:337:GLU:CG	2.48	0.43
1:B:289:HIS:HA	1:B:292:VAL:HG22	1.99	0.43
1:A:286:SER:HB2	1:A:287:PRO:CD	2.48	0.43
1:A:122:GLY:O	1:A:126:LEU:HD13	2.19	0.43
1:A:236:MSE:CE	1:A:240:GLN:OE1	2.67	0.42
1:A:149:LEU:HD21	1:A:168:LEU:CD1	2.40	0.42
1:A:330:GLN:HG2	1:B:174:GLN:HE22	1.85	0.41
1:A:285:HIS:HA	1:A:289:HIS:CE1	2.55	0.41
1:B:286:SER:CB	1:B:287:PRO:CD	2.92	0.41
1:B:282:ARG:HG3	1:B:320:LEU:HD21	2.03	0.41
1:A:155:GLU:OE2	1:A:194:ARG:NH2	2.54	0.41
2:A:641:HOH:O	1:B:330:GLN:HA	2.20	0.41
1:B:209:LEU:C	1:B:209:LEU:CD2	2.94	0.41
1:B:311:ARG:O	1:B:312:GLU:C	2.62	0.41
1:A:206:VAL:HG12	2:A:583:HOH:O	2.19	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:173:SER:O	1:B:174:GLN:C	2.63	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	266/296 (90%)	261 (98%)	5 (2%)	0	100	100
1	B	266/296 (90%)	260 (98%)	4 (2%)	2 (1%)	16	12
All	All	532/592 (90%)	521 (98%)	9 (2%)	2 (0%)	30	28

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	329	GLN
1	B	106	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	218/239 (91%)	204 (94%)	14 (6%)	16	14
1	B	218/239 (91%)	203 (93%)	15 (7%)	14	12
All	All	436/478 (91%)	407 (93%)	29 (7%)	15	12

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	130	LEU
1	A	149	LEU
1	A	168	LEU
1	A	189	LEU
1	A	194	ARG
1	A	216	VAL
1	A	241	GLN
1	A	244	GLN
1	A	263	GLU
1	A	268	LEU
1	A	275	GLN
1	A	317	LEU
1	A	324	ARG
1	A	350	SER
1	B	101	SER
1	B	104	MSE
1	B	130	LEU
1	B	195	LEU
1	B	224	LEU
1	B	227	LEU
1	B	237	ARG
1	B	245	LYS
1	B	246	LEU
1	B	253	LEU
1	B	282	ARG
1	B	293	LEU
1	B	300	VAL
1	B	315	LEU
1	B	336	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (26) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	113	GLN
1	A	136	ASN
1	A	174	GLN
1	A	219	GLN
1	A	225	GLN
1	A	240	GLN
1	A	244	GLN
1	A	275	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	276	GLN
1	A	285	HIS
1	A	289	HIS
1	A	330	GLN
1	A	335	GLN
1	B	113	GLN
1	B	117	GLN
1	B	118	GLN
1	B	136	ASN
1	B	147	HIS
1	B	167	GLN
1	B	174	GLN
1	B	244	GLN
1	B	261	HIS
1	B	276	GLN
1	B	285	HIS
1	B	329	GLN
1	B	331	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	259/296 (87%)	0.63	26 (10%) 12 13	21, 31, 49, 76	0
1	B	259/296 (87%)	0.52	17 (6%) 24 26	21, 28, 46, 61	0
All	All	518/592 (87%)	0.57	43 (8%) 17 18	21, 30, 48, 76	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	331	HIS	4.0
1	A	327	LEU	3.6
1	A	109	GLY	3.5
1	A	106	PRO	3.5
1	B	282	ARG	3.4
1	B	330	GLN	3.4
1	B	287	PRO	3.4
1	A	111	ALA	3.1
1	A	331	HIS	3.1
1	A	108	ALA	3.1
1	A	329	GLN	3.0
1	B	175	ASN	2.9
1	B	350	SER	2.8
1	A	112	GLU	2.8
1	B	108	ALA	2.8
1	B	327	LEU	2.8
1	A	330	GLN	2.7
1	B	106	PRO	2.7
1	B	323	HIS	2.7
1	B	296	LEU	2.6
1	B	334	TYR	2.6
1	A	346	GLN	2.4
1	A	350	SER	2.4
1	B	224	LEU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	101	SER	2.4
1	A	110	GLU	2.3
1	A	107	THR	2.3
1	B	107	THR	2.3
1	B	228	ARG	2.3
1	A	249	LYS	2.3
1	A	288	PHE	2.3
1	A	105	PRO	2.3
1	A	287	PRO	2.3
1	B	335	GLN	2.1
1	A	323	HIS	2.1
1	A	84	ARG	2.1
1	A	336	GLU	2.1
1	B	220	GLU	2.0
1	A	86	GLN	2.0
1	A	241	GLN	2.0
1	A	335	GLN	2.0
1	A	334	TYR	2.0
1	A	216	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.