



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 05:23 AM UTC

PDB ID : 1XRH / pdb_00001xrh
Title : Crystal Structure of Ureidoglycolate Dehydrogenase from Escherichia Coli
Authors : Rajashankar, K.R.; Kniewel, R.; Lima, C.D.; Burley, S.K.; New York SGX
Research Center for Structural Genomics (NYSGXRC)
Deposited on : 2004-10-14
Resolution : 2.25 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1898 (2.26-2.26)
Clashscore	190562	2005 (2.26-2.26)
Ramachandran outliers	187476	1965 (2.26-2.26)
Sidechain outliers	187428	1966 (2.26-2.26)
RSRZ outliers	180081	1898 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	351	% 64% 28% 5% .
1	B	351	% 66% 26% 5% .
1	C	351	65% 30% . .
1	D	351	66% 28% . .
1	E	351	% 61% 32% 5% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	351	 65% 28% . .
1	G	351	 68% 25% . .
1	H	351	 71% 23% . .

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 21480 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ureidoglycolate Dehydrogenase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	343	Total	C	N	O	S	Se	0	0	0
			2616	1647	457	497	3	12			
1	B	343	Total	C	N	O	S	Se	0	0	0
			2616	1647	457	497	3	12			
1	C	344	Total	C	N	O	S	Se	0	0	0
			2622	1650	458	499	3	12			
1	D	342	Total	C	N	O	S	Se	0	0	0
			2607	1642	456	494	3	12			
1	E	343	Total	C	N	O	S	Se	0	0	0
			2616	1647	457	497	3	12			
1	F	339	Total	C	N	O	S	Se	0	0	0
			2582	1626	453	488	3	12			
1	G	341	Total	C	N	O	S	Se	0	0	0
			2595	1633	455	492	3	12			
1	H	339	Total	C	N	O	S	Se	0	0	0
			2582	1626	453	488	3	12			

There are 120 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	SER	-	cloning artifact	UNP P77555
A	2	SER	-	cloning artifact	UNP P77555
A	3	SER	MET	cloning artifact	UNP P77555
A	94	MSE	MET	modified residue	UNP P77555
A	96	MSE	MET	modified residue	UNP P77555
A	116	MSE	MET	modified residue	UNP P77555
A	139	MSE	MET	modified residue	UNP P77555
A	145	MSE	MET	modified residue	UNP P77555
A	177	MSE	MET	modified residue	UNP P77555
A	195	MSE	MET	modified residue	UNP P77555
A	231	MSE	MET	modified residue	UNP P77555
A	232	MSE	MET	modified residue	UNP P77555
A	233	MSE	MET	modified residue	UNP P77555

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	253	MSE	MET	modified residue	UNP P77555
A	289	MSE	MET	modified residue	UNP P77555
B	1	SER	-	cloning artifact	UNP P77555
B	2	SER	-	cloning artifact	UNP P77555
B	3	SER	MET	cloning artifact	UNP P77555
B	94	MSE	MET	modified residue	UNP P77555
B	96	MSE	MET	modified residue	UNP P77555
B	116	MSE	MET	modified residue	UNP P77555
B	139	MSE	MET	modified residue	UNP P77555
B	145	MSE	MET	modified residue	UNP P77555
B	177	MSE	MET	modified residue	UNP P77555
B	195	MSE	MET	modified residue	UNP P77555
B	231	MSE	MET	modified residue	UNP P77555
B	232	MSE	MET	modified residue	UNP P77555
B	233	MSE	MET	modified residue	UNP P77555
B	253	MSE	MET	modified residue	UNP P77555
B	289	MSE	MET	modified residue	UNP P77555
C	1	SER	-	cloning artifact	UNP P77555
C	2	SER	-	cloning artifact	UNP P77555
C	3	SER	MET	cloning artifact	UNP P77555
C	94	MSE	MET	modified residue	UNP P77555
C	96	MSE	MET	modified residue	UNP P77555
C	116	MSE	MET	modified residue	UNP P77555
C	139	MSE	MET	modified residue	UNP P77555
C	145	MSE	MET	modified residue	UNP P77555
C	177	MSE	MET	modified residue	UNP P77555
C	195	MSE	MET	modified residue	UNP P77555
C	231	MSE	MET	modified residue	UNP P77555
C	232	MSE	MET	modified residue	UNP P77555
C	233	MSE	MET	modified residue	UNP P77555
C	253	MSE	MET	modified residue	UNP P77555
C	289	MSE	MET	modified residue	UNP P77555
D	1	SER	-	cloning artifact	UNP P77555
D	2	SER	-	cloning artifact	UNP P77555
D	3	SER	MET	cloning artifact	UNP P77555
D	94	MSE	MET	modified residue	UNP P77555
D	96	MSE	MET	modified residue	UNP P77555
D	116	MSE	MET	modified residue	UNP P77555
D	139	MSE	MET	modified residue	UNP P77555
D	145	MSE	MET	modified residue	UNP P77555
D	177	MSE	MET	modified residue	UNP P77555
D	195	MSE	MET	modified residue	UNP P77555

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	231	MSE	MET	modified residue	UNP P77555
D	232	MSE	MET	modified residue	UNP P77555
D	233	MSE	MET	modified residue	UNP P77555
D	253	MSE	MET	modified residue	UNP P77555
D	289	MSE	MET	modified residue	UNP P77555
E	1	SER	-	cloning artifact	UNP P77555
E	2	SER	-	cloning artifact	UNP P77555
E	3	SER	MET	cloning artifact	UNP P77555
E	94	MSE	MET	modified residue	UNP P77555
E	96	MSE	MET	modified residue	UNP P77555
E	116	MSE	MET	modified residue	UNP P77555
E	139	MSE	MET	modified residue	UNP P77555
E	145	MSE	MET	modified residue	UNP P77555
E	177	MSE	MET	modified residue	UNP P77555
E	195	MSE	MET	modified residue	UNP P77555
E	231	MSE	MET	modified residue	UNP P77555
E	232	MSE	MET	modified residue	UNP P77555
E	233	MSE	MET	modified residue	UNP P77555
E	253	MSE	MET	modified residue	UNP P77555
E	289	MSE	MET	modified residue	UNP P77555
F	1	SER	-	cloning artifact	UNP P77555
F	2	SER	-	cloning artifact	UNP P77555
F	3	SER	MET	cloning artifact	UNP P77555
F	94	MSE	MET	modified residue	UNP P77555
F	96	MSE	MET	modified residue	UNP P77555
F	116	MSE	MET	modified residue	UNP P77555
F	139	MSE	MET	modified residue	UNP P77555
F	145	MSE	MET	modified residue	UNP P77555
F	177	MSE	MET	modified residue	UNP P77555
F	195	MSE	MET	modified residue	UNP P77555
F	231	MSE	MET	modified residue	UNP P77555
F	232	MSE	MET	modified residue	UNP P77555
F	233	MSE	MET	modified residue	UNP P77555
F	253	MSE	MET	modified residue	UNP P77555
F	289	MSE	MET	modified residue	UNP P77555
G	1	SER	-	cloning artifact	UNP P77555
G	2	SER	-	cloning artifact	UNP P77555
G	3	SER	MET	cloning artifact	UNP P77555
G	94	MSE	MET	modified residue	UNP P77555
G	96	MSE	MET	modified residue	UNP P77555
G	116	MSE	MET	modified residue	UNP P77555
G	139	MSE	MET	modified residue	UNP P77555

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	145	MSE	MET	modified residue	UNP P77555
G	177	MSE	MET	modified residue	UNP P77555
G	195	MSE	MET	modified residue	UNP P77555
G	231	MSE	MET	modified residue	UNP P77555
G	232	MSE	MET	modified residue	UNP P77555
G	233	MSE	MET	modified residue	UNP P77555
G	253	MSE	MET	modified residue	UNP P77555
G	289	MSE	MET	modified residue	UNP P77555
H	1	SER	-	cloning artifact	UNP P77555
H	2	SER	-	cloning artifact	UNP P77555
H	3	SER	MET	cloning artifact	UNP P77555
H	94	MSE	MET	modified residue	UNP P77555
H	96	MSE	MET	modified residue	UNP P77555
H	116	MSE	MET	modified residue	UNP P77555
H	139	MSE	MET	modified residue	UNP P77555
H	145	MSE	MET	modified residue	UNP P77555
H	177	MSE	MET	modified residue	UNP P77555
H	195	MSE	MET	modified residue	UNP P77555
H	231	MSE	MET	modified residue	UNP P77555
H	232	MSE	MET	modified residue	UNP P77555
H	233	MSE	MET	modified residue	UNP P77555
H	253	MSE	MET	modified residue	UNP P77555
H	289	MSE	MET	modified residue	UNP P77555


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	58	Total O 58 58	0	0
2	B	102	Total O 102 102	0	0
2	C	77	Total O 77 77	0	0
2	D	113	Total O 113 113	0	0
2	E	61	Total O 61 61	0	0
2	F	70	Total O 70 70	0	0
2	G	69	Total O 69 69	0	0
2	H	94	Total O 94 94	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Ureidoglycolate Dehydrogenase




• Molecule 1: Ureidoglycolate Dehydrogenase




• Molecule 1: Ureidoglycolate Dehydrogenase






● Molecule 1: Ureidoglycolate Dehydrogenase

Chain D: 66% 28% . .



● Molecule 1: Ureidoglycolate Dehydrogenase


Chain E: 61% 32% 5% .



● Molecule 1: Ureidoglycolate Dehydrogenase


Chain F: 65% 28% . .






• Molecule 1: Ureidoglycolate Dehydrogenase

Chain G: 68% 25% . .



• Molecule 1: Ureidoglycolate Dehydrogenase

Chain H: 71% 23% . .



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	104.72Å 129.92Å 106.47Å 90.00° 95.77° 90.00°	Depositor
Resolution (Å)	19.86 – 2.25 19.86 – 2.25	Depositor EDS
% Data completeness (in resolution range)	75.6 (19.86-2.25) 87.5 (19.86-2.25)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.44 (at 2.26Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.192 , 0.250 0.206 , 0.262	Depositor DCC
R_{free} test set	9652 reflections (4.53%)	wwPDB-VP
Wilson B-factor (Å ²)	33.5	Xtriage
Anisotropy	0.157	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 52.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.016 for l,-k,h	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	21480	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.98	0/2660	1.17	12/3587 (0.3%)
1	B	1.08	0/2660	1.23	20/3587 (0.6%)
1	C	0.99	2/2666 (0.1%)	1.22	14/3595 (0.4%)
1	D	1.05	3/2651 (0.1%)	1.21	11/3575 (0.3%)
1	E	1.04	1/2660 (0.0%)	1.19	9/3587 (0.3%)
1	F	1.03	0/2625	1.23	15/3539 (0.4%)
1	G	1.03	1/2638 (0.0%)	1.15	6/3557 (0.2%)
1	H	1.02	2/2625 (0.1%)	1.21	15/3539 (0.4%)
All	All	1.03	9/21185 (0.0%)	1.20	102/28566 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	2
1	C	0	2
1	G	0	1
All	All	0	6

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	144	PRO	CA-C	6.29	1.59	1.52
1	G	231	MSE	SE-CE	6.26	2.14	1.95
1	H	147	VAL	CA-CB	5.80	1.61	1.54
1	D	181	VAL	CA-CB	5.77	1.60	1.54
1	D	320	VAL	CA-CB	5.62	1.62	1.54

The worst 5 of 102 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	306	TYR	CA-C-N	-10.74	108.70	119.76
1	F	306	TYR	C-N-CA	-10.74	108.70	119.76
1	F	237	LEU	N-CA-C	8.04	120.04	111.28
1	C	51	VAL	N-CA-C	8.00	118.10	110.42
1	C	177	MSE	N-CA-C	7.93	121.33	108.32

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	343	TYR	Sidechain
1	B	228	TYR	Sidechain
1	B	343	TYR	Sidechain
1	C	228	TYR	Sidechain
1	C	330	TYR	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2616	0	2559	92	0
1	B	2616	0	2559	89	0
1	C	2622	0	2567	79	0
1	D	2607	0	2553	76	0
1	E	2616	0	2559	110	0
1	F	2582	0	2532	79	0
1	G	2595	0	2544	89	0
1	H	2582	0	2532	53	0
2	A	58	0	0	0	0
2	B	102	0	0	5	0
2	C	77	0	0	3	0
2	D	113	0	0	4	0
2	E	61	0	0	7	0
2	F	70	0	0	1	0
2	G	69	0	0	7	0
2	H	94	0	0	3	0
All	All	21480	0	20405	613	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 613 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:50:ARG:HH21	1:E:253:MSE:HE1	1.09	1.15
1:G:232:MSE:HE2	1:G:236:VAL:HG21	1.27	1.14
1:A:153:GLU:HG3	1:B:295:ILE:HG23	1.34	1.10
1:A:232:MSE:HE3	1:B:173:LEU:HG	1.31	1.09
1:C:232:MSE:HE3	1:D:173:LEU:HD13	1.34	1.08

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	341/351 (97%)	317 (93%)	21 (6%)	3 (1%)	14	12
1	B	341/351 (97%)	323 (95%)	14 (4%)	4 (1%)	10	7
1	C	342/351 (97%)	332 (97%)	9 (3%)	1 (0%)	36	40
1	D	340/351 (97%)	329 (97%)	9 (3%)	2 (1%)	21	21
1	E	341/351 (97%)	319 (94%)	20 (6%)	2 (1%)	21	21
1	F	337/351 (96%)	321 (95%)	12 (4%)	4 (1%)	10	7
1	G	339/351 (97%)	326 (96%)	12 (4%)	1 (0%)	36	40
1	H	337/351 (96%)	325 (96%)	10 (3%)	2 (1%)	21	21
All	All	2718/2808 (97%)	2592 (95%)	107 (4%)	19 (1%)	18	17

5 of 19 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	195	MSE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	194	ASN
1	B	192	SER
1	B	194	ASN
1	G	194	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	272/267 (102%)	245 (90%)	27 (10%)	7	5
1	B	272/267 (102%)	253 (93%)	19 (7%)	14	13
1	C	273/267 (102%)	259 (95%)	14 (5%)	21	23
1	D	271/267 (102%)	247 (91%)	24 (9%)	9	7
1	E	272/267 (102%)	250 (92%)	22 (8%)	11	9
1	F	268/267 (100%)	245 (91%)	23 (9%)	10	8
1	G	270/267 (101%)	252 (93%)	18 (7%)	15	14
1	H	268/267 (100%)	251 (94%)	17 (6%)	16	16
All	All	2166/2136 (101%)	2002 (92%)	164 (8%)	12	10

5 of 164 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	123	SER
1	G	285	LEU
1	F	145	MSE
1	G	18	LEU
1	H	94	MSE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 76 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	249	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	H	128	GLN
1	F	303	GLN
1	G	194	ASN
1	H	293	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	331/351 (94%)	-0.05	2 (0%) 85 86	25, 40, 59, 75	0
1	B	331/351 (94%)	-0.19	2 (0%) 85 86	21, 33, 54, 70	0
1	C	332/351 (94%)	-0.17	1 (0%) 90 90	19, 37, 53, 68	0
1	D	330/351 (94%)	-0.29	1 (0%) 90 90	21, 34, 50, 67	0
1	E	331/351 (94%)	-0.05	4 (1%) 76 77	21, 37, 59, 75	0
1	F	327/351 (93%)	-0.12	0 100 100	24, 38, 54, 67	0
1	G	329/351 (93%)	-0.25	1 (0%) 90 90	22, 35, 58, 73	0
1	H	327/351 (93%)	-0.28	1 (0%) 90 90	19, 32, 55, 65	0
All	All	2638/2808 (93%)	-0.17	12 (0%) 87 88	19, 36, 56, 75	0

The worst 5 of 12 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	2	SER	3.4
1	G	194	ASN	2.8
1	C	2	SER	2.8
1	B	343	TYR	2.6
1	E	194	ASN	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.