



Full wwPDB NMR Structure Validation Report ⓘ

Mar 30, 2026 – 01:15 AM UTC

PDB ID : 6XRY / pdb_00006xry
BMRB ID : 30773
Title : Intrinsically disordered bacterial polar organizing protein Z, PopZ, interacts with protein binding partners through an N-terminal Molecular Recognition Feature
Authors : Nordyke, C.T.; Ahmed, Y.M.; Puterbaugh, R.Z.; Bowman, G.R.; Varga, K.
Deposited on : 2020-07-14

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
BMRB Restraints Analysis : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 63%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	141	

2 Ensemble composition and analysis

This entry contains 20 models. Model 17 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *medoid*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:1-A:18 (18)	2.46	17

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 2, 5, 7, 9, 18, 19, 20
2	3, 4, 12, 15, 17
3	8, 11, 13, 16
4	6, 10, 14

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2034 atoms, of which 979 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Polar organizing protein Z.

Mol	Chain	Residues	Atoms						Trace
1	A	141	Total	C	H	N	O	S	0
			2034	660	979	166	226	3	

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	134	LEU	-	expression tag	UNP Q9A8N4
A	135	GLU	-	expression tag	UNP Q9A8N4
A	136	HIS	-	expression tag	UNP Q9A8N4
A	137	HIS	-	expression tag	UNP Q9A8N4
A	138	HIS	-	expression tag	UNP Q9A8N4
A	139	HIS	-	expression tag	UNP Q9A8N4
A	140	HIS	-	expression tag	UNP Q9A8N4
A	141	HIS	-	expression tag	UNP Q9A8N4

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Polar organizing protein Z





4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1


- Molecule 1: Polar organizing protein Z






4.2.5 Score per residue for model 5

- Molecule 1: Polar organizing protein Z




4.2.6 Score per residue for model 6

- Molecule 1: Polar organizing protein Z




4.2.7 Score per residue for model 7

- Molecule 1: Polar organizing protein Z




4.2.8 Score per residue for model 8

- Molecule 1: Polar organizing protein Z




4.2.9 Score per residue for model 9

- Molecule 1: Polar organizing protein Z




4.2.10 Score per residue for model 10

- Molecule 1: Polar organizing protein Z




4.2.11 Score per residue for model 11

- Molecule 1: Polar organizing protein Z




4.2.12 Score per residue for model 12

- Molecule 1: Polar organizing protein Z




4.2.13 Score per residue for model 13

- Molecule 1: Polar organizing protein Z




4.2.14 Score per residue for model 14

- Molecule 1: Polar organizing protein Z




4.2.15 Score per residue for model 15

- Molecule 1: Polar organizing protein Z




4.2.16 Score per residue for model 16

- Molecule 1: Polar organizing protein Z




4.2.17 Score per residue for model 17 (medoid)

- Molecule 1: Polar organizing protein Z




4.2.18 Score per residue for model 18

- Molecule 1: Polar organizing protein Z




4.2.19 Score per residue for model 19

- Molecule 1: Polar organizing protein Z



4.2.20 Score per residue for model 20

- Molecule 1: Polar organizing protein Z



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
PONDEROSA-C/S	structure calculation	
PONDEROSA-C/S	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1007
Number of shifts mapped to atoms	1007
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	63%

6 Model quality [i](#)

6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	141	138	138	1±1
All	All	2820	2760	2760	13

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:1:MET:HE3	1:A:11:GLU:CD	0.69	2.12	2	1
1:A:17:ILE:HD13	1:A:17:ILE:O	0.64	1.93	13	1
1:A:1:MET:HE3	1:A:11:GLU:OE2	0.54	2.02	2	1
1:A:14:LEU:HD13	1:A:14:LEU:O	0.47	2.10	17	1
1:A:14:LEU:C	1:A:14:LEU:HD23	0.47	2.35	15	4
1:A:1:MET:HE3	1:A:11:GLU:OE1	0.42	2.13	2	1
1:A:10:MET:O	1:A:14:LEU:HD13	0.41	2.16	1	1
1:A:13:ILE:N	1:A:13:ILE:HD12	0.41	2.31	4	1
1:A:16:SER:O	1:A:17:ILE:HB	0.41	2.16	19	1
1:A:14:LEU:HD23	1:A:14:LEU:C	0.40	2.41	13	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	17/141 (12%)	14±2 (84±9%)	2±1 (11±6%)	1±1 (5±5%)	2	22
All	All	340/2820 (12%)	284 (84%)	38 (11%)	18 (5%)	2	22

All 8 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	18	ARG	4
1	A	17	ILE	4
1	A	6	GLN	3
1	A	8	PRO	2
1	A	4	GLN	2
1	A	7	GLU	1
1	A	5	SER	1
1	A	2	SER	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	17/118 (14%)	17±0 (99±2%)	0±0 (1±2%)	68	95
All	All	340/2360 (14%)	337 (99%)	3 (1%)	68	95

All 3 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	1	MET	1
1	A	17	ILE	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	14	LEU	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 63% for the well-defined parts and 58% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1007
Number of shifts mapped to atoms	1007
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	116	-0.04 ± 0.04	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	109	0.17 ± 0.09	None needed (< 0.5 ppm)
$^{13}\text{C}'$	114	-0.15 ± 0.05	None needed (< 0.5 ppm)
^{15}N	103	-0.99 ± 0.10	Should be applied

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 63%, i.e. 150 atoms were assigned a chemical shift out of a possible 238. 0 out of 1 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	85/88 (97%)	33/35 (94%)	36/36 (100%)	16/17 (94%)
Sidechain	65/150 (43%)	37/97 (38%)	28/48 (58%)	0/5 (0%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Overall	150/238 (63%)	70/132 (53%)	64/84 (76%)	16/22 (73%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 58%, i.e. 1007 atoms were assigned a chemical shift out of a possible 1745. 0 out of 18 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	542/659 (82%)	209/261 (80%)	230/282 (82%)	103/116 (89%)
Sidechain	465/996 (47%)	271/653 (42%)	193/330 (58%)	1/13 (8%)
Aromatic	0/90 (0%)	0/47 (0%)	0/37 (0%)	0/6 (0%)
Overall	1007/1745 (58%)	480/961 (50%)	423/649 (65%)	104/135 (77%)


7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:



8 NMR restraints analysis

No restraints data found

9 Distance violation analysis ⓘ

No distance restraints data found

10 Dihedral-angle violation analysis ⓘ

No dihedral-angle restraints found