



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 10, 2026 – 03:14 AM UTC

PDB ID : 7XRM / pdb_00007xrm
Title : Ethanolamine ammonia-lyase complexed with AdoMeCbl
Authors : Shibata, N.; Toraya, T.
Deposited on : 2022-05-10
Resolution : 2.13 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.13 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3689 (2.16-2.12)
Clashscore	190562	3812 (2.16-2.12)
Ramachandran outliers	187476	3773 (2.16-2.12)
Sidechain outliers	187428	3772 (2.16-2.12)
RSRZ outliers	180081	3691 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	453	 89% 11%
1	C	453	 89% 10% .
2	B	263	 78% 17% . .
2	D	263	 2% 75% 21% .

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 11814 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ethanolamine ammonia-lyase large subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	453	Total	C	N	O	S	0	4	0
			3484	2184	597	681	22			
1	C	453	Total	C	N	O	S	0	5	0
			3492	2190	599	680	23			

- Molecule 2 is a protein called Ethanolamine ammonia-lyase small subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	253	Total	C	N	O	S	0	2	0
			1943	1213	354	366	10			
2	D	252	Total	C	N	O	S	0	6	0
			1956	1222	351	373	10			

There are 22 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	33	MET	-	initiating methionine	UNP P19636
B	34	ASP	-	expression tag	UNP P19636
B	35	GLN	-	expression tag	UNP P19636
B	36	SER	-	expression tag	UNP P19636
B	37	SER	-	expression tag	UNP P19636
B	38	HIS	-	expression tag	UNP P19636
B	39	HIS	-	expression tag	UNP P19636
B	40	HIS	-	expression tag	UNP P19636
B	41	HIS	-	expression tag	UNP P19636
B	42	HIS	-	expression tag	UNP P19636
B	43	HIS	-	expression tag	UNP P19636
D	33	MET	-	initiating methionine	UNP P19636
D	34	ASP	-	expression tag	UNP P19636
D	35	GLN	-	expression tag	UNP P19636
D	36	SER	-	expression tag	UNP P19636
D	37	SER	-	expression tag	UNP P19636

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	38	HIS	-	expression tag	UNP P19636
D	39	HIS	-	expression tag	UNP P19636
D	40	HIS	-	expression tag	UNP P19636
D	41	HIS	-	expression tag	UNP P19636
D	42	HIS	-	expression tag	UNP P19636
D	43	HIS	-	expression tag	UNP P19636

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	C	1	Total C O 6 3 3	0	0
3	C	1	Total C O 6 3 3	0	0
3	C	1	Total C O 6 3 3	0	0
3	C	1	Total C O 6 3 3	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is AMMONIUM ION (CCD ID: NH4) (formula: H₄N).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	N	0	0
			1	1		
4	C	1	Total	N	0	0
			1	1		

- Molecule 5 is (2 {R},3 {R},4 {S},5 {R})-2-(6-aminopurin-9-yl)-5-ethyl-oxolane-3,4-diol (CCD ID: FWK) (formula: C₁₁H₁₅N₅O₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			19	11	5	3		
5	C	1	Total	C	N	O	0	0
			19	11	5	3		

- Molecule 6 is COBALAMIN (CCD ID: B12) (formula: C₆₂H₈₉CoN₁₃O₁₄P) (labeled as "Lig- and of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
6	B	1	Total	C	Co	N	O	P	0	0
			91	62	1	13	14	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
6	D	1	Total	C	Co	N	O	P	0	0
			91	62	1	13	14	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	286	Total	O	0	0
			286	286		
7	B	72	Total	O	0	0
			72	72		
7	C	259	Total	O	0	0
			259	259		
7	D	34	Total	O	0	0
			34	34		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ethanolamine ammonia-lyase large subunit




- Molecule 1: Ethanolamine ammonia-lyase large subunit



- Molecule 2: Ethanolamine ammonia-lyase small subunit



- Molecule 2: Ethanolamine ammonia-lyase small subunit



G193	L194	K195	R203	Y204	K208	I209	E210	D211	G214	E215	I216	L217	V222	L225	L238	S239	C240	P246	R247	T251	V252	E253	T257	C258	I259	P268	P269	V270	E271	A272	A273	A274	V275	I276	V277	D278	L279	A280	K281	R282	M283	Q286	E295
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	242.88Å 242.88Å 76.64Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.42 – 2.13 46.42 – 2.13	Depositor EDS
% Data completeness (in resolution range)	99.2 (46.42-2.13) 99.2 (46.42-2.13)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.00 (at 2.14Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.194 , 0.220 (Not available) , 0.221	Depositor DCC
R_{free} test set	7190 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	34.6	Xtriage
Anisotropy	0.126	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 38.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.38$, $\langle L^2 \rangle = 0.21$	Xtriage
Estimated twinning fraction	0.085 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	11814	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NH4, FWK, GOL, B12

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.13	0/3550	0.29	0/4807
1	C	0.12	0/3561	0.28	0/4820
2	B	0.15	0/1974	0.34	0/2674
2	D	0.14	0/1992	0.34	0/2699
All	All	0.13	0/11077	0.31	0/15000

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3484	0	3452	36	0
1	C	3492	0	3468	31	0
2	B	1943	0	1998	32	0
2	D	1956	0	2012	38	0
3	A	24	0	32	3	0
3	C	42	0	56	2	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
5	A	19	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	C	19	0	0	0	0
6	B	91	0	88	9	0
6	D	91	0	88	8	0
7	A	286	0	0	5	0
7	B	72	0	0	0	0
7	C	259	0	0	0	0
7	D	34	0	0	0	0
All	All	11814	0	11194	148	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 148 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:160:ASP:HB2	2:B:194:LEU:HD13	1.60	0.82
2:B:283:MET:HE3	2:B:289:SER:HB3	1.67	0.75
2:D:216:ILE:HG22	2:D:217:LEU:HD23	1.71	0.73
2:B:186:MET:HE2	2:B:199:PRO:HG3	1.70	0.72
1:C:269[B]:ARG:NH1	1:C:318:TYR:O	2.26	0.68

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	455/453 (100%)	438 (96%)	16 (4%)	1 (0%)	43	42
1	C	456/453 (101%)	437 (96%)	18 (4%)	1 (0%)	43	42
2	B	253/263 (96%)	245 (97%)	8 (3%)	0	100	100
2	D	256/263 (97%)	242 (94%)	14 (6%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	1420/1432 (99%)	1362 (96%)	56 (4%)	2 (0%)	48	49

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	163	PRO
1	C	330	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	374/370 (101%)	372 (100%)	2 (0%)	81	86
1	C	375/370 (101%)	369 (98%)	6 (2%)	55	61
2	B	209/217 (96%)	206 (99%)	3 (1%)	59	65
2	D	212/217 (98%)	211 (100%)	1 (0%)	81	86
All	All	1170/1174 (100%)	1158 (99%)	12 (1%)	70	74

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	224[B]	VAL
1	C	326	VAL
2	D	128	SER
1	C	401[A]	MET
2	B	194	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	324	ASN
1	C	368	HIS
2	D	191	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	D	62	HIS
2	B	154	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 17 ligands modelled in this entry, 2 are modelled with single atom - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	C	504	-	5,5,5	1.03	0	5,5,5	1.03	0
3	GOL	A	502	-	5,5,5	0.93	0	5,5,5	1.01	0
3	GOL	C	507	-	5,5,5	1.06	0	5,5,5	0.88	0
3	GOL	C	503	-	5,5,5	0.96	0	5,5,5	1.02	0
5	FWK	C	509	-	21,21,21	1.53	4 (19%)	30,31,31	2.14	7 (23%)
3	GOL	C	506	-	5,5,5	0.96	0	5,5,5	1.05	0
6	B12	D	301	-	94,101,101	1.46	13 (13%)	149,166,166	1.42	18 (12%)
5	FWK	A	506	-	21,21,21	1.55	4 (19%)	30,31,31	2.11	7 (23%)
3	GOL	C	502	-	5,5,5	0.89	0	5,5,5	1.09	0
3	GOL	A	504	-	5,5,5	0.97	0	5,5,5	0.97	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	B12	B	301	-	94,101,101	1.46	13 (13%)	149,166,166	1.37	21 (14%)
3	GOL	A	503	-	5,5,5	0.98	0	5,5,5	1.20	1 (20%)
3	GOL	A	501	-	5,5,5	0.99	0	5,5,5	1.05	0
3	GOL	C	501	-	5,5,5	0.94	0	5,5,5	1.05	0
3	GOL	C	505	-	5,5,5	1.00	0	5,5,5	1.00	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	C	504	-	-	1/4/4/4	-
3	GOL	A	502	-	-	0/4/4/4	-
3	GOL	C	507	-	-	0/4/4/4	-
3	GOL	C	503	-	-	3/4/4/4	-
5	FWK	C	509	-	-	1/6/22/22	0/3/3/3
3	GOL	C	506	-	-	4/4/4/4	-
6	B12	D	301	-	-	22/56/223/223	0/3/11/11
5	FWK	A	506	-	-	0/6/22/22	0/3/3/3
3	GOL	C	502	-	-	0/4/4/4	-
3	GOL	A	504	-	-	0/4/4/4	-
6	B12	B	301	-	-	20/56/223/223	0/3/11/11
3	GOL	A	503	-	-	1/4/4/4	-
3	GOL	A	501	-	-	2/4/4/4	-
3	GOL	C	501	-	-	3/4/4/4	-
3	GOL	C	505	-	-	2/4/4/4	-

The worst 5 of 34 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	D	301	B12	C8B-C9B	5.36	1.49	1.40
6	B	301	B12	C8B-C9B	5.32	1.49	1.40
6	D	301	B12	C14-N23	-5.08	1.28	1.35
6	B	301	B12	C14-N23	-4.98	1.28	1.35
5	A	506	FWK	C5-C4	4.83	1.47	1.39

The worst 5 of 54 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	506	FWK	C5-C4-N3	-6.19	118.20	126.72
5	C	509	FWK	C5-C4-N3	-6.16	118.23	126.72
6	D	301	B12	C8B-C9B-N3B	-5.63	103.91	110.00
6	B	301	B12	C8B-C9B-N3B	-5.26	104.31	110.00
5	C	509	FWK	N3-C4-N9	5.22	136.04	127.17

There are no chirality outliers.

5 of 59 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	C	501	GOL	O1-C1-C2-C3
3	C	505	GOL	O1-C1-C2-O2
3	C	505	GOL	O1-C1-C2-C3
3	C	506	GOL	C1-C2-C3-O3
6	B	301	B12	C1-C2-C26-C27

There are no ring outliers.


6 monomers are involved in 22 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	D	301	B12	8	0
6	B	301	B12	9	0
3	A	503	GOL	2	0
3	A	501	GOL	1	0
3	C	501	GOL	1	0
3	C	505	GOL	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



Ligand B12 D 301







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å²)	Q<0.9
1	A	453/453 (100%)	-1.05	0	100	100	17, 34, 51, 74	4 (0%)
1	C	453/453 (100%)	-1.00	0	100	100	17, 35, 55, 75	5 (1%)
2	B	253/263 (96%)	-0.41	0	100	100	29, 58, 77, 107	2 (0%)
2	D	252/263 (95%)	0.23	5 (1%)	65	69	28, 81, 124, 156	6 (2%)
All	All	1411/1432 (98%)	-0.69	5 (0%)	88	91	17, 41, 101, 156	17 (1%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	286	GLN	3.5
2	D	121	LEU	3.2
2	D	148	VAL	2.8
2	D	192	ALA	2.4
2	D	111	PRO	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	C	506	6/6	0.91	0.12	46,56,58,60	0
3	GOL	A	502	6/6	0.92	0.11	51,55,57,61	0
3	GOL	C	503	6/6	0.93	0.12	46,47,52,58	0
3	GOL	A	501	6/6	0.95	0.10	36,39,41,54	0
3	GOL	A	504	6/6	0.95	0.11	37,45,47,49	0
3	GOL	C	507	6/6	0.95	0.11	48,61,69,72	0
3	GOL	C	504	6/6	0.96	0.11	40,45,46,47	0
3	GOL	C	505	6/6	0.97	0.08	38,47,49,53	0
3	GOL	A	503	6/6	0.97	0.10	34,37,42,42	0
3	GOL	C	501	6/6	0.97	0.07	36,41,48,51	0
3	GOL	C	502	6/6	0.98	0.10	40,46,46,48	0
5	FWK	C	509	19/19	0.98	0.06	39,40,55,59	0
6	B12	D	301	91/91	0.98	0.07	47,62,71,76	0
5	FWK	A	506	19/19	0.99	0.04	29,35,40,42	0
4	NH4	A	505	1/1	0.99	0.05	27,27,27,27	0
6	B12	B	301	91/91	0.99	0.05	33,41,49,55	0
4	NH4	C	508	1/1	0.99	0.10	35,35,35,35	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around FWK C 509:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around B12 D 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FWK A 506:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around B12 B 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.