



Full wwPDB NMR Structure Validation Report ⓘ

Mar 9, 2026 – 12:01 PM UTC

PDB ID : 7XRW / pdb_00007xrw
BMRB ID : 36489
Title : Solution structure of T. brucei RAP1
Authors : Yang, X.; Pan, X.H.; Ji, Z.Y.; Wong, K.B.; Zhang, M.J.; Zhao, Y.X.
Deposited on : 2022-05-12

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 72%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	127	

2 Ensemble composition and analysis

This entry contains 10 models. Model 8 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:650-A:725 (76)	0.70	8

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

NmrClust was unable to cluster the ensemble.

Error message: Inconsistent models

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2005 atoms, of which 1005 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Repressor activator protein 1.

Mol	Chain	Residues	Atoms						Trace
1	A	127	Total	C	H	N	O	S	0
			2005	624	1005	184	190	2	

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	635	GLY	-	expression tag	UNP C1JEX5
A	636	PRO	-	expression tag	UNP C1JEX5
A	637	GLY	-	expression tag	UNP C1JEX5
A	638	SER	-	expression tag	UNP C1JEX5

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Repressor activator protein 1




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1


- Molecule 1: Repressor activator protein 1



4.2.2 Score per residue for model 2


- Molecule 1: Repressor activator protein 1






4.2.3 Score per residue for model 3

- Molecule 1: Repressor activator protein 1




4.2.4 Score per residue for model 4

- Molecule 1: Repressor activator protein 1




4.2.5 Score per residue for model 5

- Molecule 1: Repressor activator protein 1




4.2.6 Score per residue for model 6

- Molecule 1: Repressor activator protein 1




4.2.7 Score per residue for model 7

- Molecule 1: Repressor activator protein 1




4.2.8 Score per residue for model 8 (medoid)

- Molecule 1: Repressor activator protein 1




4.2.9 Score per residue for model 9

- Molecule 1: Repressor activator protein 1



4.2.10 Score per residue for model 10

- Molecule 1: Repressor activator protein 1



5 Refinement protocol and experimental data overview

The models were refined using the following method: *DGSA-distance geometry simulated annealing*.

Of the 90 calculated structures, 10 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	structure calculation	
X-PLOR NIH	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1212
Number of shifts mapped to atoms	1212
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	72%

6 Model quality [i](#)

6.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.16±0.01	0±0/635 (0.0± 0.0%)	1.11±0.02	1±1/858 (0.1± 0.1%)
All	All	1.16	0/6350 (0.0%)	1.11	7/8580 (0.1%)

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	679	PHE	N-CA-C	-6.00	104.07	113.02	7	4
1	A	652	GLU	N-CA-C	-5.42	105.98	113.18	10	2
1	A	709	GLU	N-CA-C	-5.18	104.35	111.81	10	1

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	619	622	622	8±2
All	All	6190	6220	6220	80

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:677:LEU:HD21	1:A:679:PHE:CD1	0.80	2.11	7	5
1:A:661:LEU:HD23	1:A:715:LEU:HD12	0.73	1.60	10	2
1:A:677:LEU:HD22	1:A:678:PRO:HD3	0.70	1.62	10	1
1:A:677:LEU:HB3	1:A:678:PRO:CD	0.68	2.17	10	1
1:A:661:LEU:HD11	1:A:719:LEU:HD21	0.62	1.71	10	1
1:A:656:ILE:CG2	1:A:719:LEU:HD23	0.56	2.31	8	6
1:A:651:PHE:CZ	1:A:707:TYR:CG	0.55	2.95	10	7
1:A:677:LEU:HD13	1:A:677:LEU:H	0.50	1.66	8	2
1:A:684:ILE:HD12	1:A:697:PHE:CZ	0.50	2.42	5	1
1:A:677:LEU:HB3	1:A:678:PRO:HD2	0.48	1.82	10	1
1:A:677:LEU:HB2	1:A:678:PRO:HD2	0.47	1.86	6	9
1:A:684:ILE:HD11	1:A:695:CYS:HB3	0.47	1.85	5	2
1:A:678:PRO:HB2	1:A:712:VAL:HG12	0.47	1.85	3	2
1:A:651:PHE:CZ	1:A:707:TYR:CD2	0.47	3.03	2	5
1:A:668:LEU:CD2	1:A:673:LEU:HD22	0.47	2.39	10	2
1:A:656:ILE:HG22	1:A:719:LEU:HD23	0.46	1.86	9	3
1:A:668:LEU:HD12	1:A:668:LEU:C	0.46	2.35	9	1
1:A:677:LEU:O	1:A:678:PRO:C	0.45	2.59	9	2
1:A:673:LEU:HD21	1:A:686:LEU:HD13	0.45	1.88	6	2
1:A:651:PHE:CZ	1:A:723:PRO:HB3	0.45	2.47	5	1
1:A:677:LEU:CD2	1:A:677:LEU:C	0.44	2.90	9	2
1:A:714:ILE:HG21	1:A:719:LEU:HD22	0.44	1.90	2	1
1:A:673:LEU:HD23	1:A:686:LEU:HD13	0.44	1.89	10	1
1:A:677:LEU:CB	1:A:678:PRO:CD	0.44	2.93	10	4
1:A:651:PHE:CE2	1:A:723:PRO:HB3	0.43	2.49	2	1
1:A:677:LEU:HD13	1:A:714:ILE:HB	0.43	1.91	7	2
1:A:677:LEU:CD2	1:A:678:PRO:HD3	0.43	2.40	10	1
1:A:677:LEU:HD23	1:A:677:LEU:O	0.43	2.12	7	1
1:A:679:PHE:CD2	1:A:686:LEU:HD11	0.43	2.49	9	1
1:A:724:VAL:HG23	1:A:724:VAL:O	0.42	2.14	1	1
1:A:668:LEU:HD22	1:A:673:LEU:HD22	0.42	1.92	10	1
1:A:668:LEU:HD21	1:A:693:ILE:HD11	0.42	1.92	1	1
1:A:676:GLN:HB3	1:A:677:LEU:HD13	0.42	1.92	8	1
1:A:651:PHE:CE1	1:A:723:PRO:HB3	0.41	2.51	5	1
1:A:677:LEU:HB2	1:A:678:PRO:CD	0.41	2.46	3	3
1:A:677:LEU:CB	1:A:678:PRO:HD2	0.41	2.46	6	1
1:A:661:LEU:HD23	1:A:717:ARG:HD2	0.40	1.93	4	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	76/127 (60%)	67±1 (88±2%)	6±2 (8±2%)	3±1 (4±1%)	4	30
All	All	760/1270 (60%)	667 (88%)	63 (8%)	30 (4%)	4	30

All 6 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	659	GLU	9
1	A	677	LEU	8
1	A	679	PHE	6
1	A	678	PRO	4
1	A	667	GLY	2
1	A	683	PRO	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	70/109 (64%)	65±1 (93±2%)	5±1 (7±2%)	17	66
All	All	700/1090 (64%)	654 (93%)	46 (7%)	17	66

All 13 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	668	LEU	9
1	A	676	GLN	8
1	A	685	LYS	8
1	A	679	PHE	5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	712	VAL	3
1	A	693	ILE	3
1	A	677	LEU	3
1	A	719	LEU	2
1	A	682	TYR	1
1	A	694	PHE	1
1	A	709	GLU	1
1	A	665	ARG	1
1	A	663	SER	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 72% for the well-defined parts and 69% for the entire structure.

7.1 Chemical shift list 1

File name: `working_cs.cif`

Chemical shift list name: *starch_output*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1212
Number of shifts mapped to atoms	1212
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	2

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	126	-0.11 ± 0.08	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	119	0.03 ± 0.11	None needed (< 0.5 ppm)
$^{13}\text{C}'$	0	—	None (insufficient data)
^{15}N	111	-0.13 ± 0.28	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 72%, i.e. 778 atoms were assigned a chemical shift out of a possible 1082. 0 out of 11 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	288/370 (78%)	146/149 (98%)	76/152 (50%)	66/69 (96%)
Sidechain	486/624 (78%)	322/402 (80%)	164/193 (85%)	0/29 (0%)

Continued on next page...

Continued from previous page...

	Total	¹H	¹³C	¹⁵N
Aromatic	4/88 (5%)	4/43 (9%)	0/45 (0%)	0/0 (—%)
Overall	778/1082 (72%)	472/594 (79%)	240/390 (62%)	66/98 (67%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 69%, i.e. 1212 atoms were assigned a chemical shift out of a possible 1748. 0 out of 17 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Backbone	473/621 (76%)	236/251 (94%)	126/254 (50%)	111/116 (96%)
Sidechain	735/1032 (71%)	480/664 (72%)	255/313 (81%)	0/55 (0%)
Aromatic	4/95 (4%)	4/47 (9%)	0/47 (0%)	0/1 (0%)
Overall	1212/1748 (69%)	720/962 (75%)	381/614 (62%)	111/172 (65%)

7.1.4 Statistically unusual chemical shifts ⓘ

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	723	PRO	HB2	-0.39	0.37 – 3.78	-7.2
1	A	723	PRO	HB3	0.05	0.25 – 3.76	-5.6

7.1.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:

