



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 05:09 PM UTC

PDB ID : 1XSP / pdb_00001xsp
Title : Crystal Structure of human DNA polymerase lambda in complex with nicked DNA and pyrophosphate
Authors : Garcia-Diaz, M.; Bebenek, K.; Krahn, J.M.; Kunkel, T.A.; Pedersen, L.C.
Deposited on : 2004-10-19
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	T	11	 82% 18%
2	P	7	 71% 29%
3	D	4	 50% 50%
4	A	335	 9% 75% 21% ..

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 3363 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(*CP*GP*GP*CP*CP*GP*TP*AP*CP*TP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	T	11	Total	C	N	O	P	0	0	0
			222	106	41	65	10			

- Molecule 2 is a DNA chain called 5'-D(*CP*AP*GP*TP*AP*CP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	P	7	Total	C	N	O	P	0	0	0
			141	68	28	39	6			

- Molecule 3 is a DNA chain called 5'-D(P*GP*CP*CP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	4	Total	C	N	O	P	0	0	0
			83	38	16	25	4			

- Molecule 4 is a protein called DNA polymerase lambda.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	A	327	Total	C	N	O	S	0	5	0
			2570	1610	470	479	11			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	241	MET	-	initiating methionine	UNP Q9UGP5
A	543	ALA	CYS	engineered mutation	UNP Q9UGP5

- Molecule 5 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	D	1	Total Na 1 1	0	0
5	A	3	Total Na 3 3	0	0

- Molecule 6 is PYROPHOSPHATE (CCD ID: PPV) (formula: $\text{H}_4\text{O}_7\text{P}_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total O P 9 7 2	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	T	66	Total O 66 66	0	0
7	P	28	Total O 28 28	0	0
7	D	11	Total O 11 11	0	0
7	A	229	Total O 229 229	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5'-D(*CP*GP*GP*CP*CP*GP*TP*AP*CP*TP*G)-3'





- Molecule 2: 5'-D(*CP*AP*GP*TP*AP*CP*G)-3'



- Molecule 3: 5'-D(P*GP*CP*CP*G)-3'



- Molecule 4: DNA polymerase lambda



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	55.84Å 62.76Å 140.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.72 – 2.20 41.72 – 2.20	Depositor EDS
% Data completeness (in resolution range)	94.0 (41.72-2.20) 94.0 (41.72-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.55 (at 2.20Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.206 , 0.242 0.213 , 0.246	Depositor DCC
R_{free} test set	1230 reflections (4.80%)	wwPDB-VP
Wilson B-factor (Å ²)	35.1	Xtriage
Anisotropy	0.403	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 48.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3363	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, PPV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	T	0.28	0/248	0.67	0/381
2	P	0.29	0/158	0.57	0/242
3	D	0.53	0/92	0.51	0/138
4	A	0.51	0/2621	0.95	11/3540 (0.3%)
All	All	0.49	0/3119	0.90	11/4301 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
4	A	0	1

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
4	A	453	GLN	OE1-CD-NE2	-9.83	112.77	122.60
4	A	449	ASP	O-C-N	-8.94	109.38	122.43
4	A	298	GLU	CA-C-N	7.31	135.50	121.54
4	A	298	GLU	C-N-CA	7.31	135.50	121.54
4	A	545	VAL	N-CA-C	-6.99	107.07	113.71
4	A	453	GLN	CG-CD-NE2	6.68	126.42	116.40
4	A	298	GLU	O-C-N	6.40	129.45	122.15
4	A	273	LYS	N-CA-C	6.13	117.96	111.28
4	A	290	HIS	N-CA-C	6.00	120.61	112.88
4	A	553	THR	CA-C-N	5.08	124.87	119.28
4	A	553	THR	C-N-CA	5.08	124.87	119.28

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
4	A	449	ASP	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	T	222	0	125	1	0
2	P	141	0	80	1	0
3	D	83	0	45	1	0
4	A	2570	0	2524	54	0
5	A	3	0	0	0	0
5	D	1	0	0	0	0
6	A	9	0	0	0	0
7	A	229	0	0	7	0
7	D	11	0	0	0	0
7	P	28	0	0	0	0
7	T	66	0	0	0	0
All	All	3363	0	2774	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:256:ILE:HD13	4:A:313:ILE:HG23	1.59	0.84
4:A:417:SER:HB2	4:A:422[B]:LYS:HD3	1.60	0.84
4:A:291:LYS:HE3	4:A:298:GLU:OE2	1.88	0.74
4:A:488:ARG:HD2	7:A:731:HOH:O	1.91	0.70
4:A:396:GLU:HG3	4:A:414:ALA:HB2	1.72	0.70
4:A:256:ILE:HG12	4:A:316:ILE:HG21	1.75	0.69
4:A:438:ARG:O	4:A:441:ARG:HG3	1.96	0.65
4:A:387:MET:HE1	4:A:428:VAL:HG23	1.79	0.63
4:A:320:GLY:HA2	7:A:692:HOH:O	1.98	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:302:ILE:HB	4:A:305:ILE:HD12	1.80	0.63
4:A:537:VAL:HB	4:A:546:GLY:HA3	1.82	0.60
4:A:519:LEU:HD23	4:A:565:LEU:HD11	1.83	0.60
4:A:289:PHE:CE2	4:A:291:LYS:O	2.56	0.58
4:A:289:PHE:HE2	4:A:291:LYS:O	1.86	0.58
4:A:259:LYS:NZ	7:A:778:HOH:O	2.37	0.57
4:A:351:MET:O	4:A:355:GLN:HG3	2.05	0.56
4:A:256:ILE:HG12	4:A:316:ILE:CG2	2.35	0.56
4:A:397:GLN:O	4:A:401:LYS:HG3	2.05	0.56
4:A:539:ASN:HD21	4:A:543:ALA:HB3	1.70	0.55
4:A:449:ASP:O	4:A:453:GLN:HG3	2.07	0.54
4:A:417:SER:CB	4:A:422[B]:LYS:HD3	2.37	0.53
4:A:521:LYS:HE2	4:A:538:ARG:NE	2.25	0.52
4:A:298:GLU:O	4:A:301:SER:HB2	2.10	0.51
4:A:346[B]:THR:HG23	7:A:604:HOH:O	2.10	0.50
4:A:261:GLU:HG2	4:A:283:ILE:HD13	1.93	0.49
4:A:302:ILE:HD12	4:A:305:ILE:HD12	1.94	0.49
2:P:6:DC:H2'	2:P:7:DG:C8	2.48	0.49
4:A:253:ASN:OD1	4:A:292:PRO:HA	2.13	0.49
4:A:431:LEU:HD12	4:A:432:ILE:N	2.29	0.48
4:A:256:ILE:HG23	4:A:316:ILE:HD12	1.96	0.48
4:A:434:HIS:O	4:A:496:TYR:HB2	2.14	0.48
4:A:521:LYS:HE2	4:A:538:ARG:HE	1.78	0.47
4:A:390:GLU:HB3	7:A:681:HOH:O	2.14	0.47
4:A:539:ASN:ND2	4:A:543:ALA:HB3	2.30	0.47
4:A:285:ALA:O	4:A:289:PHE:HB2	2.15	0.47
4:A:513:ASN:O	4:A:517:ARG:HG3	2.15	0.46
4:A:519:LEU:CD2	4:A:565:LEU:HD21	2.46	0.46
3:D:3:DC:H2''	3:D:4:DG:C8	2.50	0.46
4:A:453:GLN:C	4:A:455:GLY:H	2.23	0.45
4:A:332:VAL:HB	4:A:333:PRO:HD3	1.98	0.45
4:A:436:ASP:OD1	4:A:437:GLY:N	2.49	0.45
4:A:538:ARG:HD3	4:A:542:GLY:O	2.16	0.45
4:A:261:GLU:CG	4:A:283:ILE:HD13	2.47	0.44
4:A:427:ASP:C	4:A:427:ASP:OD1	2.61	0.43
4:A:279:TYR:O	4:A:283:ILE:HG13	2.17	0.43
4:A:565:LEU:O	7:A:766:HOH:O	2.22	0.43
4:A:311:GLU:O	4:A:314:ILE:HB	2.19	0.42
4:A:501:CYS:SG	4:A:531:ALA:HA	2.60	0.42
4:A:505:TYR:CG	4:A:529:GLU:HB3	2.55	0.42
4:A:263:LEU:HG	4:A:267:TYR:CE2	2.54	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:431:LEU:HD11	4:A:494:VAL:HG22	2.02	0.41
4:A:568:ARG:HD2	7:A:769:HOH:O	2.19	0.41
4:A:254:LEU:O	4:A:258:GLU:HB2	2.21	0.41
4:A:453:GLN:C	4:A:455:GLY:N	2.79	0.41
4:A:257:THR:O	4:A:261:GLU:HG3	2.21	0.41
4:A:259:LYS:HB3	4:A:325:LEU:HD11	2.04	0.40
1:T:8:DA:H2''	1:T:9:DC:O5'	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
4	A	330/335 (98%)	315 (96%)	14 (4%)	1 (0%)	36 42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	A	407	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
4	A	267/280 (95%)	266 (100%)	1 (0%)	84 92

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	A	506	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
4	A	270	GLN
4	A	355	GLN
4	A	400	GLN
4	A	404	GLN
4	A	470	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	PPV	A	579	5	6,8,8	1.02	1 (16%)	12,13,13	1.31	1 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PPV	A	579	5	-	0/6/6/6	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	579	PPV	P2-O12	-2.17	1.46	1.54

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	579	PPV	O32-P2-OPP	-2.92	94.86	104.64

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	T	11/11 (100%)	-1.04	0 100 100	25, 27, 33, 40	0
2	P	7/7 (100%)	-1.22	0 100 100	22, 22, 25, 27	0
3	D	4/4 (100%)	-0.19	0 100 100	35, 40, 40, 43	0
4	A	327/335 (97%)	0.40	30 (9%) 14 12	13, 38, 88, 98	5 (1%)
All	All	349/357 (97%)	0.32	30 (8%) 16 14	13, 38, 87, 98	5 (1%)

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	A	293	VAL	4.1
4	A	575	TRP	4.0
4	A	249	ALA	3.9
4	A	250	THR	3.4
4	A	511[A]	HIS	3.2
4	A	286	LEU	3.2
4	A	299	ALA	3.2
4	A	296	TYR	3.0
4	A	254	LEU	2.8
4	A	297	GLN	2.8
4	A	321	HIS	2.7
4	A	292	PRO	2.7
4	A	257	THR	2.6
4	A	253	ASN	2.6
4	A	290	HIS	2.5
4	A	289	PHE	2.4
4	A	251	ASN	2.4
4	A	316	ILE	2.4
4	A	322	LEU	2.4
4	A	319	SER	2.4
4	A	437	GLY	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	A	252	HIS	2.3
4	A	294	THR	2.3
4	A	320	GLY	2.3
4	A	300	CYS	2.2
4	A	439	SER	2.2
4	A	256	ILE	2.2
4	A	317	LEU	2.2
4	A	255	HIS	2.1
4	A	411	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	NA	D	503	1/1	0.70	0.37	79,79,79,79	0
5	NA	A	578	1/1	0.87	0.10	43,43,43,43	0
5	NA	A	577	1/1	0.92	0.07	29,29,29,29	0
5	NA	A	576	1/1	0.92	0.05	28,28,28,28	0
6	PPV	A	579	9/9	0.92	0.10	32,40,49,49	0

6.5 Other polymers [i](#)

There are no such residues in this entry.