



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 14, 2026 – 11:26 AM UTC

PDB ID : 4XSZ / pdb_00004xsz
Title : Crystal structure of CBR 9393 bound to Escherichia coli RNA polymerase holoenzyme
Authors : Bae, B.; Darst, S.A.
Deposited on : 2015-01-22
Resolution : 3.68 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 3.68 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1263 (3.80-3.56)
Clashscore	190562	1305 (3.80-3.56)
Ramachandran outliers	187476	1259 (3.80-3.56)
Sidechain outliers	187428	1256 (3.80-3.56)
RSRZ outliers	180081	1262 (3.80-3.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	239	
1	B	239	
1	G	239	
1	H	239	
2	C	1342	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	I	1342	65% 31%
3	D	1407	53% 26% 17%
3	J	1407	56% 27% 5% 12%
4	E	91	74% 19% 5%
4	K	91	64% 20% 13%
5	F	522	59% 27% 10%
5	L	522	61% 25% 10%

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 55744 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	224	Total	C	N	O	S	0	0	0
			1730	1076	308	340	6			
1	B	220	Total	C	N	O	S	0	0	0
			1687	1053	298	330	6			
1	G	228	Total	C	N	O	S	0	0	0
			1750	1088	312	344	6			
1	H	217	Total	C	N	O	S	0	0	0
			1667	1041	293	327	6			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	235	GLU	-	expression tag	UNP A7ZSI4
A	236	VAL	-	expression tag	UNP A7ZSI4
A	237	LEU	-	expression tag	UNP A7ZSI4
A	238	PHE	-	expression tag	UNP A7ZSI4
A	239	GLN	-	expression tag	UNP A7ZSI4
B	235	GLU	-	expression tag	UNP A7ZSI4
B	236	VAL	-	expression tag	UNP A7ZSI4
B	237	LEU	-	expression tag	UNP A7ZSI4
B	238	PHE	-	expression tag	UNP A7ZSI4
B	239	GLN	-	expression tag	UNP A7ZSI4
G	235	GLU	-	expression tag	UNP A7ZSI4
G	236	VAL	-	expression tag	UNP A7ZSI4
G	237	LEU	-	expression tag	UNP A7ZSI4
G	238	PHE	-	expression tag	UNP A7ZSI4
G	239	GLN	-	expression tag	UNP A7ZSI4
H	235	GLU	-	expression tag	UNP A7ZSI4
H	236	VAL	-	expression tag	UNP A7ZSI4
H	237	LEU	-	expression tag	UNP A7ZSI4
H	238	PHE	-	expression tag	UNP A7ZSI4
H	239	GLN	-	expression tag	UNP A7ZSI4

- Molecule 2 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	1340	Total	C	N	O	S	0	0	0
			10570	6631	1841	2055	43			
2	I	1340	Total	C	N	O	S	0	0	0
			10566	6629	1840	2054	43			

- Molecule 3 is a protein called DNA-directed RNA polymerase subunit beta'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	1166	Total	C	N	O	S	0	0	0
			9107	5723	1634	1704	46			
3	J	1236	Total	C	N	O	S	0	0	0
			9638	6058	1726	1807	47			

- Molecule 4 is a protein called DNA-directed RNA polymerase subunit omega.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	E	89	Total	C	N	O	S	0	0	0
			691	421	129	140	1			
4	K	79	Total	C	N	O	S	0	0	0
			627	382	118	126	1			

- Molecule 5 is a protein called RNA polymerase sigma factor RpoD.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	F	470	Total	C	N	O	S	0	0	0
			3822	2394	680	725	23			
5	L	469	Total	C	N	O	S	0	0	0
			3821	2393	679	726	23			

- Molecule 6 is 4-[3-(4-fluorophenyl)-1H-pyrazol-4-yl]-N-[2-(piperazin-1-yl)ethyl]-2-(trifluoromethyl)aniline (CCD ID: 42U) (formula: C₂₂H₂₃F₄N₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	C	1	Total 31	C 22	F 4	N 5	0	0
6	I	1	Total 31	C 22	F 4	N 5	0	0

- Molecule 7 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	D	1	Total Mg 1 1	0	0
7	J	1	Total Mg 1 1	0	0


- Molecule 8 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	D	2	Total Zn 2 2	0	0
8	J	2	Total Zn 2 2	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA-directed RNA polymerase subunit alpha




- Molecule 1: DNA-directed RNA polymerase subunit alpha




- Molecule 1: DNA-directed RNA polymerase subunit alpha






• Molecule 2: DNA-directed RNA polymerase subunit beta


Chain I: 65% 31%






• Molecule 3: DNA-directed RNA polymerase subunit beta'


Chain D: 53% 26% 17%







• Molecule 3: DNA-directed RNA polymerase subunit beta'

Chain J: 56% 27% 5% 12%






• Molecule 4: DNA-directed RNA polymerase subunit omega




• Molecule 4: DNA-directed RNA polymerase subunit omega



• Molecule 5: RNA polymerase sigma factor RpoD



● Molecule 5: RNA polymerase sigma factor RpoD



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	184.44Å 206.33Å 309.61Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.93 – 3.68 39.93 – 3.68	Depositor EDS
% Data completeness (in resolution range)	88.5 (39.93-3.68) 88.4 (39.93-3.68)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.52 (at 3.66Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.240 , 0.286 0.262 , 0.302	Depositor DCC
R_{free} test set	5687 reflections (4.06%)	wwPDB-VP
Wilson B-factor (Å ²)	139.9	Xtriage
Anisotropy	0.373	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 58.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	55744	wwPDB-VP
Average B, all atoms (Å ²)	97.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: 42U, MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.36	0/1751	0.87	2/2373 (0.1%)
1	B	0.38	0/1707	0.87	1/2314 (0.0%)
1	G	0.37	0/1771	0.91	4/2401 (0.2%)
1	H	0.38	0/1686	0.86	1/2285 (0.0%)
2	C	0.36	1/10739 (0.0%)	0.87	20/14489 (0.1%)
2	I	0.36	0/10735	0.85	19/14484 (0.1%)
3	D	0.36	0/9246	0.87	16/12478 (0.1%)
3	J	0.36	0/9785	0.87	23/13206 (0.2%)
4	E	0.38	0/693	0.89	1/935 (0.1%)
4	K	0.39	0/629	0.87	1/847 (0.1%)
5	F	0.39	0/3873	0.88	11/5206 (0.2%)
5	L	0.39	0/3872	0.88	8/5205 (0.2%)
All	All	0.37	1/56487 (0.0%)	0.87	107/76223 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	C	0	3
2	I	0	3
3	D	0	3
3	J	0	3
5	F	0	1
5	L	0	1
All	All	0	14

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	512	SER	CA-C	-5.43	1.46	1.52

The worst 5 of 107 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	109	ALA	CA-C-N	14.55	135.50	119.93
2	C	109	ALA	C-N-CA	14.55	135.50	119.93
1	G	232	VAL	N-CA-C	10.05	123.33	109.46
3	J	858	VAL	CA-C-N	8.63	130.63	119.84
3	J	858	VAL	C-N-CA	8.63	130.63	119.84

There are no chirality outliers.

5 of 14 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	109	ALA	Peptide
2	C	1161	LEU	Peptide
2	C	236	LYS	Peptide
3	D	1168	GLU	Peptide
3	D	1184	ASP	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1730	0	1756	51	0
1	B	1687	0	1700	48	0
1	G	1750	0	1764	47	0
1	H	1667	0	1689	55	0
2	C	10570	0	10582	284	0
2	I	10566	0	10576	264	0
3	D	9107	0	9309	256	0
3	J	9638	0	9853	271	0
4	E	691	0	695	18	0
4	K	627	0	634	10	0
5	F	3822	0	3885	98	0
5	L	3821	0	3884	93	0
6	C	31	0	23	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	I	31	0	23	0	0
7	D	1	0	0	0	0
7	J	1	0	0	0	0
8	D	2	0	0	0	0
8	J	2	0	0	0	0
All	All	55744	0	56373	1359	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 1359 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:D:660:GLU:HB3	3:D:685:ILE:HD12	1.50	0.94
3:J:660:GLU:HB3	3:J:685:ILE:HD12	1.51	0.93
1:G:45:ARG:NH1	1:H:34:GLY:O	2.11	0.83
3:D:1372:ARG:HH21	3:J:854:ALA:HB3	1.42	0.83
2:C:10:ARG:HD3	2:C:1181:PRO:HG2	1.62	0.81

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	222/239 (93%)	194 (87%)	26 (12%)	2 (1%)	14	45
1	B	216/239 (90%)	192 (89%)	24 (11%)	0	100	100
1	G	226/239 (95%)	196 (87%)	28 (12%)	2 (1%)	14	45
1	H	213/239 (89%)	190 (89%)	23 (11%)	0	100	100
2	C	1338/1342 (100%)	1231 (92%)	102 (8%)	5 (0%)	30	60

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	I	1338/1342 (100%)	1230 (92%)	103 (8%)	5 (0%)	30	60
3	D	1162/1407 (83%)	1077 (93%)	81 (7%)	4 (0%)	36	65
3	J	1230/1407 (87%)	1145 (93%)	81 (7%)	4 (0%)	36	65
4	E	87/91 (96%)	82 (94%)	5 (6%)	0	100	100
4	K	77/91 (85%)	74 (96%)	3 (4%)	0	100	100
5	F	464/522 (89%)	429 (92%)	35 (8%)	0	100	100
5	L	463/522 (89%)	427 (92%)	36 (8%)	0	100	100
All	All	7036/7680 (92%)	6467 (92%)	547 (8%)	22 (0%)	36	65

5 of 22 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	1159	VAL
2	I	1159	VAL
2	C	170	VAL
2	I	170	VAL
3	J	340	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	191/206 (93%)	170 (89%)	21 (11%)	6	26
1	B	184/206 (89%)	163 (89%)	21 (11%)	5	24
1	G	191/206 (93%)	173 (91%)	18 (9%)	8	31
1	H	183/206 (89%)	164 (90%)	19 (10%)	7	28
2	C	1155/1157 (100%)	1027 (89%)	128 (11%)	6	25
2	I	1154/1157 (100%)	1028 (89%)	126 (11%)	6	26
3	D	975/1168 (84%)	859 (88%)	116 (12%)	5	23
3	J	1036/1168 (89%)	913 (88%)	123 (12%)	5	23
4	E	72/75 (96%)	64 (89%)	8 (11%)	6	25

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	K	67/75 (89%)	61 (91%)	6 (9%)	9	32
5	F	417/462 (90%)	372 (89%)	45 (11%)	6	26
5	L	418/462 (90%)	373 (89%)	45 (11%)	6	26
All	All	6043/6548 (92%)	5367 (89%)	676 (11%)	6	25

5 of 676 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	I	859	GLU
3	J	788	LEU
2	I	1073	LYS
2	I	857	VAL
3	J	292	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 98 such sidechains are listed below:

Mol	Chain	Res	Type
1	H	132	HIS
3	J	206	ASN
2	I	513	GLN
2	I	1136	GLN
3	J	594	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	42U	I	1401	-	33,34,34	1.46	8 (24%)	44,48,48	3.12	21 (47%)
6	42U	C	1401	-	33,34,34	1.49	9 (27%)	44,48,48	3.09	19 (43%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	42U	I	1401	-	-	7/20/28/28	0/4/4/4
6	42U	C	1401	-	-	7/20/28/28	0/4/4/4

The worst 5 of 17 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	C	1401	42U	C13-C23	3.16	1.43	1.40
6	I	1401	42U	C13-C23	2.95	1.43	1.40
6	C	1401	42U	C10-C09	2.73	1.52	1.48
6	C	1401	42U	C25-C23	2.53	1.56	1.50
6	I	1401	42U	C12-C13	2.49	1.43	1.39

The worst 5 of 40 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	I	1401	42U	C12-C13-C23	-9.14	110.85	119.87
6	C	1401	42U	C12-C13-C23	-8.91	111.08	119.87
6	I	1401	42U	C11-C10-C24	-7.38	110.71	119.25
6	C	1401	42U	C11-C10-C24	-7.31	110.79	119.25
6	I	1401	42U	C04-C05-C08	6.24	130.96	120.74

There are no chirality outliers.



5 of 14 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	C	1401	42U	N14-C15-C16-N17
6	I	1401	42U	N14-C15-C16-N17
6	C	1401	42U	C13-C23-C25-F27
6	I	1401	42U	C13-C23-C25-F27
6	C	1401	42U	C13-C23-C25-F28

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	224/239 (93%)	-0.61	0 100 100	54, 87, 128, 162	0
1	B	220/239 (92%)	-0.45	1 (0%) 87 68	71, 115, 149, 168	0
1	G	228/239 (95%)	-0.48	0 100 100	74, 113, 149, 176	0
1	H	217/239 (90%)	-0.39	0 100 100	73, 129, 157, 174	0
2	C	1340/1342 (99%)	-0.58	6 (0%) 88 72	29, 79, 134, 171	0
2	I	1340/1342 (99%)	-0.54	2 (0%) 92 87	32, 96, 144, 171	0
3	D	1166/1407 (82%)	-0.57	6 (0%) 87 68	24, 78, 144, 177	0
3	J	1236/1407 (87%)	-0.55	5 (0%) 88 72	33, 95, 148, 172	0
4	E	89/91 (97%)	-0.70	1 (1%) 78 53	40, 86, 123, 133	0
4	K	79/91 (86%)	-0.64	0 100 100	60, 92, 142, 160	0
5	F	470/522 (90%)	-0.47	0 100 100	45, 121, 158, 182	0
5	L	469/522 (89%)	-0.51	0 100 100	49, 117, 159, 180	0
All	All	7078/7680 (92%)	-0.54	21 (0%) 90 76	24, 96, 149, 182	0

The worst 5 of 21 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	1000	LEU	3.4
2	C	920	VAL	3.2
3	D	499	ILE	2.9
2	C	929	ILE	2.8
3	J	522	GLY	2.7

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)



In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	42U	C	1401	31/31	0.86	0.09	31,71,93,105	0
7	MG	J	1501	1/1	0.86	0.08	75,75,75,75	0
6	42U	I	1401	31/31	0.92	0.07	48,83,117,134	0
7	MG	D	1501	1/1	0.96	0.03	42,42,42,42	0
8	ZN	D	1503	1/1	0.98	0.06	76,76,76,76	0
8	ZN	J	1503	1/1	0.98	0.11	126,126,126,126	0
8	ZN	J	1502	1/1	0.99	0.08	160,160,160,160	0
8	ZN	D	1502	1/1	0.99	0.14	176,176,176,176	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 42U C 1401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 42U I 1401:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.