



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2026 – 11:20 AM UTC

PDB ID : 5XS8 / pdb_00005xs8
Title : Crystal structure of solute-binding protein complexed with unsaturated chondroitin disaccharide with two sulfate groups at C-4 and C-6 positions of GalNAc
Authors : Oiki, S.; Kamochi, R.; Mikami, B.; Murata, K.; Hashimoto, W.
Deposited on : 2017-06-12
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	505	85%8%•6%
1	B	505	88%6%6%
1	C	505	85%8%•6%
1	D	505	87%7%6%
2	E	2	100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	2	 100%
2	G	2	 50%  50%
2	H	2	 100%

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 16765 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Extracellular solute-binding protein family 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	474	Total	C	N	O	S	0	0	0
			3833	2466	633	728	6			
1	B	474	Total	C	N	O	S	0	0	0
			3833	2466	633	728	6			
1	C	474	Total	C	N	O	S	0	0	0
			3833	2466	633	728	6			
1	D	474	Total	C	N	O	S	0	0	0
			3833	2466	633	728	6			

- Molecule 2 is an oligosaccharide called 4-deoxy-alpha-L-threo-hex-4-enopyranuronic acid-(1-3)-2-acetamido-2-deoxy-4,6-di-O-sulfo-beta-D-galactopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	2	Total	C	N	O	S	0	0	0
			34	14	1	17	2			
2	F	2	Total	C	N	O	S	0	0	0
			34	14	1	17	2			
2	G	2	Total	C	N	O	S	0	0	0
			34	14	1	17	2			
2	H	2	Total	C	N	O	S	0	0	0
			34	14	1	17	2			

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total 1	Ca 1	0	0
3	C	1	Total 1	Ca 1	0	0
3	D	1	Total 1	Ca 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	327	Total 327	O 327	0	0
4	B	321	Total 321	O 321	0	0
4	C	326	Total 326	O 326	0	0
4	D	319	Total 319	O 319	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Extracellular solute-binding protein family 1




- Molecule 1: Extracellular solute-binding protein family 1




- Molecule 1: Extracellular solute-binding protein family 1



- Molecule 1: Extracellular solute-binding protein family 1





- Molecule 2: 4-deoxy-alpha-L-threo-hex-4-enopyranuronic acid-(1-3)-2-acetamido-2-deoxy-4,6-di-O-sulfo-beta-D-galactopyranose

Chain E:  100%

8EX1
GCD2

- Molecule 2: 4-deoxy-alpha-L-threo-hex-4-enopyranuronic acid-(1-3)-2-acetamido-2-deoxy-4,6-di-O-sulfo-beta-D-galactopyranose

Chain F:  100%

8EX1
GCD2

- Molecule 2: 4-deoxy-alpha-L-threo-hex-4-enopyranuronic acid-(1-3)-2-acetamido-2-deoxy-4,6-di-O-sulfo-beta-D-galactopyranose

Chain G:  50%  50%

8EX1
GCD2

- Molecule 2: 4-deoxy-alpha-L-threo-hex-4-enopyranuronic acid-(1-3)-2-acetamido-2-deoxy-4,6-di-O-sulfo-beta-D-galactopyranose

Chain H:  100%

8EX1
GCD2

4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	49.74Å 69.21Å 165.97Å 89.99° 90.04° 90.03°	Depositor
Resolution (Å)	40.40 – 1.95 40.40 – 1.95	Depositor EDS
% Data completeness (in resolution range)	97.0 (40.40-1.95) 97.0 (40.40-1.95)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.53 (at 1.95Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.173 , 0.215 0.173 , 0.215	Depositor DCC
R_{free} test set	7826 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	23.2	Xtriage
Anisotropy	0.254	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 28.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.477 for h,-k,-l 0.477 for -h,k,-l 0.477 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	16765	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GCD, CA, 8EX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.06	1/3923 (0.0%)	0.96	4/5308 (0.1%)
1	B	1.06	4/3923 (0.1%)	1.00	4/5308 (0.1%)
1	C	1.06	3/3923 (0.1%)	0.98	7/5308 (0.1%)
1	D	1.05	1/3923 (0.0%)	0.99	4/5308 (0.1%)
All	All	1.06	9/15692 (0.1%)	0.98	19/21232 (0.1%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	141	MET	SD-CE	-8.61	1.58	1.79
1	B	402	GLY	N-CA	7.33	1.56	1.45
1	B	401	LEU	C-N	-6.82	1.24	1.33
1	C	141	MET	SD-CE	-5.99	1.64	1.79
1	C	401	LEU	C-N	-5.94	1.25	1.33
1	C	409	TYR	C-O	-5.77	1.17	1.24
1	B	402	GLY	CA-C	5.70	1.59	1.51
1	D	401	LEU	C-N	-5.69	1.25	1.33
1	B	53	LYS	CD-CE	5.50	1.69	1.52

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	401	LEU	O-C-N	-8.84	110.83	122.59
1	D	401	LEU	O-C-N	-8.46	111.34	122.59
1	C	401	LEU	O-C-N	-8.19	111.69	122.59
1	D	401	LEU	CA-C-N	-8.18	105.42	121.53
1	D	401	LEU	C-N-CA	-8.18	105.42	121.53
1	C	401	LEU	CA-C-N	-7.50	106.75	121.53
1	C	401	LEU	C-N-CA	-7.50	106.75	121.53
1	B	401	LEU	CA-C-N	-7.16	105.49	121.18

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	401	LEU	C-N-CA	-7.16	105.49	121.18
1	B	402	GLY	N-CA-C	6.58	124.78	115.30
1	C	311	GLU	N-CA-CB	-6.10	100.35	110.23
1	A	253	LYS	CB-CG-CD	-5.87	97.80	111.30
1	C	311	GLU	CB-CA-C	5.71	118.95	109.53
1	D	195	LYS	CD-CE-NZ	-5.51	94.26	111.90
1	A	124	GLU	N-CA-CB	-5.33	102.28	110.44
1	C	402	GLY	N-CA-C	5.33	122.71	115.30
1	A	28	LYS	CA-C-N	-5.19	114.19	121.72
1	A	28	LYS	C-N-CA	-5.19	114.19	121.72
1	C	264	ARG	N-CA-C	5.04	121.53	110.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3833	0	3788	32	0
1	B	3833	0	3788	17	0
1	C	3833	0	3788	27	0
1	D	3833	0	3788	19	0
2	E	34	0	5	0	0
2	F	34	0	5	0	0
2	G	34	0	5	1	0
2	H	34	0	5	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	327	0	0	8	0
4	B	321	0	0	2	0
4	C	326	0	0	1	0
4	D	319	0	0	3	0
All	All	16765	0	15172	92	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (92) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:141:MET:SD	4:A:975:HOH:O	2.20	0.98
1:C:95:LEU:HD23	1:C:141:MET:HE2	1.48	0.95
1:A:426:MET:HE2	4:A:823:HOH:O	1.68	0.92
1:B:426:MET:HE2	4:B:882:HOH:O	1.73	0.89
1:A:155:GLN:OE1	1:A:210:LYS:NZ	2.10	0.85
1:A:80:GLU:OE2	1:C:491:LYS:NZ	2.10	0.85
1:B:294:ASN:HD21	1:B:385:GLY:HA2	1.42	0.84
1:A:28:LYS:O	4:A:701:HOH:O	1.95	0.83
1:D:155:GLN:OE1	1:D:210:LYS:NZ	2.12	0.82
1:B:155:GLN:OE1	1:B:210:LYS:NZ	2.15	0.80
1:C:442:LYS:HE2	1:C:446:LYS:NZ	1.97	0.78
1:C:155:GLN:OE1	1:C:210:LYS:NZ	2.16	0.74
1:A:422:TYR:HB3	1:A:426:MET:HE3	1.70	0.74
1:A:166:LYS:HE3	4:A:976:HOH:O	1.89	0.73
1:A:28:LYS:HB2	4:A:701:HOH:O	1.89	0.71
1:B:422:TYR:HB3	1:B:426:MET:HE3	1.72	0.71
1:A:426:MET:HE1	4:A:895:HOH:O	1.90	0.70
1:C:101:GLU:OE2	1:C:442:LYS:HE3	1.92	0.70
1:D:442:LYS:O	1:D:446:LYS:HG2	1.92	0.69
1:A:442:LYS:O	1:A:446:LYS:HG2	1.94	0.67
1:A:206:ASN:HB3	1:A:266:LEU:HD21	1.76	0.66
1:D:144:ASN:HB2	1:D:402:GLY:H	1.59	0.66
1:D:50:LYS:HE2	4:D:983:HOH:O	1.96	0.64
1:D:231:MET:HE3	1:D:233:LYS:HD2	1.81	0.62
1:C:95:LEU:CD2	1:C:141:MET:HE2	2.26	0.62
1:C:144:ASN:HB2	1:C:402:GLY:H	1.65	0.62
1:B:101:GLU:OE2	1:B:442:LYS:HE3	1.99	0.61
1:A:50:LYS:HE2	4:A:984:HOH:O	2.00	0.61
1:C:442:LYS:HE2	1:C:446:LYS:HZ2	1.63	0.61
1:C:442:LYS:HE2	1:C:446:LYS:HZ1	1.65	0.60
1:C:326:LEU:O	1:C:400:ARG:HD3	2.03	0.59
1:A:206:ASN:HB3	1:A:266:LEU:CD2	2.32	0.58
1:B:326:LEU:O	1:B:400:ARG:HD3	2.04	0.56
1:C:210:LYS:HE3	2:G:1:8EX:O1S	2.05	0.56
1:A:422:TYR:HB3	1:A:426:MET:CE	2.36	0.56
1:B:143:PRO:HB3	1:B:325:TYR:CD2	2.40	0.55
1:A:491:LYS:NZ	1:C:80:GLU:OE2	2.40	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:144:ASN:HB2	1:B:402:GLY:H	1.73	0.54
1:C:141:MET:CE	1:C:328:GLY:HA3	2.37	0.54
1:C:143:PRO:HB3	1:C:325:TYR:CD2	2.44	0.53
1:A:445:SER:O	1:A:449:VAL:HG13	2.08	0.52
1:D:326:LEU:O	1:D:400:ARG:HD3	2.09	0.52
1:A:326:LEU:O	1:A:400:ARG:HD3	2.10	0.52
1:A:266:LEU:H	1:A:266:LEU:HD22	1.75	0.51
1:A:82:LYS:HD3	1:A:334:ASP:CG	2.35	0.51
1:B:153:VAL:HG22	1:B:431:VAL:HG21	1.93	0.51
1:D:101:GLU:OE2	1:D:442:LYS:HE2	2.11	0.50
1:D:444:LYS:HE3	4:D:725:HOH:O	2.12	0.50
1:D:494:GLN:O	1:D:498:GLU:HG3	2.12	0.50
1:D:442:LYS:NZ	1:D:446:LYS:HE3	2.28	0.49
1:B:246:LYS:HE2	1:B:474:ASP:OD2	2.13	0.48
1:C:242:LYS:NZ	1:C:474:ASP:OD1	2.45	0.48
1:A:316:ALA:HB1	4:A:785:HOH:O	2.13	0.48
1:D:316:ALA:HB1	4:D:747:HOH:O	2.13	0.47
1:A:494:GLN:O	1:A:498:GLU:HG3	2.15	0.47
1:D:143:PRO:HB3	1:D:325:TYR:CD2	2.50	0.47
1:A:143:PRO:HB3	1:A:325:TYR:CD2	2.49	0.46
1:B:227:GLU:HG2	1:B:228:GLU:N	2.31	0.46
1:C:92:SER:OG	1:C:141:MET:CE	2.63	0.46
1:B:52:GLU:OE2	1:B:58:LYS:HD3	2.16	0.46
1:A:242:LYS:HD2	1:A:477:TRP:CG	2.50	0.45
1:D:169:LEU:HD23	1:D:169:LEU:HA	1.66	0.45
1:B:72:GLN:HG3	1:D:454:ALA:CB	2.47	0.44
1:B:426:MET:HG2	4:B:1010:HOH:O	2.16	0.44
1:D:257:ILE:CG2	1:D:261:ILE:HD13	2.47	0.44
1:C:131:ASP:HB3	1:C:325:TYR:CE1	2.53	0.43
1:C:111:ILE:HA	1:C:115:ALA:HB3	1.99	0.43
1:C:227:GLU:HG2	1:C:228:GLU:N	2.31	0.43
1:D:111:ILE:HA	1:D:115:ALA:HB3	2.00	0.43
1:B:49:VAL:O	1:B:53:LYS:HE3	2.18	0.43
1:A:122:PHE:HB3	1:A:129:LYS:HD3	2.01	0.43
1:C:163:TRP:CG	1:C:190:PRO:HB2	2.54	0.43
1:A:365:GLU:CD	1:A:365:GLU:N	2.78	0.42
1:C:494:GLN:O	1:C:498:GLU:HG3	2.19	0.42
1:D:47:VAL:HG13	1:D:356:LEU:HG	2.00	0.42
1:A:47:VAL:HG13	1:A:356:LEU:HG	2.01	0.42
1:C:85:ASP:HB3	1:C:341:LEU:HD11	2.02	0.42
1:C:223:PRO:HA	1:C:236:PRO:HG3	2.02	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:141:MET:HB2	1:A:141:MET:HE3	1.70	0.42
1:A:257:ILE:CG2	1:A:261:ILE:HD13	2.49	0.42
1:A:366:TYR:CD2	1:A:373:PRO:HB3	2.54	0.42
1:C:298:LYS:HB2	1:C:298:LYS:HE2	1.80	0.42
1:C:308:LEU:O	1:C:309:PRO:C	2.63	0.41
1:D:183:LYS:HE3	1:D:254:GLU:OE2	2.20	0.41
1:D:28:LYS:O	1:D:58:LYS:HB3	2.19	0.41
1:B:256:LEU:N	1:B:256:LEU:HD23	2.36	0.41
1:A:227:GLU:HG2	1:A:228:GLU:N	2.36	0.41
1:A:266:LEU:HD22	1:A:266:LEU:N	2.34	0.41
1:C:153:VAL:HG13	4:C:847:HOH:O	2.21	0.40
1:C:242:LYS:HD2	1:C:477:TRP:CG	2.56	0.40
1:A:111:ILE:HA	1:A:115:ALA:HB3	2.03	0.40
1:B:143:PRO:HB3	1:B:325:TYR:CG	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	472/505 (94%)	462 (98%)	10 (2%)	0	100	100
1	B	472/505 (94%)	464 (98%)	8 (2%)	0	100	100
1	C	472/505 (94%)	460 (98%)	12 (2%)	0	100	100
1	D	472/505 (94%)	462 (98%)	10 (2%)	0	100	100
All	All	1888/2020 (94%)	1848 (98%)	40 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	408/433 (94%)	402 (98%)	6 (2%)	57	54
1	B	408/433 (94%)	404 (99%)	4 (1%)	68	67
1	C	408/433 (94%)	403 (99%)	5 (1%)	63	61
1	D	408/433 (94%)	403 (99%)	5 (1%)	63	61
All	All	1632/1732 (94%)	1612 (99%)	20 (1%)	63	61

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	82	LYS
1	A	141	MET
1	A	261	ILE
1	A	380	LEU
1	A	409	TYR
1	A	449	VAL
1	B	141	MET
1	B	261	ILE
1	B	409	TYR
1	B	487	LEU
1	C	95	LEU
1	C	261	ILE
1	C	409	TYR
1	C	468	LEU
1	C	487	LEU
1	D	78	LEU
1	D	95	LEU
1	D	141	MET
1	D	261	ILE
1	D	409	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	405	GLN
1	A	494	GLN
1	B	161	GLN
1	B	294	ASN
1	B	313	ASN
1	B	405	GLN
1	B	464	GLN
1	B	494	GLN
1	C	65	GLN
1	C	113	GLN
1	C	398	GLN
1	C	405	GLN
1	C	464	GLN
1	D	313	ASN
1	D	398	GLN
1	D	405	GLN
1	D	464	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	8EX	E	1	2	23,23,23	1.56	5 (21%)	31,35,35	2.67	11 (35%)
2	GCD	E	2	2	10,11,12	2.07	3 (30%)	12,15,17	1.38	3 (25%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	8EX	F	1	2	23,23,23	2.10	8 (34%)	31,35,35	2.90	12 (38%)
2	GCD	F	2	2	10,11,12	2.08	1 (10%)	12,15,17	1.63	1 (8%)
2	8EX	G	1	2	23,23,23	2.29	8 (34%)	31,35,35	2.90	15 (48%)
2	GCD	G	2	2	10,11,12	2.24	1 (10%)	12,15,17	1.59	2 (16%)
2	8EX	H	1	2	23,23,23	1.88	7 (30%)	31,35,35	2.67	11 (35%)
2	GCD	H	2	2	10,11,12	2.45	3 (30%)	12,15,17	1.43	2 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	8EX	E	1	2	-	4/15/35/35	0/1/1/1
2	GCD	E	2	2	-	0/4/17/20	0/1/1/1
2	8EX	F	1	2	-	3/15/35/35	0/1/1/1
2	GCD	F	2	2	-	0/4/17/20	0/1/1/1
2	8EX	G	1	2	-	7/15/35/35	0/1/1/1
2	GCD	G	2	2	-	0/4/17/20	0/1/1/1
2	8EX	H	1	2	-	5/15/35/35	0/1/1/1
2	GCD	H	2	2	-	0/4/17/20	0/1/1/1

All (36) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	2	GCD	O5-C5	6.17	1.45	1.37
2	G	2	GCD	O5-C5	6.09	1.45	1.37
2	F	2	GCD	O5-C5	5.80	1.45	1.37
2	G	1	8EX	C8-C7	-5.67	1.38	1.50
2	F	1	8EX	O1S-S1	5.48	1.69	1.45
2	E	2	GCD	O5-C5	5.16	1.44	1.37
2	H	1	8EX	C8-C7	4.49	1.59	1.50
2	F	1	8EX	C8-C7	4.38	1.59	1.50
2	G	1	8EX	O7-C7	4.04	1.32	1.23
2	G	1	8EX	O1S-S1	3.96	1.62	1.45
2	G	1	8EX	C2-N2	3.58	1.51	1.45
2	G	1	8EX	O3S-S1	3.51	1.60	1.45
2	E	1	8EX	C3-C2	-3.50	1.46	1.53
2	E	1	8EX	O3S-S1	3.35	1.59	1.45
2	F	1	8EX	C2-N2	3.20	1.50	1.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	2	GCD	C3-C4	3.15	1.54	1.50
2	H	1	8EX	O3S-S1	3.01	1.58	1.45
2	H	1	8EX	C2-N2	2.98	1.50	1.45
2	F	1	8EX	O3S-S1	2.82	1.57	1.45
2	G	1	8EX	O6-C6	-2.70	1.36	1.46
2	F	1	8EX	C3-C2	-2.68	1.48	1.53
2	E	2	GCD	C3-C4	2.67	1.53	1.50
2	F	1	8EX	O6-S1	-2.65	1.49	1.56
2	H	1	8EX	C3-C2	-2.48	1.48	1.53
2	H	2	GCD	O5-C1	-2.42	1.41	1.45
2	E	1	8EX	C8-C7	2.31	1.55	1.50
2	H	1	8EX	O2S-S1	2.31	1.64	1.50
2	G	1	8EX	O4-C4	-2.31	1.41	1.46
2	H	1	8EX	O6-C6	-2.31	1.38	1.46
2	F	1	8EX	O5-C1	2.26	1.48	1.42
2	H	1	8EX	O1S-S1	2.26	1.55	1.45
2	E	2	GCD	O6B-C6	-2.18	1.24	1.30
2	G	1	8EX	C3-C2	-2.14	1.49	1.53
2	E	1	8EX	O2S-S1	2.12	1.63	1.50
2	E	1	8EX	O6-C6	-2.09	1.38	1.46
2	F	1	8EX	O6-C6	-2.04	1.38	1.46

All (57) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	1	8EX	O1-C1-C2	-8.31	91.96	109.22
2	F	1	8EX	O1-C1-C2	-7.83	92.96	109.22
2	H	1	8EX	C1-O5-C5	-7.13	99.87	113.65
2	G	1	8EX	O1-C1-C2	-6.89	94.91	109.22
2	F	1	8EX	O4S-S2-O6S	-6.74	84.99	108.56
2	H	1	8EX	O1-C1-C2	-6.69	95.31	109.22
2	G	1	8EX	C1-O5-C5	-6.44	101.19	113.65
2	G	1	8EX	O2S-S1-O1S	5.23	126.84	108.56
2	E	1	8EX	C1-O5-C5	-5.13	103.73	113.65
2	G	1	8EX	O1-C1-O5	-4.97	95.64	110.41
2	F	1	8EX	O4S-S2-O4	-4.97	94.94	106.37
2	E	1	8EX	O1-C1-O5	-4.97	95.65	110.41
2	G	1	8EX	C1-C2-N2	-4.85	105.11	110.73
2	F	1	8EX	C1-O5-C5	-4.83	104.31	113.65
2	H	1	8EX	O2S-S1-O1S	4.82	125.43	108.56
2	H	1	8EX	O1-C1-O5	-4.70	96.46	110.41
2	F	2	GCD	O3-C3-C2	-4.56	101.31	109.44

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	1	8EX	C1-C2-N2	-4.36	105.68	110.73
2	G	1	8EX	O6-S1-O3S	4.14	119.56	106.92
2	F	1	8EX	C1-C2-C3	-4.12	104.92	110.54
2	H	1	8EX	C4-O4-S2	3.87	128.31	119.04
2	F	1	8EX	O2S-S1-O1S	3.79	121.81	108.56
2	F	1	8EX	O1-C1-O5	-3.64	99.59	110.41
2	F	1	8EX	C1-C2-N2	-3.53	106.64	110.73
2	G	2	GCD	O3-C3-C2	-3.50	103.20	109.44
2	H	1	8EX	O6-S1-O3S	3.48	117.53	106.92
2	H	1	8EX	C1-C2-C3	-3.37	105.94	110.54
2	H	1	8EX	O3S-S1-O1S	-3.31	99.47	112.24
2	G	1	8EX	O5-C1-C2	-3.29	106.21	109.52
2	E	1	8EX	O6-S1-O3S	3.28	116.92	106.92
2	E	1	8EX	O3S-S1-O1S	-3.27	99.63	112.24
2	E	1	8EX	O2S-S1-O1S	3.23	119.85	108.56
2	G	1	8EX	C1-C2-C3	-3.20	106.17	110.54
2	H	1	8EX	O2S-S1-O6	-3.08	99.29	106.37
2	G	1	8EX	O6-S1-O1S	-3.01	97.74	106.92
2	H	2	GCD	O3-C3-C2	-2.94	104.20	109.44
2	F	1	8EX	O6S-S2-O5S	2.84	123.21	112.24
2	G	1	8EX	C6-C5-C4	-2.75	106.72	113.35
2	F	1	8EX	O2S-S1-O3S	-2.72	99.02	108.56
2	E	1	8EX	C4-O4-S2	2.69	125.48	119.04
2	E	1	8EX	O5-C1-C2	-2.59	106.92	109.52
2	E	2	GCD	O3-C3-C2	-2.56	104.87	109.44
2	H	1	8EX	C6-C5-C4	-2.45	107.45	113.35
2	G	1	8EX	C2-N2-C7	2.44	128.83	123.11
2	G	1	8EX	O4S-S2-O5S	2.43	117.06	108.56
2	G	2	GCD	O5-C5-C6	2.42	116.29	111.85
2	H	2	GCD	O6B-C6-C5	2.27	119.21	114.06
2	G	1	8EX	O2S-S1-O3S	-2.27	100.60	108.56
2	F	1	8EX	O5-C1-C2	-2.23	107.28	109.52
2	E	1	8EX	C1-C2-C3	-2.20	107.54	110.54
2	E	2	GCD	O6A-C6-C5	-2.15	115.31	120.57
2	E	2	GCD	O6B-C6-C5	2.11	118.85	114.06
2	G	1	8EX	O2S-S1-O6	-2.10	101.55	106.37
2	H	1	8EX	C1-C2-N2	-2.07	108.32	110.73
2	G	1	8EX	C8-C7-N2	-2.05	112.72	116.12
2	F	1	8EX	O6-S1-O3S	2.03	113.11	106.92
2	E	1	8EX	O4S-S2-O6S	2.00	115.56	108.56

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	G	1	8EX	C4-O4-S2-O5S
2	G	1	8EX	C8-C7-N2-C2
2	G	1	8EX	O7-C7-N2-C2
2	E	1	8EX	C4-O4-S2-O5S
2	E	1	8EX	C4-O4-S2-O6S
2	F	1	8EX	C4-O4-S2-O5S
2	G	1	8EX	C4-O4-S2-O6S
2	H	1	8EX	C4-O4-S2-O5S
2	H	1	8EX	C4-O4-S2-O6S
2	G	1	8EX	C6-O6-S1-O2S
2	E	1	8EX	C4-O4-S2-O4S
2	F	1	8EX	C4-O4-S2-O4S
2	G	1	8EX	C4-O4-S2-O4S
2	H	1	8EX	C4-O4-S2-O4S
2	F	1	8EX	C6-O6-S1-O2S
2	H	1	8EX	C6-O6-S1-O1S
2	G	1	8EX	C6-O6-S1-O1S
2	H	1	8EX	C5-C6-O6-S1
2	E	1	8EX	C5-C6-O6-S1


There are no ring outliers.

1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	G	1	8EX	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.


Oligosaccharide Chain E




Bond lengths



Bond angles




Torsions




Rings


Oligosaccharide Chain F




Oligosaccharide Chain G




Bond lengths




Bond angles



Torsions



Rings



5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	474/505 (93%)	-1.07	0 100 100	14, 23, 36, 67	0
1	B	474/505 (93%)	-1.08	0 100 100	14, 23, 35, 58	0
1	C	474/505 (93%)	-1.07	0 100 100	14, 23, 36, 61	0
1	D	474/505 (93%)	-1.08	0 100 100	14, 23, 36, 66	0
All	All	1896/2020 (93%)	-1.08	0 100 100	14, 23, 36, 67	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	8EX	E	1	23/23	0.99	0.04	16,19,35,43	0
2	GCD	E	2	11/12	0.99	0.03	14,16,18,20	0
2	8EX	F	1	23/23	0.99	0.04	15,20,38,47	0
2	GCD	F	2	11/12	0.99	0.03	14,16,18,20	0
2	8EX	G	1	23/23	0.99	0.04	9,21,40,49	0
2	GCD	G	2	11/12	0.99	0.03	13,16,18,21	0
2	8EX	H	1	23/23	0.99	0.04	15,20,37,46	0
2	GCD	H	2	11/12	0.99	0.03	14,16,18,19	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain F:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain G:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CA	A	601	1/1	1.00	0.02	27,27,27,27	0
3	CA	B	601	1/1	1.00	0.02	26,26,26,26	0
3	CA	C	601	1/1	1.00	0.01	27,27,27,27	0
3	CA	D	601	1/1	1.00	0.02	27,27,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.