



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 09:42 PM UTC

PDB ID : 8XSA / pdb_00008xsa
Title : Crystal structure of the DNA-bound AHR-ARNT heterodimer in complex with Indigo
Authors : Diao, X.; Shang, Q.; Wu, D.
Deposited on : 2024-01-09
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	382	4% 62% 15% 23%
2	B	395	5% 72% 11% 17%
3	C	21	48% 52%
4	D	21	57% 43%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6076 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aryl hydrocarbon receptor nuclear translocator.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	295	Total	C	N	O	S	0	1	0
			2385	1503	421	443	18			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	84	MET	-	initiating methionine	UNP P27540

- Molecule 2 is a protein called Aryl hydrocarbon receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	327	Total	C	N	O	S	0	0	0
			2664	1708	471	471	14			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	25	MET	-	initiating methionine	UNP I3LF82
B	414	HIS	-	expression tag	UNP I3LF82
B	415	HIS	-	expression tag	UNP I3LF82
B	416	HIS	-	expression tag	UNP I3LF82
B	417	HIS	-	expression tag	UNP I3LF82
B	418	HIS	-	expression tag	UNP I3LF82
B	419	HIS	-	expression tag	UNP I3LF82


- Molecule 3 is a DNA chain called DNAF.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	21	Total	C	N	O	P	0	0	0
			430	204	84	122	20			

- Molecule 4 is a DNA chain called DNAR.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	21	Total	C	N	O	P	0	0	0
			425	203	76	126	20			

- Molecule 5 is 2,3-DIHYDROXY-1,4-DITHIOBUTANE (CCD ID: DTT) (formula: $C_4H_{10}O_2S_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	O	S	0	0
			8	4	2	2		

- Molecule 6 is 2-(3-oxidanyl-1H-indol-2-yl)indol-3-one (CCD ID: A1LWK) (formula: $C_{16}H_{10}N_2O_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	B	1	Total	C	N	O	0	0
			20	16	2	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	71	Total	O	0	0
			71	71		
7	B	67	Total	O	0	0
			67	67		
7	C	4	Total	O	0	0
			4	4		
7	D	2	Total	O	0	0
			2	2		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aryl hydrocarbon receptor nuclear translocator



- Molecule 2: Aryl hydrocarbon receptor




- Molecule 3: DNAF



- Molecule 4: DNAR

Chain D:  57% 43%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	68.02Å 98.84Å 80.31Å 90.00° 90.41° 90.00°	Depositor
Resolution (Å)	30.48 – 2.60 30.48 – 2.60	Depositor EDS
% Data completeness (in resolution range)	94.1 (30.48-2.60) 89.9 (30.48-2.60)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.40 (at 2.61Å)	Xtriage
Refinement program	PHENIX 1.14	Depositor
R, R_{free}	0.195 , 0.249 0.196 , 0.249	Depositor DCC
R_{free} test set	2010 reflections (6.49%)	wwPDB-VP
Wilson B-factor (Å ²)	37.2	Xtriage
Anisotropy	0.134	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 52.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.038 for h,-k,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6076	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A1LWK, DTT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.30	0/2435	0.50	1/3284 (0.0%)
2	B	0.18	0/2717	0.40	0/3658
3	C	0.20	0/483	0.37	0/744
4	D	0.21	0/475	0.42	0/731
All	All	0.24	0/6110	0.44	1/8417 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	117	PRO	N-CA-C	5.24	123.26	112.47

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2385	0	2339	41	0
2	B	2664	0	2697	32	0
3	C	430	0	236	10	1
4	D	425	0	238	9	1
5	A	8	0	10	0	0
6	B	20	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	71	0	0	0	0
7	B	67	0	0	3	0
7	C	4	0	0	1	0
7	D	2	0	0	0	0
All	All	6076	0	5520	82	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:85:ASP:O	1:A:89:LEU:HG	1.85	0.77
2:B:312:GLU:HA	2:B:315:MET:HE2	1.68	0.75
2:B:256:ILE:HD12	2:B:256:ILE:N	2.03	0.74
4:D:13:DA:H2''	4:D:14:DT:H5''	1.72	0.72
2:B:275:ILE:HD12	2:B:353:THR:HG21	1.73	0.69
2:B:98:GLN:NE2	7:B:603:HOH:O	2.32	0.61
1:A:215:PRO:O	1:A:218:VAL:HG12	2.01	0.60
4:D:11:DC:H2''	4:D:12:DG:C8	2.37	0.60
1:A:208:THR:HG22	1:A:210:TYR:H	1.67	0.60
3:C:4:DC:H2''	3:C:5:DG:C8	2.37	0.60
1:A:361:THR:HG22	1:A:362:GLU:H	1.66	0.59
1:A:343:LEU:O	1:A:344:GLN:HB2	2.02	0.59
2:B:343:GLU:OE1	2:B:364:ASN:ND2	2.25	0.58
1:A:362:GLU:HB3	1:A:462:THR:HG23	1.85	0.58
4:D:12:DG:H2''	4:D:13:DA:C8	2.39	0.57
3:C:7:DG:H1	4:D:16:DC:H5	1.51	0.56
2:B:130:THR:HG23	2:B:261:LEU:O	2.06	0.56
3:C:7:DG:H22	4:D:16:DC:H5	1.54	0.56
4:D:9:DC:H2''	4:D:10:DG:C8	2.41	0.56
1:A:203:GLU:O	1:A:203:GLU:HG3	2.05	0.55
2:B:386:ASP:O	2:B:390:LYS:HG3	2.06	0.55
1:A:409:ARG:O	1:A:413:GLN:HG2	2.07	0.54
1:A:401:HIS:HB2	1:A:428:ARG:HB2	1.89	0.54
1:A:205:PHE:CE1	2:B:106:ARG:HD3	2.42	0.54
3:C:2:DA:H2'	3:C:3:DT:C6	2.43	0.54
2:B:362:GLN:OE1	2:B:382:ARG:NH2	2.41	0.54
4:D:15:DG:H2''	4:D:16:DC:O2	2.08	0.53
1:A:449:PRO:HG2	2:B:335:HIS:ND1	2.24	0.52
1:A:257:MET:HG2	1:A:258:GLY:H	1.74	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:378:HIS:HD2	1:A:388:PRO:HG2	1.74	0.51
4:D:14:DT:H2''	4:D:15:DG:H5'	1.92	0.51
2:B:294:THR:HA	2:B:318:THR:HA	1.93	0.51
1:A:88:ARG:HH21	1:A:89:LEU:HD21	1.77	0.49
1:A:446:PHE:HB2	1:A:458:ILE:HD11	1.93	0.49
1:A:204:TRP:O	1:A:207:SER:HB3	2.12	0.49
3:C:11:DC:N4	7:C:101:HOH:O	2.41	0.49
1:A:96:GLU:O	1:A:100:ARG:HG2	2.14	0.48
2:B:233:LEU:HD23	2:B:235:MET:HG2	1.94	0.48
1:A:418:LEU:HB3	1:A:421:GLN:HG3	1.94	0.48
2:B:124:PHE:CE2	2:B:235:MET:HE1	2.48	0.48
2:B:149:GLN:O	2:B:153:ILE:HG13	2.14	0.48
1:A:94:HIS:HE1	3:C:16:DG:O6	1.97	0.47
1:A:377:ASP:OD1	1:A:378:HIS:N	2.47	0.47
1:A:172:ALA:HB3	2:B:266:ILE:HD12	1.94	0.47
4:D:14:DT:H2'	4:D:15:DG:H21	1.78	0.47
3:C:19:DA:H2''	3:C:20:DA:C8	2.50	0.47
1:A:363:PHE:HB2	1:A:377:ASP:HB3	1.96	0.47
3:C:6:DG:H2'	3:C:6:DG:N3	2.31	0.46
2:B:330:TYR:CZ	2:B:396:ARG:HD3	2.50	0.45
2:B:343:GLU:OE2	2:B:366:ARG:NE	2.43	0.45
1:A:127:ASP:OD1	1:A:130:THR:HG23	2.17	0.45
2:B:330:TYR:CE2	2:B:396:ARG:HD3	2.52	0.45
2:B:390:LYS:HB2	2:B:390:LYS:HE3	1.77	0.44
2:B:250:LYS:C	2:B:257:LEU:HD13	2.42	0.44
2:B:408:GLU:HA	7:B:613:HOH:O	2.17	0.43
2:B:338:MET:HE2	2:B:338:MET:HB3	1.92	0.43
1:A:446:PHE:HB2	1:A:458:ILE:CD1	2.48	0.43
1:A:174:GLY:HA2	1:A:341:GLY:O	2.18	0.43
1:A:401:HIS:O	1:A:405:GLN:HG3	2.19	0.43
2:B:399:LYS:HD2	2:B:403:MET:HE1	2.01	0.43
1:A:268:ARG:HG2	1:A:303:PHE:CE2	2.54	0.43
1:A:443:SER:HA	1:A:458:ILE:O	2.19	0.43
1:A:208:THR:HG22	1:A:210:TYR:N	2.33	0.43
1:A:397:VAL:HG11	1:A:409:ARG:HE	1.84	0.43
1:A:209:LEU:HG	1:A:221:LEU:HD21	2.00	0.42
1:A:387:GLN:HB2	1:A:390:GLU:HG3	2.00	0.42
1:A:392:LEU:HD11	2:B:408:GLU:HB2	2.01	0.42
2:B:75:SER:OG	2:B:154:HIS:HB2	2.20	0.42
1:A:432:LYS:C	1:A:434:GLN:H	2.27	0.42
1:A:432:LYS:HB3	1:A:432:LYS:HE2	1.83	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:127:ASP:HB2	3:C:11:DC:OP1	2.19	0.41
1:A:170:GLU:HG3	2:B:242:LYS:HD3	2.01	0.41
2:B:397:THR:HB	2:B:398:LEU:H	1.77	0.41
1:A:181:CYS:HB3	1:A:225:LEU:O	2.20	0.41
1:A:458:ILE:HG12	2:B:328:MET:HE1	2.01	0.41
2:B:275:ILE:HB	2:B:359:THR:HB	2.02	0.41
3:C:1:DC:H2'	3:C:2:DA:C8	2.56	0.41
2:B:106:ARG:NH1	7:B:607:HOH:O	2.54	0.41
1:A:179:VAL:HG11	1:A:209:LEU:HB2	2.03	0.40
1:A:210:TYR:CD1	1:A:218:VAL:HG23	2.56	0.40
2:B:118:LEU:HD12	2:B:121:LEU:HD12	2.03	0.40
2:B:276:LEU:CD2	2:B:386:ASP:H	2.35	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:1:DC:O2	4:D:1:DG:N2[1_655]	2.18	0.02

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	284/382 (74%)	270 (95%)	13 (5%)	1 (0%)	30	51
2	B	317/395 (80%)	309 (98%)	7 (2%)	1 (0%)	36	58
All	All	601/777 (77%)	579 (96%)	20 (3%)	2 (0%)	36	58

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	273	PRO

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	117	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	270/346 (78%)	268 (99%)	2 (1%)	76	89
2	B	290/347 (84%)	289 (100%)	1 (0%)	86	94
All	All	560/693 (81%)	557 (100%)	3 (0%)	81	92

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	84	MET
1	A	202	SER
2	B	256	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	93	ASN
2	B	98	GLN
2	B	245	HIS
2	B	381	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	DTT	A	501	-	7,7,7	0.81	0	4,8,8	1.16	0
6	A1LWK	B	501	-	23,23,23	2.28	7 (30%)	28,34,34	1.67	4 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	DTT	A	501	-	-	6/8/8/8	-
6	A1LWK	B	501	-	-	0/4/16/16	0/4/4/4

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	501	A1LWK	C12-N11	6.15	1.52	1.39
6	B	501	A1LWK	C10-N11	3.91	1.45	1.32
6	B	501	A1LWK	C08-C10	3.59	1.52	1.44
6	B	501	A1LWK	C08-N09	-3.32	1.33	1.38
6	B	501	A1LWK	C05-C04	-3.05	1.37	1.41
6	B	501	A1LWK	O20-C07	2.60	1.40	1.33
6	B	501	A1LWK	O19-C18	-2.12	1.18	1.23

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	501	A1LWK	C17-C12-N11	-6.47	108.08	112.91
6	B	501	A1LWK	C10-C08-N09	2.65	127.65	118.67
6	B	501	A1LWK	C03-C04-C05	-2.57	119.70	122.19
6	B	501	A1LWK	C08-C10-N11	2.20	125.44	118.71

There are no chirality outliers.


All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	501	DTT	S1-C1-C2-O2
5	A	501	DTT	S1-C1-C2-C3
5	A	501	DTT	C1-C2-C3-C4
5	A	501	DTT	C1-C2-C3-O3
5	A	501	DTT	O2-C2-C3-C4
5	A	501	DTT	C2-C3-C4-S4

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	295/382 (77%)	0.19	16 (5%) 31 26	16, 41, 78, 126	1 (0%)
2	B	327/395 (82%)	0.08	20 (6%) 27 22	24, 37, 70, 119	0
3	C	21/21 (100%)	0.26	0 100 100	50, 63, 97, 100	0
4	D	21/21 (100%)	0.30	0 100 100	43, 68, 88, 94	0
All	All	664/819 (81%)	0.14	36 (5%) 31 26	16, 41, 80, 126	1 (0%)

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	330	GLY	5.4
2	B	280	THR	5.0
2	B	213	PHE	4.8
2	B	282	ASN	4.3
2	B	273	PRO	4.2
2	B	272	PRO	3.9
1	A	84	MET	3.9
2	B	277	GLU	3.9
2	B	281	LYS	3.7
2	B	278	ILE	3.6
1	A	257	MET	3.4
1	A	119	CYS	3.3
1	A	89	LEU	3.2
1	A	121	ALA	3.2
2	B	33	SER	3.2
2	B	397	THR	3.0
2	B	274	SER	2.9
2	B	100	ASN	2.9
1	A	344	GLN	2.7
1	A	125	LYS	2.7
1	A	270	GLY	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	275	ILE	2.5
2	B	88	LYS	2.5
2	B	223	ARG	2.5
1	A	227	THR	2.4
2	B	232	PHE	2.4
1	A	271	SER	2.4
1	A	302	HIS	2.4
1	A	117	PRO	2.3
1	A	360	PRO	2.3
1	A	258	GLY	2.3
1	A	118	THR	2.2
2	B	96	GLY	2.1
2	B	256	ILE	2.1
2	B	276	LEU	2.1
2	B	355	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	DTT	A	501	8/8	0.89	0.17	70,73,83,87	0
6	A1LWK	B	501	20/20	0.95	0.09	35,43,48,65	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around A1LWK B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.