



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 07:13 PM UTC

PDB ID : 4XTR / pdb_00004xtr
Title : Structure of Get3 bound to the transmembrane domain of Pep12
Authors : Mateja, A.; Paduch, M.; Chang, H.-Y.; Szydlowska, A.; Kossiakoff, A.A.;
Hegde, R.S.; Keenan, R.J.
Deposited on : 2015-01-23
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2260 (2.04-2.04)
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)
RSRZ outliers	180081	2260 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	354	 6% 81% 5% 14%
1	B	354	 4% 81% 6% 12%
2	C	230	 2% 91% 6% 3%
2	E	230	 96% 4%
3	D	217	 3% 94% 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	217	2% 95% 5%
4	G	37	38% 43% 16% 41%

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 24293 atoms, of which 11564 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATPase GET3.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	305	Total	C	H	N	O	S	0	2	0
			4815	1530	2400	401	466	18			
1	B	311	Total	C	H	N	O	S	0	1	0
			4873	1543	2432	404	476	18			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	57	ASN	ASP	engineered mutation	UNP Q12154
B	57	ASN	ASP	engineered mutation	UNP Q12154

- Molecule 2 is a protein called Antibody Heavy chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	C	217	Total	C	H	N	O	S	0	1	0
			3241	1037	1600	278	320	6			
2	E	222	Total	C	H	N	O	S	0	1	0
			3306	1055	1633	284	328	6			

- Molecule 3 is a protein called Antibody Light chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	D	216	Total	C	H	N	O	S	0	0	0
			3269	1038	1611	276	338	6			
3	F	216	Total	C	H	N	O	S	0	0	0
			3269	1038	1611	276	338	6			


- Molecule 4 is a protein called Pep12p.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
4	G	22	Total	C	H	N	O	S	0	0	0
			427	141	231	30	23	2			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	252	MET	-	initiating methionine	UNP E7M086
G	253	GLY	-	expression tag	UNP E7M086
G	254	SER	-	expression tag	UNP E7M086
G	255	HIS	-	expression tag	UNP E7M086
G	256	HIS	-	expression tag	UNP E7M086
G	257	HIS	-	expression tag	UNP E7M086
G	258	HIS	-	expression tag	UNP E7M086
G	259	HIS	-	expression tag	UNP E7M086
G	260	HIS	-	expression tag	UNP E7M086
G	261	SER	-	expression tag	UNP E7M086

- Molecule 5 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
5	A	1	Total	C	H	N	O	P	0	1
			38	10	11	5	10	2		
5	B	1	Total	C	H	N	O	P	0	1
			39	10	12	5	10	2		


- Molecule 6 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Mg	0	0
			1	1		
6	B	1	Total	Mg	0	0
			1	1		

- Molecule 7 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total Zn 1 1	0	0

- Molecule 8 is ADENOSINE-5'-TRIPHOSPHATE (CCD ID: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
8	A	1	Total 42	C 10	H 11	N 5	O 13	P 3	0	1
8	B	1	Total 43	C 10	H 12	N 5	O 13	P 3	0	1

- Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	104	Total O 104 104	0	0
9	B	149	Total O 149 149	0	0
9	C	162	Total O 162 162	0	0
9	D	164	Total O 164 164	0	0
9	E	211	Total O 211 211	0	0
9	F	137	Total O 137 137	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	G	1	Total	O	0	0
			1	1		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: ATPase GET3




• Molecule 1: ATPase GET3




• Molecule 2: Antibody Heavy chain





• Molecule 2: Antibody Heavy chain




• Molecule 3: Antibody Light chain



• Molecule 3: Antibody Light chain



• Molecule 4: Pep12p



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	102.87Å 112.03Å 153.69Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	53.95 – 2.05 53.95 – 2.05	Depositor EDS
% Data completeness (in resolution range)	97.8 (53.95-2.05) 92.1 (53.95-2.05)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.74 (at 2.05Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1839)	Depositor
R, R_{free}	0.187 , 0.216 0.190 , 0.218	Depositor DCC
R_{free} test set	5469 reflections (4.28%)	wwPDB-VP
Wilson B-factor (Å ²)	34.9	Xtriage
Anisotropy	0.474	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 52.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	24293	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ATP, ADP, MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.28	0/2457	0.60	0/3310
1	B	0.30	0/2482	0.61	0/3345
2	C	0.28	0/1682	0.57	0/2293
2	E	0.27	0/1715	0.59	0/2338
3	D	0.29	0/1694	0.57	0/2299
3	F	0.28	0/1694	0.57	0/2299
4	G	0.29	0/200	0.76	0/269
All	All	0.28	0/11924	0.59	0/16153

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2415	2400	2397	8	0
1	B	2441	2432	2431	9	0
2	C	1641	1600	1598	4	0
2	E	1673	1633	1632	0	0
3	D	1658	1611	1611	5	0
3	F	1658	1611	1611	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	G	196	231	230	4	0
5	A	27	11	12	0	0
5	B	27	12	12	0	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
7	A	1	0	0	0	0
8	A	31	11	12	0	0
8	B	31	12	12	0	0
9	A	104	0	0	0	0
9	B	149	0	0	1	0
9	C	162	0	0	2	0
9	D	164	0	0	0	0
9	E	211	0	0	0	0
9	F	137	0	0	2	0
9	G	1	0	0	0	0
All	All	12729	11564	11558	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:320:GLU:OE1	1:B:322:ARG:NH1	2.01	0.93
3:D:111:ARG:NH1	3:D:112:THR:O	2.10	0.83
1:A:277:ALA:O	1:A:279:ASN:N	2.17	0.77
3:F:192:HIS:O	3:F:214:ARG:NH1	2.24	0.70
1:B:307:GLU:OE2	9:B:599:HOH:O	2.15	0.63
3:F:141:ASN:ND2	9:F:412:HOH:O	2.31	0.63
3:D:108:GLU:OE2	3:D:176:TYR:OH	2.17	0.56
2:C:107:ARG:NH1	9:C:439:HOH:O	2.37	0.56
1:B:220:LYS:NZ	1:B:224:GLU:OE2	2.44	0.51
1:A:275:LEU:O	1:A:277:ALA:N	2.42	0.50
1:B:97:MET:HE1	4:G:282:ILE:HG23	1.96	0.47
3:F:25:ARG:NE	3:F:71:ASP:OD2	2.37	0.47
4:G:281:PHE:CZ	4:G:285:ILE:HD11	2.50	0.47
4:G:277:VAL:O	4:G:281:PHE:N	2.49	0.46
3:F:145:ARG:NH1	9:F:413:HOH:O	2.49	0.45
4:G:279:LEU:HA	4:G:282:ILE:HD12	1.99	0.45
1:A:64:ASP:OD1	1:B:254:ARG:NH1	2.48	0.45
1:A:215:LYS:NZ	1:A:218:GLU:OE2	2.46	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:D:5:MET:HE3	3:D:24:CYS:SG	2.58	0.43
3:F:15:SER:N	3:F:18:ASP:OD2	2.39	0.43
1:B:186:LEU:HD23	1:B:186:LEU:C	2.43	0.43
1:B:168:ALA:O	1:B:173:THR:OG1	2.36	0.43
1:A:246[B]:PHE:CG	1:A:247:LEU:N	2.83	0.43
2:C:90:ARG:NH1	9:C:357:HOH:O	2.43	0.43
2:C:210:LYS:N	2:C:211:PRO:HD2	2.33	0.43
1:A:320:GLU:OE2	1:B:298:TYR:OH	2.28	0.43
1:A:277:ALA:O	1:A:279:ASN:ND2	2.52	0.42
3:D:135:VAL:HG13	3:D:182:LEU:HB3	2.01	0.42
3:F:5:MET:HE3	3:F:24:CYS:SG	2.58	0.42
2:C:107:ARG:NH2	3:D:96:SER:O	2.53	0.41
1:A:246[B]:PHE:CZ	1:B:322:ARG:HB3	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	297/354 (84%)	285 (96%)	10 (3%)	2 (1%)	18	10
1	B	304/354 (86%)	297 (98%)	5 (2%)	2 (1%)	18	10
2	C	214/230 (93%)	208 (97%)	6 (3%)	0	100	100
2	E	221/230 (96%)	218 (99%)	3 (1%)	0	100	100
3	D	214/217 (99%)	209 (98%)	5 (2%)	0	100	100
3	F	214/217 (99%)	210 (98%)	4 (2%)	0	100	100
4	G	20/37 (54%)	19 (95%)	1 (5%)	0	100	100
All	All	1484/1639 (90%)	1446 (97%)	34 (2%)	4 (0%)	36	30

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	170	THR
1	A	170	THR
1	A	278	GLU
1	B	285	CYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	273/309 (88%)	267 (98%)	6 (2%)	45	44
1	B	276/309 (89%)	266 (96%)	10 (4%)	31	26
2	C	182/193 (94%)	179 (98%)	3 (2%)	55	56
2	E	186/193 (96%)	183 (98%)	3 (2%)	55	56
3	D	191/192 (100%)	187 (98%)	4 (2%)	47	46
3	F	191/192 (100%)	191 (100%)	0	100	100
4	G	22/36 (61%)	21 (96%)	1 (4%)	24	18
All	All	1321/1424 (93%)	1294 (98%)	27 (2%)	48	48

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	126	LEU
1	A	216	LEU
1	A	254	ARG
1	A	283	HIS
1	A	332	GLN
1	A	351	GLU
1	B	100	MET
1	B	132	SER
1	B	149	ILE
1	B	151	ARG
1	B	181	ASN
1	B	193	ILE
1	B	194	THR
1	B	228	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	266	VAL
1	B	353	LYS
2	C	32	LEU
2	C	103	ARG
2	C	206	ASN
3	D	108	GLU
3	D	130	SER
3	D	136	VAL
3	D	202	GLN
2	E	32	LEU
2	E	114[A]	GLN
2	E	114[B]	GLN
4	G	276	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	301	GLN
1	A	332	GLN
1	B	61	ASN
1	B	301	GLN
1	B	335	ASN
2	C	173	HIS
3	D	163	GLN
2	E	85	GLN
2	E	87	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 7 ligands modelled in this entry, 3 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	ADP	B	401[B]	6	28,29,29	3.50	10 (35%)	43,45,45	2.13	10 (23%)
5	ADP	A	401[B]	6	28,29,29	3.50	10 (35%)	43,45,45	2.13	10 (23%)
8	ATP	B	403[A]	6	32,33,33	2.60	10 (31%)	48,52,52	2.21	12 (25%)
8	ATP	A	404[A]	6	32,33,33	2.62	12 (37%)	48,52,52	2.23	12 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	ADP	B	401[B]	6	-	0/16/32/32	0/3/3/3
5	ADP	A	401[B]	6	-	0/16/32/32	0/3/3/3
8	ATP	B	403[A]	6	-	1/22/38/38	0/3/3/3
8	ATP	A	404[A]	6	-	0/22/38/38	0/3/3/3

All (42) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	401[B]	ADP	PA-O3A	9.93	1.70	1.59
5	B	401[B]	ADP	PA-O3A	9.81	1.70	1.59
5	B	401[B]	ADP	O4'-C1'	8.64	1.62	1.42
5	A	401[B]	ADP	O4'-C1'	8.61	1.61	1.42
5	A	401[B]	ADP	C2'-C1'	-6.94	1.31	1.53
5	B	401[B]	ADP	C2'-C1'	-6.91	1.31	1.53
5	B	401[B]	ADP	C6-N6	6.38	1.50	1.34
5	A	401[B]	ADP	C6-N6	6.37	1.50	1.34
8	B	403[A]	ATP	PB-O3A	5.79	1.65	1.59
8	A	404[A]	ATP	PB-O3A	5.78	1.65	1.59
5	B	401[B]	ADP	O4'-C4'	-5.73	1.32	1.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	401[B]	ADP	O4'-C4'	-5.71	1.32	1.45
8	B	403[A]	ATP	C2-N1	-5.49	1.24	1.33
8	A	404[A]	ATP	C2-N1	-5.47	1.24	1.33
8	A	404[A]	ATP	C8-N7	5.23	1.41	1.31
8	B	403[A]	ATP	C8-N7	5.22	1.41	1.31
8	A	404[A]	ATP	C5-C4	5.20	1.48	1.39
8	B	403[A]	ATP	C5-C4	5.19	1.48	1.39
8	A	404[A]	ATP	C8-N9	3.80	1.44	1.37
8	B	403[A]	ATP	C8-N9	3.77	1.44	1.37
8	B	403[A]	ATP	C6-N6	3.74	1.43	1.34
8	A	404[A]	ATP	C6-N6	3.74	1.43	1.34
8	B	403[A]	ATP	C4-N3	3.73	1.41	1.34
8	A	404[A]	ATP	C4-N3	3.73	1.41	1.34
8	A	404[A]	ATP	C6-N1	-3.67	1.25	1.35
8	B	403[A]	ATP	C6-N1	-3.65	1.25	1.35
5	A	401[B]	ADP	O2'-C2'	3.13	1.50	1.43
5	B	401[B]	ADP	O2'-C2'	3.12	1.50	1.43
8	A	404[A]	ATP	C2-N3	-2.78	1.29	1.33
8	B	403[A]	ATP	C2-N3	-2.77	1.29	1.33
5	B	401[B]	ADP	C5-C4	-2.25	1.35	1.39
5	B	401[B]	ADP	C5-N7	-2.25	1.35	1.39
8	B	403[A]	ATP	C2'-C3'	-2.23	1.47	1.53
5	A	401[B]	ADP	C5-C4	-2.22	1.35	1.39
5	A	401[B]	ADP	C5-N7	-2.21	1.35	1.39
5	B	401[B]	ADP	C2-N1	2.21	1.37	1.33
8	A	404[A]	ATP	C2'-C3'	-2.18	1.47	1.53
5	A	401[B]	ADP	C2-N1	2.14	1.37	1.33
5	B	401[B]	ADP	PA-O5'	2.13	1.67	1.59
5	A	401[B]	ADP	PA-O5'	2.10	1.67	1.59
8	A	404[A]	ATP	PG-O2G	-2.08	1.47	1.54
8	A	404[A]	ATP	O4'-C1'	2.06	1.46	1.42

All (44) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	A	404[A]	ATP	C5-C4-N3	-7.60	116.26	126.72
8	B	403[A]	ATP	C5-C4-N3	-7.44	116.48	126.72
5	A	401[B]	ADP	N6-C6-N1	-6.28	104.39	118.38
5	B	401[B]	ADP	N6-C6-N1	-6.24	104.49	118.38
8	A	404[A]	ATP	N3-C4-N9	5.89	137.18	127.17
8	B	403[A]	ATP	N3-C4-N9	5.86	137.13	127.17
5	B	401[B]	ADP	N3-C2-N1	-5.78	119.83	128.58

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	401[B]	ADP	N3-C2-N1	-5.72	119.92	128.58
8	B	403[A]	ATP	C2-N1-C6	5.69	128.08	118.73
8	A	404[A]	ATP	C2-N1-C6	5.59	127.92	118.73
5	A	401[B]	ADP	C5-C6-N6	5.06	135.80	123.29
5	B	401[B]	ADP	C5-C6-N6	5.05	135.79	123.29
5	A	401[B]	ADP	C5-C4-N3	-4.80	120.10	126.72
5	B	401[B]	ADP	C5-C4-N3	-4.78	120.13	126.72
5	B	401[B]	ADP	N9-C8-N7	-4.07	108.16	113.94
5	A	401[B]	ADP	N9-C8-N7	-4.04	108.20	113.94
8	B	403[A]	ATP	N3-C2-N1	-3.91	122.66	128.58
8	A	404[A]	ATP	N3-C2-N1	-3.86	122.73	128.58
5	A	401[B]	ADP	C2-N3-C4	3.34	119.98	111.83
8	A	404[A]	ATP	C2-N3-C4	3.34	119.98	111.83
5	B	401[B]	ADP	C2-N3-C4	3.34	119.98	111.83
8	A	404[A]	ATP	N9-C8-N7	-3.28	109.28	113.94
8	B	403[A]	ATP	C2-N3-C4	3.27	119.82	111.83
8	B	403[A]	ATP	N9-C8-N7	-3.26	109.32	113.94
8	B	403[A]	ATP	O3G-PG-O3B	3.01	114.73	104.64
8	A	404[A]	ATP	O3G-PG-O3B	2.97	114.60	104.64
5	B	401[B]	ADP	C5-N7-C8	2.91	108.02	103.45
5	A	401[B]	ADP	C5-N7-C8	2.90	108.01	103.45
8	B	403[A]	ATP	O2G-PG-O3B	2.88	114.29	104.64
8	B	403[A]	ATP	C4-N9-C8	2.86	108.75	105.74
8	A	404[A]	ATP	C4-N9-C8	2.79	108.67	105.74
8	A	404[A]	ATP	O2G-PG-O3B	2.74	113.83	104.64
5	A	401[B]	ADP	N3-C4-N9	2.69	131.75	127.17
5	B	401[B]	ADP	N3-C4-N9	2.68	131.73	127.17
8	A	404[A]	ATP	O2B-PB-O1B	-2.39	101.33	112.44
8	A	404[A]	ATP	O2A-PA-O1A	-2.34	101.58	112.44
5	A	401[B]	ADP	C4-C5-N7	-2.33	107.92	110.58
8	B	403[A]	ATP	O2B-PB-O1B	-2.30	101.73	112.44
5	B	401[B]	ADP	C4-C5-N7	-2.29	107.96	110.58
8	B	403[A]	ATP	O2A-PA-O1A	-2.27	101.90	112.44
5	A	401[B]	ADP	C5-C4-N9	2.15	108.15	105.81
5	B	401[B]	ADP	C5-C4-N9	2.14	108.14	105.81
8	A	404[A]	ATP	C6-C5-N7	2.13	136.20	132.09
8	B	403[A]	ATP	C6-C5-N7	2.09	136.12	132.09

There are no chirality outliers.


All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	B	403[A]	ATP	PA-O3A-PB-O2B


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand ADP A 401 (B)



Ligand ATP B 403 (A)





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	305/354 (86%)	0.38	23 (7%) 20 20	24, 55, 134, 149	2 (0%)
1	B	311/354 (87%)	0.07	15 (4%) 35 36	20, 44, 110, 147	1 (0%)
2	C	217/230 (94%)	0.11	5 (2%) 61 63	19, 46, 105, 143	1 (0%)
2	E	222/230 (96%)	-0.31	1 (0%) 87 89	19, 38, 66, 110	1 (0%)
3	D	216/217 (99%)	0.16	6 (2%) 55 57	28, 52, 100, 145	0
3	F	216/217 (99%)	0.10	5 (2%) 61 63	30, 54, 78, 120	0
4	G	22/37 (59%)	2.50	14 (63%) 0 0	98, 125, 136, 155	0
All	All	1509/1639 (92%)	0.13	69 (4%) 37 37	19, 49, 112, 155	5 (0%)

All (69) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	G	279	LEU	5.8
1	A	246[A]	PHE	5.4
1	A	155	GLY	5.0
1	B	123	GLY	4.9
4	G	285	ILE	4.2
4	G	267	TRP	4.0
1	A	190	PHE	4.0
1	B	190	PHE	4.0
1	A	212	ILE	3.9
4	G	275	LEU	3.8
1	A	126	LEU	3.8
4	G	274	VAL	3.6
4	G	281	PHE	3.4
1	A	129	LEU	3.3
1	A	249	LEU	3.3
3	D	157	LEU	3.2
1	B	186	LEU	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	353	LYS	2.9
1	A	5	VAL	2.9
1	A	294	MET	2.9
1	B	101	ALA	2.8
3	F	3	ILE	2.8
1	B	278	GLU	2.8
3	F	77	SER	2.7
1	A	216	LEU	2.7
3	D	24	CYS	2.7
1	A	189	LYS	2.7
1	A	125	ALA	2.7
1	B	194	THR	2.7
4	G	280	LEU	2.6
1	B	103	SER	2.6
3	F	24	CYS	2.6
1	A	124	GLY	2.6
4	G	282	ILE	2.5
2	C	159	VAL	2.5
1	B	193	ILE	2.5
1	A	247	LEU	2.4
1	A	211	ASP	2.4
1	A	159	THR	2.4
1	A	194	THR	2.4
1	A	100	MET	2.3
2	C	137	SER	2.3
1	A	276	PHE	2.3
2	E	225	CYS	2.3
1	A	277	ALA	2.3
4	G	278	MET	2.3
2	C	161	VAL	2.3
3	D	2	ASP	2.3
1	B	191	GLY	2.3
2	C	224	SER	2.2
4	G	269	VAL	2.2
1	B	346	VAL	2.2
4	G	270	TYR	2.2
2	C	223	LYS	2.2
3	D	200	THR	2.2
1	B	212	ILE	2.1
3	F	132	THR	2.1
3	D	135	VAL	2.1
3	D	184	LEU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	F	109	ILE	2.1
4	G	283	PHE	2.1
4	G	287	LYS	2.1
4	G	271	LEU	2.0
1	B	124	GLY	2.0
1	B	100	MET	2.0
1	B	195	ASN	2.0
1	A	186	LEU	2.0
1	A	283	HIS	2.0
1	A	344	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	ADP	A	401[B]	27/27	0.94	0.07	30,36,46,48	38
8	ATP	A	404[A]	31/31	0.94	0.08	30,36,45,48	42
8	ATP	B	403[A]	31/31	0.94	0.08	25,34,45,52	43
5	ADP	B	401[B]	27/27	0.95	0.08	30,36,45,50	39
6	MG	A	402	1/1	0.96	0.07	41,41,41,41	0
6	MG	B	402	1/1	0.96	0.05	35,35,35,35	0
7	ZN	A	403	1/1	0.97	0.04	68,68,68,68	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ADP A 401 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ATP A 404 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around ATP B 403 (A):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ADP B 401 (B):

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.