



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 08:36 PM UTC

PDB ID : 7XTO / pdb_00007xto
Title : Structure of CIA2 reveal the Mechanism of soil bacterial derived chlorinase
Authors : Liu, Y.H.; Liu, Y.; Li, Y.
Deposited on : 2022-05-17
Resolution : 2.48 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7589 (2.50-2.46)
Clashscore	190562	8295 (2.50-2.46)
Ramachandran outliers	187476	8164 (2.50-2.46)
Sidechain outliers	187428	8166 (2.50-2.46)
RSRZ outliers	180081	7593 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	276	
1	B	276	
1	C	276	
1	D	276	
1	E	276	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	276	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into four segments: a red segment on the left labeled '20%', a large green segment labeled '77%', a yellow segment labeled '12%', and a grey segment on the far right labeled '12%'.</p>

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 10672 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called soil bacterial derived chlorinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	245	Total 1706	C 1083	N 293	O 322	S 8	0	0	0
1	C	248	Total 1711	C 1084	N 293	O 326	S 8	0	0	0
1	D	248	Total 1747	C 1106	N 303	O 330	S 8	0	0	0
1	B	252	Total 1752	C 1113	N 299	O 332	S 8	0	0	0
1	E	247	Total 1705	C 1076	N 298	O 323	S 8	0	0	0
1	F	244	Total 1698	C 1075	N 293	O 322	S 8	0	0	0


- Molecule 2 is 1-DEAZA-ADENOSINE (CCD ID: 1DA) (formula: C₁₁H₁₄N₄O₄).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			19	11	4	4		
2	C	1	Total	C	N	O	0	0
			19	11	4	4		
2	D	1	Total	C	N	O	0	0
			19	11	4	4		
2	B	1	Total	C	N	O	0	0
			19	11	4	4		
2	E	1	Total	C	N	O	0	0
			19	11	4	4		
2	F	1	Total	C	N	O	0	0
			19	11	4	4		

- Molecule 3 is water.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	37	Total	O	0	0
			37	37		
3	C	39	Total	O	0	0
			39	39		
3	D	56	Total	O	0	0
			56	56		
3	B	28	Total	O	0	0
			28	28		
3	E	31	Total	O	0	0
			31	31		
3	F	48	Total	O	0	0
			48	48		





• Molecule 1: soil bacterial derived chlorinase



• Molecule 1: soil bacterial derived chlorinase



• Molecule 1: soil bacterial derived chlorinase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	181.38Å 104.42Å 124.97Å 90.00° 108.88° 90.00°	Depositor
Resolution (Å)	24.99 – 2.48 24.99 – 2.48	Depositor EDS
% Data completeness (in resolution range)	86.8 (24.99-2.48) 87.1 (24.99-2.48)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.63 (at 2.48Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.278 , 0.304 0.279 , 0.304	Depositor DCC
R_{free} test set	3353 reflections (4.27%)	wwPDB-VP
Wilson B-factor (Å ²)	36.1	Xtrriage
Anisotropy	0.856	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 78.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	10672	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 56.41 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.7473e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section:
1DA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.20	0/1745	0.51	3/2407 (0.1%)
1	B	0.21	0/1792	0.51	3/2473 (0.1%)
1	C	0.20	0/1750	0.50	3/2417 (0.1%)
1	D	0.21	0/1785	0.50	3/2460 (0.1%)
1	E	0.26	0/1744	0.59	6/2404 (0.2%)
1	F	0.21	0/1735	0.50	3/2389 (0.1%)
All	All	0.21	0/10551	0.52	21/14550 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	E	0	1
All	All	0	2

There are no bond length outliers.

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	266	PRO	N-CA-C	7.18	127.26	112.47
1	F	47	ALA	CA-C-N	6.29	134.35	123.91
1	F	47	ALA	C-N-CA	6.29	134.35	123.91
1	B	47	ALA	CA-C-N	6.29	134.35	123.91
1	B	47	ALA	C-N-CA	6.29	134.35	123.91

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	198	ALA	Mainchain
1	E	198	ALA	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1706	0	1578	22	0
1	B	1752	0	1625	19	0
1	C	1711	0	1570	20	0
1	D	1747	0	1639	20	0
1	E	1705	0	1554	20	0
1	F	1698	0	1552	23	0
2	A	19	0	14	1	0
2	B	19	0	14	2	0
2	C	19	0	14	1	0
2	D	19	0	14	2	0
2	E	19	0	14	2	0
2	F	19	0	14	1	0
3	A	37	0	0	0	0
3	B	28	0	0	0	0
3	C	39	0	0	1	0
3	D	56	0	0	0	0
3	E	31	0	0	0	1
3	F	48	0	0	0	1
All	All	10672	0	9602	115	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 115 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:103:LEU:HB3	1:F:107:LEU:HD23	1.52	0.91
1:E:88:ARG:NH2	1:E:114:GLU:OE1	2.07	0.86
1:E:129:THR:HG21	1:F:242:SER:HA	1.58	0.84
1:C:252:ARG:NH1	3:C:401:HOH:O	2.12	0.83

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:129:THR:HG21	1:C:242:SER:HA	1.60	0.82

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:421:HOH:O	3:F:440:HOH:O[4_555]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	237/276 (86%)	230 (97%)	7 (3%)	0	100	100
1	B	244/276 (88%)	237 (97%)	7 (3%)	0	100	100
1	C	240/276 (87%)	232 (97%)	8 (3%)	0	100	100
1	D	238/276 (86%)	231 (97%)	7 (3%)	0	100	100
1	E	237/276 (86%)	227 (96%)	8 (3%)	2 (1%)	16	28
1	F	232/276 (84%)	225 (97%)	7 (3%)	0	100	100
All	All	1428/1656 (86%)	1382 (97%)	44 (3%)	2 (0%)	48	66

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	266	PRO
1	E	267	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	161/217 (74%)	161 (100%)	0	100	100
1	B	166/217 (76%)	165 (99%)	1 (1%)	78	90
1	C	161/217 (74%)	159 (99%)	2 (1%)	63	81
1	D	170/217 (78%)	169 (99%)	1 (1%)	78	90
1	E	158/217 (73%)	158 (100%)	0	100	100
1	F	159/217 (73%)	159 (100%)	0	100	100
All	All	975/1302 (75%)	971 (100%)	4 (0%)	84	92

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	88	ARG
1	C	237	LEU
1	D	237	LEU
1	B	154	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	1DA	A	301	-	20,21,21	1.35	3 (15%)	29,31,31	2.01	11 (37%)
2	1DA	D	301	-	20,21,21	1.35	3 (15%)	29,31,31	2.01	11 (37%)
2	1DA	F	301	-	20,21,21	1.35	3 (15%)	29,31,31	2.01	11 (37%)
2	1DA	E	301	-	20,21,21	1.35	3 (15%)	29,31,31	2.01	11 (37%)
2	1DA	C	301	-	20,21,21	1.35	3 (15%)	29,31,31	2.01	11 (37%)
2	1DA	B	301	-	20,21,21	1.35	3 (15%)	29,31,31	2.01	11 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1DA	A	301	-	-	2/6/22/22	0/3/3/3
2	1DA	D	301	-	-	2/6/22/22	0/3/3/3
2	1DA	F	301	-	-	2/6/22/22	0/3/3/3
2	1DA	E	301	-	-	2/6/22/22	0/3/3/3
2	1DA	C	301	-	-	2/6/22/22	0/3/3/3
2	1DA	B	301	-	-	2/6/22/22	0/3/3/3

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	1DA	C5-N7	-2.83	1.33	1.39
2	B	301	1DA	C5-N7	-2.82	1.33	1.39
2	F	301	1DA	C5-N7	-2.82	1.33	1.39
2	E	301	1DA	C5-N7	-2.82	1.33	1.39
2	D	301	1DA	C5-N7	-2.82	1.33	1.39

The worst 5 of 66 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	301	1DA	C5-C4-N3	-4.65	120.31	126.72
2	B	301	1DA	C5-C4-N3	-4.65	120.31	126.72
2	E	301	1DA	C5-C4-N3	-4.65	120.31	126.72
2	C	301	1DA	C5-C4-N3	-4.65	120.31	126.72
2	A	301	1DA	C5-C4-N3	-4.65	120.31	126.72

There are no chirality outliers.

5 of 12 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	A	301	1DA	O4'-C4'-C5'-O5'
2	C	301	1DA	O4'-C4'-C5'-O5'
2	D	301	1DA	O4'-C4'-C5'-O5'
2	B	301	1DA	O4'-C4'-C5'-O5'
2	E	301	1DA	O4'-C4'-C5'-O5'



There are no ring outliers.


6 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	1DA	1	0
2	D	301	1DA	2	0
2	F	301	1DA	1	0
2	E	301	1DA	2	0
2	C	301	1DA	1	0
2	B	301	1DA	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	245/276 (88%)	1.12	48 (19%) 3 2	24, 40, 71, 100	0
1	B	252/276 (91%)	0.80	30 (11%) 9 7	24, 40, 75, 93	0
1	C	248/276 (89%)	1.03	40 (16%) 4 3	24, 40, 75, 112	0
1	D	248/276 (89%)	1.29	53 (21%) 2 2	24, 40, 74, 94	0
1	E	247/276 (89%)	1.31	55 (22%) 2 2	24, 40, 73, 94	0
1	F	244/276 (88%)	1.42	56 (22%) 2 1	24, 40, 73, 112	0
All	All	1484/1656 (89%)	1.16	282 (19%) 3 2	24, 40, 74, 112	0

The worst 5 of 282 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	127	PRO	6.8
1	D	127	PRO	6.0
1	E	166	ASP	5.9
1	F	126	HIS	5.9
1	B	264	ASP	5.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	1DA	C	301	19/19	0.90	0.12	20,35,53,69	0
2	1DA	D	301	19/19	0.90	0.12	20,35,53,69	0
2	1DA	B	301	19/19	0.90	0.12	20,35,53,69	0
2	1DA	E	301	19/19	0.90	0.12	20,35,53,69	0
2	1DA	F	301	19/19	0.91	0.12	20,35,53,69	0
2	1DA	A	301	19/19	0.92	0.12	20,35,53,69	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around 1DA D 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 1DA B 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around 1DA E 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 1DA F 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.