



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 07:02 PM EDT

PDB ID : 6XUN / pdb_00006xun
Title : Ab 5b1 bound to CA19-9
Authors : Diskin, R.; Borenstein-Katz, A.
Deposited on : 2020-01-20
Resolution : 2.41 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6062 (2.44-2.40)
Clashscore	190562	6562 (2.44-2.40)
Ramachandran outliers	187476	6481 (2.44-2.40)
Sidechain outliers	187428	6482 (2.44-2.40)
RSRZ outliers	180081	6066 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	230	
1	C	230	
1	H	230	
2	B	218	
2	D	218	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	L	218	% 94% 6%
3	E	4	25% 50% 25%
3	F	4	25% 25% 50%
3	G	4	25% 75%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 10655 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	H	220	Total	C	N	O	S	0	0	0
			1674	1055	290	322	7			
1	A	218	Total	C	N	O	S	0	1	0
			1665	1051	288	319	7			
1	C	221	Total	C	N	O	S	0	1	0
			1686	1063	292	324	7			

- Molecule 2 is a protein called Light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	217	Total	C	N	O	S	0	1	0
			1651	1026	284	336	5			
2	D	217	Total	C	N	O	S	0	1	0
			1651	1026	284	336	5			
2	L	217	Total	C	N	O	S	0	1	0
			1651	1026	284	336	5			

- Molecule 3 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-3)-[alpha-L-fucopyranose-(1-4)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	4	Total	C	N	O	0	0	0
			56	31	2	23			
3	F	4	Total	C	N	O	0	0	0
			56	31	2	23			
3	G	4	Total	C	N	O	0	0	0
			56	31	2	23			

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	D	1	Total	C	O	0	0
			6	3	3		
4	C	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	H	84	Total	O	0	0
			84	84		
5	B	76	Total	O	0	0
			76	76		
5	D	62	Total	O	0	0
			62	62		
5	L	52	Total	O	0	0
			52	52		
5	A	112	Total	O	0	0
			112	112		
5	C	99	Total	O	0	0
			99	99		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Heavy chain



- Molecule 1: Heavy chain



- Molecule 1: Heavy chain



- Molecule 2: Light chain




- Molecule 2: Light chain





• Molecule 2: Light chain



• Molecule 3: N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-3)-[alpha-L-fucopyranose-(1-4)]2-acetamido-2-deoxy-beta-D-glucopyranose



• Molecule 3: N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-3)-[alpha-L-fucopyranose-(1-4)]2-acetamido-2-deoxy-beta-D-glucopyranose



• Molecule 3: N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-3)-[alpha-L-fucopyranose-(1-4)]2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, α , β , γ	152.54Å 152.54Å 60.89Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.58 – 2.41 47.58 – 2.41	Depositor EDS
% Data completeness (in resolution range)	99.8 (47.58-2.41) 99.7 (47.58-2.41)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.47 (at 2.42Å)	Xtriage
Refinement program	PHENIX 1.16_3549	Depositor
R, R_{free}	0.169 , 0.215 0.170 , 0.216	Depositor DCC
R_{free} test set	2006 reflections (3.28%)	wwPDB-VP
Wilson B-factor (Å ²)	41.5	Xtriage
Anisotropy	0.049	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 36.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.010 for -h,-k,l 0.014 for h,-h-k,-l 0.000 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10655	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NAG, FUC, GAL, SIA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.27	0/1709	0.50	0/2323
1	C	0.27	0/1730	0.46	0/2350
1	H	0.25	0/1715	0.45	0/2331
2	B	0.29	0/1691	0.53	1/2300 (0.0%)
2	D	0.23	0/1691	0.46	0/2300
2	L	0.22	0/1691	0.44	0/2300
All	All	0.26	0/10227	0.47	1/13904 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	126	LYS	CB-CG-CD	-8.77	91.14	111.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1665	0	1624	13	0
1	C	1686	0	1647	21	0
1	H	1674	0	1630	7	0
2	B	1651	0	1588	13	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	1651	0	1588	13	0
2	L	1651	0	1588	8	0
3	E	56	0	49	1	0
3	F	56	0	49	1	0
3	G	56	0	49	0	0
4	B	12	0	16	0	0
4	C	6	0	8	0	0
4	D	6	0	8	1	0
5	A	112	0	0	4	0
5	B	76	0	0	3	0
5	C	99	0	0	4	1
5	D	62	0	0	2	0
5	H	84	0	0	2	0
5	L	52	0	0	2	0
All	All	10655	0	9844	75	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:117:LYS:NZ	5:A:404:HOH:O	2.15	0.80
1:A:100:SER:OG	5:A:401:HOH:O	2.00	0.80
1:C:43:LYS:O	5:C:401:HOH:O	2.04	0.75
2:B:72:SER:OG	5:B:401:HOH:O	2.03	0.75
2:L:177:SER:OG	5:L:301:HOH:O	2.05	0.74
1:A:46:GLU:OE1	5:A:402:HOH:O	2.05	0.74
2:B:190:LYS:HE3	2:B:210:ASN:HB3	1.71	0.71
1:C:68:THR:OG1	5:C:402:HOH:O	2.12	0.68
5:H:401:HOH:O	3:E:4:FUC:O4	2.11	0.67
1:C:127:SER:OG	1:C:128:SER:N	2.20	0.67
1:C:56:ARG:HG3	1:C:100(A):THR:HG23	1.76	0.66
2:D:40:PRO:O	5:D:402:HOH:O	2.12	0.66
2:B:1:GLN:HG3	2:B:2:SER:H	1.61	0.65
1:A:165:THR:O	5:A:403:HOH:O	2.15	0.65
1:C:40:PRO:HG2	1:C:43:LYS:HB2	1.83	0.61
1:H:126:PRO:HG3	1:H:138:LEU:HB3	1.82	0.60
1:C:76:ASN:OD1	5:C:403:HOH:O	2.16	0.60
1:C:126:PRO:HD2	1:C:213:PRO:HA	1.84	0.60
2:D:80:SER:HA	2:D:106:VAL:HG21	1.83	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:161:GLU:OE1	5:L:303:HOH:O	2.17	0.58
1:H:126:PRO:HD2	1:H:213:PRO:HA	1.85	0.58
1:H:82:MET:HB3	1:H:82(C):LEU:HD21	1.85	0.58
2:B:39:LEU:HD23	2:B:84:ALA:HB2	1.85	0.57
1:C:129:LYS:HD2	1:C:136:ALA:HA	1.86	0.57
2:D:61:ARG:NH1	2:D:82:ASP:OD2	2.28	0.56
3:F:2:GAL:H62	3:F:4:FUC:H3	1.87	0.56
2:B:18:ARG:NH1	5:B:403:HOH:O	2.26	0.55
2:L:49:TYR:HD2	2:L:50:ARG:HG3	1.70	0.54
1:A:95:ASP:HA	1:A:100(E):GLU:O	2.08	0.54
1:C:127:SER:HB3	1:C:129:LYS:HE3	1.91	0.53
1:C:19:ARG:NH2	5:C:408:HOH:O	2.41	0.53
2:L:40:PRO:HB3	2:L:165:GLU:HG3	1.92	0.52
2:D:116:PHE:CD2	1:C:129:LYS:HE2	2.45	0.51
1:A:39:GLN:HB2	1:A:45:LEU:HD23	1.92	0.51
2:L:167:ASP:OD1	2:L:169:LYS:HG3	2.11	0.50
1:A:126:PRO:HG3	1:A:138:LEU:HB3	1.93	0.50
2:D:132:VAL:HG13	2:D:179:LEU:HB3	1.92	0.50
1:C:159:LEU:HD21	1:C:182:VAL:HG21	1.93	0.50
2:D:199:GLN:NE2	5:D:403:HOH:O	2.21	0.49
1:A:159:LEU:HD21	1:A:182:VAL:HG21	1.93	0.49
2:B:4:LEU:HB2	2:B:99:GLY:HA2	1.95	0.49
2:D:14:THR:HG22	2:D:15:PRO:HD2	1.95	0.49
2:B:61:ARG:HB3	2:B:76:SER:O	2.12	0.48
2:L:44:PRO:O	2:L:45:LYS:HE2	2.14	0.48
1:C:59:TYR:HB2	1:C:64:LYS:HD3	1.96	0.47
1:H:1:GLN:N	5:H:406:HOH:O	2.41	0.47
1:C:96:ILE:HG12	1:C:101:GLU:HB2	1.97	0.46
1:H:11:SER:HB2	1:H:147:PRO:HG3	1.97	0.46
1:A:11:SER:HB3	1:A:147:PRO:HG3	1.98	0.46
1:A:214:LYS:HG2	1:A:215:SER:N	2.31	0.45
2:D:92:ASP:O	2:D:95(B):GLY:HA2	2.17	0.45
2:B:126:LYS:NZ	5:B:407:HOH:O	2.49	0.44
2:D:16:GLY:O	2:D:77:GLY:HA2	2.17	0.44
1:C:129:LYS:HD2	1:C:137:ALA:H	1.82	0.44
2:B:54:ARG:HD3	2:B:62:PHE:O	2.17	0.43
2:B:126:LYS:O	2:B:126:LYS:HD3	2.17	0.43
2:D:166:GLN:HG3	2:D:173:TYR:CZ	2.54	0.43
1:C:168:ALA:HB2	1:C:178:LEU:HD23	2.01	0.43
1:C:95:ASP:HA	1:C:100(E):GLU:O	2.19	0.43
4:D:301:GOL:H32	1:C:101:GLU:HG3	2.01	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:35:TRP:HB2	2:D:48:ILE:HB	2.02	0.42
1:C:119:PRO:HB3	1:C:145:TYR:HB3	2.00	0.42
2:B:154:LEU:HG	2:B:155:GLN:N	2.35	0.42
2:B:136:LEU:HB2	2:B:175:LEU:HB3	2.00	0.42
1:A:56:ARG:HA	1:A:56:ARG:HD3	1.91	0.42
2:D:118:PHE:CD2	1:C:124:LEU:HB3	2.55	0.41
1:C:139:GLY:HA2	1:C:154:TRP:CH2	2.56	0.41
1:H:214:LYS:HA	1:H:214:LYS:HD2	1.77	0.41
1:A:13:GLN:HB2	1:A:16:ARG:HG2	2.02	0.41
1:H:214:LYS:HZ2	1:H:214:LYS:HG3	1.71	0.40
2:D:40:PRO:CG	2:D:165:GLU:HG2	2.52	0.40
2:L:49:TYR:CD2	2:L:50:ARG:HG3	2.53	0.40
2:B:184:ALA:O	2:B:188:LYS:HG3	2.22	0.40
2:L:163:VAL:HG22	2:L:175:LEU:HD12	2.03	0.40
1:A:82:MET:HB3	1:A:82(C):LEU:HD21	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:C:406:HOH:O	5:C:490:HOH:O[2_555]	2.11	0.09

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	215/230 (94%)	211 (98%)	4 (2%)	0	100	100
1	C	218/230 (95%)	213 (98%)	4 (2%)	1 (0%)	24	35
1	H	216/230 (94%)	211 (98%)	5 (2%)	0	100	100
2	B	216/218 (99%)	210 (97%)	6 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	D	216/218 (99%)	210 (97%)	6 (3%)	0	100	100
2	L	216/218 (99%)	211 (98%)	5 (2%)	0	100	100
All	All	1297/1344 (96%)	1266 (98%)	30 (2%)	1 (0%)	48	63

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	127	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	184/194 (95%)	184 (100%)	0	100	100
1	C	187/194 (96%)	187 (100%)	0	100	100
1	H	185/194 (95%)	185 (100%)	0	100	100
2	B	187/187 (100%)	187 (100%)	0	100	100
2	D	187/187 (100%)	187 (100%)	0	100	100
2	L	187/187 (100%)	187 (100%)	0	100	100
All	All	1117/1143 (98%)	1117 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	H	39	GLN
1	H	164	HIS
2	B	138	ASN
2	B	210	ASN
2	D	53	GLN
2	D	147	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	L	38	GLN
2	L	138	ASN
2	L	147	GLN
1	A	192	GLN
1	C	199	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	E	1	3	15,15,15	0.23	0	21,21,21	0.42	0
3	GAL	E	2	3	11,11,12	0.62	0	15,15,17	1.14	1 (6%)
3	SIA	E	3	3	20,20,21	1.76	3 (15%)	21,28,31	1.45	2 (9%)
3	FUC	E	4	3	10,10,11	0.94	1 (10%)	14,14,16	0.92	0
3	NAG	F	1	3	15,15,15	0.24	0	21,21,21	0.36	0
3	GAL	F	2	3	11,11,12	1.02	1 (9%)	15,15,17	0.85	0
3	SIA	F	3	3	20,20,21	1.80	3 (15%)	21,28,31	1.40	3 (14%)
3	FUC	F	4	3	10,10,11	0.99	1 (10%)	14,14,16	0.81	0
3	NAG	G	1	3	15,15,15	0.16	0	21,21,21	0.41	0
3	GAL	G	2	3	11,11,12	0.73	0	15,15,17	1.03	1 (6%)
3	SIA	G	3	3	20,20,21	1.72	3 (15%)	21,28,31	1.52	2 (9%)
3	FUC	G	4	3	10,10,11	1.10	1 (10%)	14,14,16	0.72	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	E	1	3	-	1/6/26/26	0/1/1/1
3	GAL	E	2	3	-	0/2/19/22	0/1/1/1
3	SIA	E	3	3	-	3/18/34/38	0/1/1/1
3	FUC	E	4	3	-	-	0/1/1/1
3	NAG	F	1	3	-	0/6/26/26	0/1/1/1
3	GAL	F	2	3	-	2/2/19/22	0/1/1/1
3	SIA	F	3	3	-	2/18/34/38	0/1/1/1
3	FUC	F	4	3	-	-	0/1/1/1
3	NAG	G	1	3	-	0/6/26/26	0/1/1/1
3	GAL	G	2	3	-	0/2/19/22	0/1/1/1
3	SIA	G	3	3	-	4/18/34/38	0/1/1/1
3	FUC	G	4	3	-	-	0/1/1/1

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	3	SIA	C2-C1	6.70	1.60	1.52
3	E	3	SIA	C2-C1	6.49	1.60	1.52
3	G	3	SIA	C2-C1	6.11	1.59	1.52
3	G	4	FUC	O5-C1	-3.01	1.38	1.43
3	F	2	GAL	O5-C1	-2.71	1.39	1.43
3	E	4	FUC	O5-C1	-2.46	1.39	1.43
3	G	3	SIA	C7-C6	2.45	1.56	1.52
3	F	4	FUC	O5-C1	-2.33	1.39	1.43
3	E	3	SIA	O6-C2	2.13	1.47	1.43
3	E	3	SIA	C7-C6	2.06	1.55	1.52
3	G	3	SIA	O6-C2	2.06	1.47	1.43
3	F	3	SIA	O1A-C1	2.04	1.28	1.22
3	F	3	SIA	O6-C2	2.04	1.47	1.43

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	G	3	SIA	O1A-C1-C2	-4.66	112.79	122.85
3	E	3	SIA	O1A-C1-C2	-4.54	113.03	122.85
3	F	3	SIA	O1A-C1-C2	-4.16	113.87	122.85
3	E	2	GAL	C1-O5-C5	3.28	116.58	112.19
3	F	3	SIA	O6-C2-C3	-2.62	107.03	110.56

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	G	3	SIA	O1B-C1-O1A	2.50	129.75	124.08
3	G	2	GAL	O2-C2-C3	-2.47	105.03	110.15
3	E	3	SIA	O1B-C1-O1A	2.26	129.21	124.08
3	F	3	SIA	O1B-C1-O1A	2.18	129.03	124.08

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	G	3	SIA	O8-C8-C9-O9
3	G	3	SIA	C7-C8-C9-O9
3	F	2	GAL	O5-C5-C6-O6
3	F	2	GAL	C4-C5-C6-O6
3	G	3	SIA	C6-C7-C8-O8
3	F	3	SIA	C6-C7-C8-O8
3	E	3	SIA	C7-C8-C9-O9
3	G	3	SIA	O7-C7-C8-O8
3	E	1	NAG	C4-C5-C6-O6
3	E	3	SIA	O8-C8-C9-O9
3	F	3	SIA	O7-C7-C8-O8
3	E	3	SIA	C6-C7-C8-O8


There are no ring outliers.

3 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	2	GAL	1	0
3	E	4	FUC	1	0
3	F	4	FUC	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

Oligosaccharide Chain E



Oligosaccharide Chain F





5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	B	302	-	5,5,5	0.97	0	5,5,5	1.01	0
4	GOL	C	305	-	5,5,5	1.04	0	5,5,5	1.09	0
4	GOL	D	301	-	5,5,5	0.86	0	5,5,5	1.29	0
4	GOL	B	301	-	5,5,5	0.87	0	5,5,5	0.95	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	B	302	-	-	2/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	C	305	-	-	2/4/4/4	-
4	GOL	D	301	-	-	4/4/4/4	-
4	GOL	B	301	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	301	GOL	O1-C1-C2-C3
4	C	305	GOL	C1-C2-C3-O3
4	B	302	GOL	C1-C2-C3-O3
4	D	301	GOL	C1-C2-C3-O3
4	D	301	GOL	O1-C1-C2-O2
4	C	305	GOL	O2-C2-C3-O3
4	B	302	GOL	O2-C2-C3-O3
4	D	301	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	301	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	218/230 (94%)	-0.54	1 (0%) 87 86	24, 35, 65, 84	1 (0%)
1	C	221/230 (96%)	-0.29	7 (3%) 50 47	28, 39, 83, 102	1 (0%)
1	H	220/230 (95%)	-0.21	6 (2%) 56 52	31, 43, 75, 111	0
2	B	217/218 (99%)	-0.32	1 (0%) 87 86	28, 43, 58, 90	1 (0%)
2	D	217/218 (99%)	-0.14	1 (0%) 87 86	31, 47, 69, 86	1 (0%)
2	L	217/218 (99%)	-0.16	3 (1%) 73 70	30, 47, 62, 97	1 (0%)
All	All	1310/1344 (97%)	-0.28	19 (1%) 72 69	24, 44, 69, 111	5 (0%)

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	134	GLY	3.6
2	L	1	GLN	2.9
1	A	215	SER	2.9
1	C	127	SER	2.8
2	D	3	VAL	2.7
1	H	102	TYR	2.6
1	H	214	LYS	2.5
1	H	135	THR	2.5
1	H	191	THR	2.5
2	L	2	SER	2.5
1	C	187	SER	2.4
1	C	215	SER	2.4
2	L	169	LYS	2.3
1	H	216	CYS	2.3
2	B	126	LYS	2.2
1	C	129	LYS	2.2
1	H	134	GLY	2.1
1	C	191	THR	2.1
1	C	135	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	E	1	15/15	-	-	37,44,54,58	0
3	GAL	E	2	11/12	-	-	43,45,49,52	0
3	SIA	E	3	20/21	-	-	40,48,59,62	0
3	FUC	E	4	10/11	-	-	39,44,47,53	0
3	NAG	F	1	15/15	-	-	31,36,43,48	0
3	GAL	F	2	11/12	-	-	37,39,45,49	0
3	SIA	F	3	20/21	-	-	36,44,51,52	0
3	FUC	F	4	10/11	-	-	31,36,38,38	0
3	SIA	G	3	20/21	0.93	0.09	36,43,49,50	0
3	NAG	G	1	15/15	0.94	0.08	31,40,45,47	0
3	FUC	G	4	10/11	0.95	0.09	32,38,39,40	0
3	GAL	G	2	11/12	0.96	0.06	36,40,45,45	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain E:


$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain F:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	B	302	6/6	0.80	0.16	52,53,58,62	0
4	GOL	B	301	6/6	0.81	0.14	49,60,66,69	0
4	GOL	D	301	6/6	0.87	0.13	55,58,63,67	0
4	GOL	C	305	6/6	0.94	0.10	38,48,51,54	0

6.5 Other polymers [i](#)

There are no such residues in this entry.