



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 25, 2026 – 08:49 AM EDT

PDB ID : 4XVW / pdb_00004xvw
Title : Crystal structure of Proteus mirabilis ScsC in a compact conformation
Authors : Kurth, F.; Furlong, E.J.; Premkumar, L.; Martin, J.L.
Deposited on : 2015-01-27
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	224	 78% 20% ..
1	B	224	 83% 14% .
1	C	224	 87% 11% ..
1	D	224	 84% 14% .
1	E	224	 67% 27% . .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	224	 78%19%..
1	G	224	 62%29%7%.
1	H	224	 80%17%..
1	I	224	 64%22%.12%
1	J	224	 69%24%..
1	K	224	 84%14%.
1	L	224	 54%22%.21%
1	M	224	 69%25%..
1	N	224	 71%25%..
1	O	224	 71%27%.
1	P	224	 71%25%..
1	Q	224	 65%29%..
1	R	224	 71%25%..
1	T	224	 87%11%..
1	U	224	 83%16%..
1	V	224	 76%22%..
1	W	224	 70%26%..
1	X	224	 59%34%5%.
1	Y	224	 79%17%..

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 83204 atoms, of which 42074 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DsbA-like protein.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
1	A	222	Total	C	H	N	O	S	Se	0	3	0
			3527	1095	1790	295	339	4	4			
1	B	220	Total	C	H	N	O	S	Se	0	5	0
			3500	1088	1774	292	337	4	5			
1	C	222	Total	C	H	N	O	S	Se	0	3	0
			3527	1095	1790	295	339	4	4			
1	D	222	Total	C	H	N	O	S	Se	0	3	0
			3527	1095	1790	295	339	4	4			
1	E	215	Total	C	H	N	O	S	Se	0	3	0
			3437	1068	1745	286	330	4	4			
1	F	222	Total	C	H	N	O	S	Se	0	3	0
			3527	1095	1790	295	339	4	4			
1	G	219	Total	C	H	N	O	S	Se	0	3	0
			3488	1083	1769	292	336	4	4			
1	H	222	Total	C	H	N	O	S	Se	0	3	0
			3527	1095	1790	295	339	4	4			
1	I	198	Total	C	H	N	O	S	Se	0	0	0
			3118	972	1579	261	300	2	4			
1	J	219	Total	C	H	N	O	S	Se	0	0	0
			3451	1074	1750	288	333	2	4			
1	K	220	Total	C	H	N	O	S	Se	0	5	0
			3500	1088	1774	292	337	4	5			
1	L	176	Total	C	H	N	O	S	Se	0	0	0
			2800	874	1428	224	268	2	4			
1	M	220	Total	C	H	N	O	S	Se	0	3	0
			3482	1083	1764	291	336	4	4			
1	N	220	Total	C	H	N	O	S	Se	0	0	0
			3473	1080	1763	290	334	2	4			
1	O	220	Total	C	H	N	O	S	Se	0	3	0
			3482	1083	1764	291	336	4	4			
1	P	222	Total	C	H	N	O	S	Se	0	0	0
			3506	1089	1781	293	337	2	4			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
1	Q	221	Total	C	H	N	O	S	Se	0	3	0
			3504	1089	1777	293	337	4	4			
1	R	218	Total	C	H	N	O	S	Se	0	3	0
			3474	1079	1763	290	334	4	4			
1	T	221	Total	C	H	N	O	S	Se	0	3	0
			3504	1089	1777	293	337	4	4			
1	U	222	Total	C	H	N	O	S	Se	0	3	0
			3527	1095	1790	295	339	4	4			
1	V	221	Total	C	H	N	O	S	Se	0	3	0
			3504	1089	1777	293	337	4	4			
1	W	221	Total	C	H	N	O	S	Se	0	3	0
			3504	1089	1777	293	337	4	4			
1	X	222	Total	C	H	N	O	S	Se	0	0	0
			3507	1089	1782	293	337	2	4			
1	Y	222	Total	C	H	N	O	S	Se	0	3	0
			3527	1095	1790	295	339	4	4			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	SER	-	expression tag	UNP C2LPE2
A	2	ASN	-	expression tag	UNP C2LPE2
B	1	SER	-	expression tag	UNP C2LPE2
B	2	ASN	-	expression tag	UNP C2LPE2
C	1	SER	-	expression tag	UNP C2LPE2
C	2	ASN	-	expression tag	UNP C2LPE2
D	1	SER	-	expression tag	UNP C2LPE2
D	2	ASN	-	expression tag	UNP C2LPE2
E	1	SER	-	expression tag	UNP C2LPE2
E	2	ASN	-	expression tag	UNP C2LPE2
F	1	SER	-	expression tag	UNP C2LPE2
F	2	ASN	-	expression tag	UNP C2LPE2
G	1	SER	-	expression tag	UNP C2LPE2
G	2	ASN	-	expression tag	UNP C2LPE2
H	1	SER	-	expression tag	UNP C2LPE2
H	2	ASN	-	expression tag	UNP C2LPE2
I	1	SER	-	expression tag	UNP C2LPE2
I	2	ASN	-	expression tag	UNP C2LPE2
J	1	SER	-	expression tag	UNP C2LPE2
J	2	ASN	-	expression tag	UNP C2LPE2
K	1	SER	-	expression tag	UNP C2LPE2
K	2	ASN	-	expression tag	UNP C2LPE2
L	1	SER	-	expression tag	UNP C2LPE2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
L	2	ASN	-	expression tag	UNP C2LPE2
M	1	SER	-	expression tag	UNP C2LPE2
M	2	ASN	-	expression tag	UNP C2LPE2
N	1	SER	-	expression tag	UNP C2LPE2
N	2	ASN	-	expression tag	UNP C2LPE2
O	1	SER	-	expression tag	UNP C2LPE2
O	2	ASN	-	expression tag	UNP C2LPE2
P	1	SER	-	expression tag	UNP C2LPE2
P	2	ASN	-	expression tag	UNP C2LPE2
Q	1	SER	-	expression tag	UNP C2LPE2
Q	2	ASN	-	expression tag	UNP C2LPE2
R	1	SER	-	expression tag	UNP C2LPE2
R	2	ASN	-	expression tag	UNP C2LPE2
T	1	SER	-	expression tag	UNP C2LPE2
T	2	ASN	-	expression tag	UNP C2LPE2
U	1	SER	-	expression tag	UNP C2LPE2
U	2	ASN	-	expression tag	UNP C2LPE2
V	1	SER	-	expression tag	UNP C2LPE2
V	2	ASN	-	expression tag	UNP C2LPE2
W	1	SER	-	expression tag	UNP C2LPE2
W	2	ASN	-	expression tag	UNP C2LPE2
X	1	SER	-	expression tag	UNP C2LPE2
X	2	ASN	-	expression tag	UNP C2LPE2
Y	1	SER	-	expression tag	UNP C2LPE2
Y	2	ASN	-	expression tag	UNP C2LPE2

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	7	Total O 7 7	0	0
2	B	10	Total O 10 10	0	0
2	C	18	Total O 18 18	0	0
2	D	11	Total O 11 11	0	0
2	E	14	Total O 14 14	0	0
2	F	17	Total O 17 17	0	0
2	G	17	Total O 17 17	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	H	13	Total 13	O 13	0	0
2	I	13	Total 13	O 13	0	0
2	J	8	Total 8	O 8	0	0
2	K	7	Total 7	O 7	0	0
2	L	14	Total 14	O 14	0	0
2	M	11	Total 11	O 11	0	0
2	N	11	Total 11	O 11	0	0
2	O	19	Total 19	O 19	0	0
2	P	16	Total 16	O 16	0	0
2	Q	17	Total 17	O 17	0	0
2	R	9	Total 9	O 9	0	0
2	T	10	Total 10	O 10	0	0
2	U	6	Total 6	O 6	0	0
2	V	9	Total 9	O 9	0	0
2	W	6	Total 6	O 6	0	0
2	X	9	Total 9	O 9	0	0
2	Y	9	Total 9	O 9	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DsbA-like protein




- Molecule 1: DsbA-like protein



- Molecule 1: DsbA-like protein




- Molecule 1: DsbA-like protein




- Molecule 1: DsbA-like protein

Chain E:  67% 27% ..




- Molecule 1: DsbA-like protein

Chain F:  78% 19% ..




- Molecule 1: DsbA-like protein

Chain G:  62% 29% 7% .




- Molecule 1: DsbA-like protein

Chain H:  80% 17% ..




- Molecule 1: DsbA-like protein

Chain I:  64% 22% 12% .



- Molecule 1: DsbA-like protein

Chain J: 69% 24% . . .




- Molecule 1: DsbA-like protein

Chain K: 84% 14% .




- Molecule 1: DsbA-like protein

Chain L: 54% 22% . 21%




- Molecule 1: DsbA-like protein

Chain M: 69% 25% . . .




- Molecule 1: DsbA-like protein

Chain N:  71% 25%




• Molecule 1: DsbA-like protein

Chain O:  71% 27%




• Molecule 1: DsbA-like protein

Chain P:  71% 25%




• Molecule 1: DsbA-like protein

Chain Q:  65% 29%



• Molecule 1: DsbA-like protein

Chain R:  71% 25%



• Molecule 1: DsbA-like protein

Chain T:  87% 11% ..




- Molecule 1: DsbA-like protein

Chain U:  83% 16% ..




- Molecule 1: DsbA-like protein

Chain V:  76% 22% ..



- Molecule 1: DsbA-like protein

Chain W:  70% 26% ..



- Molecule 1: DsbA-like protein

Chain X:  59% 34% 5% .




- Molecule 1: DsbA-like protein

Chain Y:

79%

17%

••



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	137.47Å 163.88Å 181.94Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	91.15 – 2.60 91.15 – 2.60	Depositor EDS
% Data completeness (in resolution range)	98.5 (91.15-2.60) 98.5 (91.15-2.60)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.16 (at 2.61Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.248 , 0.282 0.245 , 0.276	Depositor DCC
R_{free} test set	12080 reflections (4.33%)	wwPDB-VP
Wilson B-factor (Å ²)	45.0	Xtriage
Anisotropy	0.635	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 42.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.40$, $\langle L^2 \rangle = 0.22$	Xtriage
Estimated twinning fraction	0.467 for h,-k,-l	Xtriage
Reported twinning fraction	0.500 for H, K, L 0.500 for h,-k,-l	Depositor
Outliers	0 of 243406 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	83204	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 31.03 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.1846e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/1765	0.86	7/2377 (0.3%)
1	B	0.36	0/1762	0.78	1/2376 (0.0%)
1	C	0.38	0/1765	0.81	0/2377
1	D	0.36	0/1765	0.78	1/2377 (0.0%)
1	E	0.42	0/1720	0.91	4/2315 (0.2%)
1	F	0.38	0/1765	0.88	5/2377 (0.2%)
1	G	0.48	0/1747	1.06	14/2352 (0.6%)
1	H	0.40	0/1765	0.90	5/2377 (0.2%)
1	I	0.41	0/1558	0.99	3/2101 (0.1%)
1	J	0.43	0/1720	0.98	13/2320 (0.6%)
1	K	0.37	0/1763	0.77	0/2377
1	L	0.42	0/1386	0.88	5/1867 (0.3%)
1	M	0.44	0/1746	0.98	6/2355 (0.3%)
1	N	0.43	0/1729	0.98	8/2331 (0.3%)
1	O	0.45	0/1746	1.00	4/2355 (0.2%)
1	P	0.40	0/1744	0.96	9/2349 (0.4%)
1	Q	0.48	0/1755	1.03	7/2366 (0.3%)
1	R	0.42	0/1739	0.93	2/2341 (0.1%)
1	T	0.34	0/1755	0.78	1/2366 (0.0%)
1	U	0.38	0/1765	0.81	2/2377 (0.1%)
1	V	0.38	0/1755	0.92	3/2366 (0.1%)
1	W	0.41	0/1755	0.98	6/2366 (0.3%)
1	X	0.47	0/1744	1.07	9/2349 (0.4%)
1	Y	0.40	0/1765	0.87	2/2377 (0.1%)
All	All	0.41	0/41479	0.92	117/55891 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	1

Continued on next page...

Continued from previous page...

Mol	Chain	#Chirality outliers	#Planarity outliers
1	F	0	1
1	J	0	2
1	Q	0	1
1	X	0	1
All	All	0	6

There are no bond length outliers.

The worst 5 of 117 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	W	47	PHE	N-CA-C	13.59	129.35	112.23
1	O	42	GLU	N-CA-C	-10.46	99.69	111.82
1	V	100	GLN	N-CA-C	-10.42	100.00	111.36
1	X	100	GLN	N-CA-C	-8.94	101.45	111.82
1	O	54	GLU	N-CA-C	-8.69	102.75	112.57

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	47	PHE	Peptide
1	F	99	GLU	Peptide
1	J	140	HIS	Peptide
1	J	144	MSE	Peptide
1	Q	149	MSE	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1737	1790	1776	32	0
1	B	1726	1774	1748	26	0
1	C	1737	1790	1776	17	2
1	D	1737	1790	1776	22	0
1	E	1692	1745	1731	55	0
1	F	1737	1790	1776	31	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	1719	1769	1755	69	2
1	H	1737	1790	1776	30	2
1	I	1539	1579	1579	37	0
1	J	1701	1750	1750	40	2
1	K	1726	1774	1751	23	0
1	L	1372	1428	1428	32	0
1	M	1718	1764	1750	52	1
1	N	1710	1763	1763	44	0
1	O	1718	1764	1750	54	0
1	P	1725	1781	1781	47	0
1	Q	1727	1777	1763	56	0
1	R	1711	1763	1749	42	0
1	T	1727	1777	1763	20	0
1	U	1737	1790	1776	25	1
1	V	1727	1777	1763	33	0
1	W	1727	1777	1763	41	0
1	X	1725	1782	1781	69	0
1	Y	1737	1790	1776	46	0
2	A	7	0	0	0	0
2	B	10	0	0	0	0
2	C	18	0	0	0	0
2	D	11	0	0	0	0
2	E	14	0	0	0	0
2	F	17	0	0	0	0
2	G	17	0	0	2	0
2	H	13	0	0	0	0
2	I	13	0	0	2	0
2	J	8	0	0	0	0
2	K	7	0	0	0	0
2	L	14	0	0	0	0
2	M	11	0	0	0	0
2	N	11	0	0	0	0
2	O	19	0	0	1	0
2	P	16	0	0	1	0
2	Q	17	0	0	1	0
2	R	9	0	0	1	0
2	T	10	0	0	0	0
2	U	6	0	0	0	0
2	V	9	0	0	0	0
2	W	6	0	0	0	0
2	X	9	0	0	1	0
2	Y	9	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	41130	42074	41800	813	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 813 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:O:124:ALA:HA	1:O:169:LEU:HD21	1.57	0.86
1:J:100:GLN:OE1	1:J:208:TYR:OH	1.97	0.83
1:I:95:GLU:OE2	1:I:141:GLN:NE2	2.11	0.83
1:O:14:ARG:NH2	1:P:23:SER:OG	2.14	0.81
1:M:57:ALA:O	1:M:109:LYS:NZ	2.13	0.81

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:161:SER:OG	1:H:99:GLU:OE2[1_655]	1.95	0.25
1:H:86:TYR:OH	1:J:187:LEU:O[1_455]	2.12	0.08
1:G:134:LYS:NZ	1:M:134:LYS:O[2_456]	2.13	0.07
1:G:148:THR:OG1	1:J:148:THR:OG1[1_455]	2.17	0.03
1:C:7:ALA:H	1:U:3:ALA:O[1_554]	1.59	0.01

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	223/224 (100%)	212 (95%)	8 (4%)	3 (1%)	9 21
1	B	223/224 (100%)	216 (97%)	6 (3%)	1 (0%)	30 51

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	223/224 (100%)	214 (96%)	8 (4%)	1 (0%)	30	51
1	D	223/224 (100%)	216 (97%)	7 (3%)	0	100	100
1	E	216/224 (96%)	203 (94%)	12 (6%)	1 (0%)	24	46
1	F	223/224 (100%)	214 (96%)	7 (3%)	2 (1%)	14	30
1	G	220/224 (98%)	208 (94%)	10 (4%)	2 (1%)	14	30
1	H	223/224 (100%)	212 (95%)	9 (4%)	2 (1%)	14	30
1	I	196/224 (88%)	188 (96%)	6 (3%)	2 (1%)	12	28
1	J	217/224 (97%)	210 (97%)	5 (2%)	2 (1%)	14	30
1	K	223/224 (100%)	215 (96%)	8 (4%)	0	100	100
1	L	172/224 (77%)	162 (94%)	6 (4%)	4 (2%)	5	9
1	M	221/224 (99%)	210 (95%)	10 (4%)	1 (0%)	24	46
1	N	218/224 (97%)	210 (96%)	7 (3%)	1 (0%)	24	46
1	O	221/224 (99%)	213 (96%)	7 (3%)	1 (0%)	24	46
1	P	220/224 (98%)	210 (96%)	8 (4%)	2 (1%)	14	30
1	Q	222/224 (99%)	198 (89%)	20 (9%)	4 (2%)	6	14
1	R	219/224 (98%)	208 (95%)	7 (3%)	4 (2%)	6	14
1	T	222/224 (99%)	216 (97%)	5 (2%)	1 (0%)	24	46
1	U	223/224 (100%)	214 (96%)	8 (4%)	1 (0%)	30	51
1	V	222/224 (99%)	211 (95%)	9 (4%)	2 (1%)	14	30
1	W	222/224 (99%)	211 (95%)	10 (4%)	1 (0%)	24	46
1	X	220/224 (98%)	196 (89%)	20 (9%)	4 (2%)	6	14
1	Y	223/224 (100%)	206 (92%)	10 (4%)	7 (3%)	3	5
All	All	5235/5376 (97%)	4973 (95%)	213 (4%)	49 (1%)	14	30

5 of 49 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	223	LYS
1	F	148	THR
1	I	148	THR
1	L	145	GLN
1	N	148	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	192/187 (103%)	191 (100%)	1 (0%)	81	92
1	B	192/187 (103%)	190 (99%)	2 (1%)	68	86
1	C	192/187 (103%)	186 (97%)	6 (3%)	35	63
1	D	192/187 (103%)	189 (98%)	3 (2%)	55	79
1	E	189/187 (101%)	186 (98%)	3 (2%)	55	79
1	F	192/187 (103%)	186 (97%)	6 (3%)	35	63
1	G	191/187 (102%)	187 (98%)	4 (2%)	47	73
1	H	192/187 (103%)	187 (97%)	5 (3%)	40	68
1	I	170/187 (91%)	166 (98%)	4 (2%)	43	70
1	J	187/187 (100%)	178 (95%)	9 (5%)	23	47
1	K	192/187 (103%)	188 (98%)	4 (2%)	47	73
1	L	156/187 (83%)	150 (96%)	6 (4%)	29	56
1	M	190/187 (102%)	183 (96%)	7 (4%)	30	57
1	N	188/187 (100%)	183 (97%)	5 (3%)	39	67
1	O	190/187 (102%)	189 (100%)	1 (0%)	81	92
1	P	189/187 (101%)	186 (98%)	3 (2%)	55	79
1	Q	191/187 (102%)	185 (97%)	6 (3%)	35	63
1	R	190/187 (102%)	185 (97%)	5 (3%)	40	68
1	T	191/187 (102%)	187 (98%)	4 (2%)	47	73
1	U	192/187 (103%)	191 (100%)	1 (0%)	81	92
1	V	191/187 (102%)	187 (98%)	4 (2%)	47	73
1	W	191/187 (102%)	182 (95%)	9 (5%)	23	48
1	X	189/187 (101%)	182 (96%)	7 (4%)	30	57
1	Y	192/187 (103%)	189 (98%)	3 (2%)	55	79
All	All	4521/4488 (101%)	4413 (98%)	108 (2%)	43	70

5 of 108 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	M	134	LYS
1	Q	149	MSE
1	X	97	ILE
1	M	218	GLN
1	O	46	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 46 such sidechains are listed below:

Mol	Chain	Res	Type
1	R	36	GLN
1	V	178	GLN
1	R	163	ASN
1	T	49	GLN
1	W	60	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	218/224 (97%)	-1.39	0 100 100	17, 40, 60, 84	2 (0%)
1	B	216/224 (96%)	-1.34	0 100 100	23, 44, 62, 89	2 (0%)
1	C	218/224 (97%)	-1.29	0 100 100	22, 49, 65, 80	2 (0%)
1	D	218/224 (97%)	-1.30	0 100 100	19, 46, 62, 72	2 (0%)
1	E	211/224 (94%)	-1.14	0 100 100	31, 67, 86, 100	2 (0%)
1	F	218/224 (97%)	-1.31	0 100 100	21, 46, 69, 89	2 (0%)
1	G	215/224 (95%)	-1.06	0 100 100	28, 67, 88, 104	2 (0%)
1	H	218/224 (97%)	-1.09	0 100 100	28, 62, 90, 102	2 (0%)
1	I	194/224 (86%)	-1.06	0 100 100	40, 67, 86, 98	0
1	J	215/224 (95%)	-1.07	0 100 100	42, 67, 93, 114	0
1	K	216/224 (96%)	-1.35	0 100 100	21, 44, 69, 86	2 (0%)
1	L	172/224 (76%)	-1.09	0 100 100	38, 62, 79, 93	0
1	M	216/224 (96%)	-1.04	0 100 100	27, 69, 93, 124	2 (0%)
1	N	216/224 (96%)	-0.99	0 100 100	36, 71, 94, 123	0
1	O	216/224 (96%)	-1.08	0 100 100	32, 66, 94, 110	2 (0%)
1	P	218/224 (97%)	-1.08	0 100 100	40, 70, 91, 100	0
1	Q	217/224 (96%)	-1.07	0 100 100	28, 67, 97, 107	2 (0%)
1	R	214/224 (95%)	-1.11	0 100 100	29, 68, 90, 104	2 (0%)
1	T	217/224 (96%)	-1.26	0 100 100	25, 48, 68, 81	2 (0%)
1	U	218/224 (97%)	-1.25	0 100 100	25, 49, 70, 95	2 (0%)
1	V	217/224 (96%)	-1.13	0 100 100	31, 59, 84, 98	2 (0%)
1	W	217/224 (96%)	-1.06	0 100 100	42, 66, 89, 122	2 (0%)
1	X	218/224 (97%)	-1.03	0 100 100	32, 68, 91, 109	0
1	Y	218/224 (97%)	-1.23	0 100 100	27, 49, 73, 87	2 (0%)

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
All	All	5131/5376 (95%)	-1.16	0 100 100	17, 59, 87, 124	36 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.