



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 06:13 PM UTC

PDB ID : 5XWB / pdb_00005xwb
Title : Crystal Structure of 5-Enolpyruvulshikimate-3-phosphate Synthase from a Psychrophilic Bacterium, Colwellia psychrerythraea
Authors : Lee, J.H.; Kim, H.J.; Choi, J.M.; Kim, D.-W.; Seo, Y.-S.
Deposited on : 2017-06-29
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	446	 88% 6% . .
1	B	446	 85% 9% . .

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 6875 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-phosphoshikimate 1-carboxyvinyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	426	Total	C	N	O	S	0	0	0
			3241	2053	537	632	19			
1	B	426	Total	C	N	O	S	0	0	0
			3241	2053	537	632	19			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP Q482G5
A	-18	GLY	-	expression tag	UNP Q482G5
A	-17	SER	-	expression tag	UNP Q482G5
A	-16	SER	-	expression tag	UNP Q482G5
A	-15	HIS	-	expression tag	UNP Q482G5
A	-14	HIS	-	expression tag	UNP Q482G5
A	-13	HIS	-	expression tag	UNP Q482G5
A	-12	HIS	-	expression tag	UNP Q482G5
A	-11	HIS	-	expression tag	UNP Q482G5
A	-10	HIS	-	expression tag	UNP Q482G5
A	-9	SER	-	expression tag	UNP Q482G5
A	-8	SER	-	expression tag	UNP Q482G5
A	-7	GLY	-	expression tag	UNP Q482G5
A	-6	LEU	-	expression tag	UNP Q482G5
A	-5	VAL	-	expression tag	UNP Q482G5
A	-4	PRO	-	expression tag	UNP Q482G5
A	-3	ARG	-	expression tag	UNP Q482G5
A	-2	GLY	-	expression tag	UNP Q482G5
A	-1	SER	-	expression tag	UNP Q482G5
A	0	HIS	-	expression tag	UNP Q482G5
B	-19	MET	-	expression tag	UNP Q482G5
B	-18	GLY	-	expression tag	UNP Q482G5
B	-17	SER	-	expression tag	UNP Q482G5
B	-16	SER	-	expression tag	UNP Q482G5
B	-15	HIS	-	expression tag	UNP Q482G5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-14	HIS	-	expression tag	UNP Q482G5
B	-13	HIS	-	expression tag	UNP Q482G5
B	-12	HIS	-	expression tag	UNP Q482G5
B	-11	HIS	-	expression tag	UNP Q482G5
B	-10	HIS	-	expression tag	UNP Q482G5
B	-9	SER	-	expression tag	UNP Q482G5
B	-8	SER	-	expression tag	UNP Q482G5
B	-7	GLY	-	expression tag	UNP Q482G5
B	-6	LEU	-	expression tag	UNP Q482G5
B	-5	VAL	-	expression tag	UNP Q482G5
B	-4	PRO	-	expression tag	UNP Q482G5
B	-3	ARG	-	expression tag	UNP Q482G5
B	-2	GLY	-	expression tag	UNP Q482G5
B	-1	SER	-	expression tag	UNP Q482G5
B	0	HIS	-	expression tag	UNP Q482G5


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	201	Total O 201 201	0	0
2	B	192	Total O 192 192	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 3-phosphoshikimate 1-carboxyvinyltransferase



- Molecule 1: 3-phosphoshikimate 1-carboxyvinyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	61.80Å 80.03Å 82.27Å 90.00° 96.46° 90.00°	Depositor
Resolution (Å)	50.00 – 2.20 50.00 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.7 (50.00-2.20) 99.7 (50.00-2.20)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	8.69 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.146 , 0.210 0.157 , 0.215	Depositor DCC
R_{free} test set	2024 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	29.1	Xtriage
Anisotropy	0.082	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 40.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6875	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.10	3/3295 (0.1%)	1.04	2/4469 (0.0%)
1	B	1.12	3/3295 (0.1%)	1.06	5/4469 (0.1%)
All	All	1.11	6/6590 (0.1%)	1.05	7/8938 (0.1%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	170	SER	N-CA	6.23	1.54	1.46
1	B	263	VAL	C-O	-5.61	1.18	1.24
1	A	37	GLY	N-CA	5.31	1.51	1.45
1	A	371	PRO	C-O	-5.22	1.20	1.24
1	B	115	LEU	N-CA	-5.21	1.39	1.46
1	A	406	ASP	C-O	5.16	1.26	1.23

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	172	GLN	N-CA-C	-6.73	104.69	113.17
1	B	414	PHE	N-CA-CB	-5.76	104.20	110.71
1	B	27	ARG	CG-CD-NE	5.31	123.69	112.00
1	A	179	MET	N-CA-C	5.21	116.96	111.28
1	B	413	THR	CA-C-N	-5.09	117.64	122.37
1	B	413	THR	C-N-CA	-5.09	117.64	122.37
1	A	26	ASN	N-CA-C	5.02	116.83	111.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3241	0	3279	12	1
1	B	3241	0	3279	24	1
2	A	201	0	0	0	0
2	B	192	0	0	2	0
All	All	6875	0	6558	34	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:280:VAL:HG22	1:B:307:MET:HE1	1.77	0.66
1:B:310:ASN:ND2	2:B:503:HOH:O	2.29	0.65
1:A:168:SER:O	1:A:169:ILE:HG22	1.97	0.64
1:B:129:LEU:HB2	1:B:173:PHE:CD1	2.33	0.64
1:B:128:HIS:HB2	1:B:173:PHE:CE2	2.33	0.62
1:A:55:ASN:HD21	1:B:307:MET:HE3	1.66	0.60
1:A:31:ILE:HD11	1:A:179:MET:HE3	1.86	0.58
1:B:97:THR:HG22	1:B:172:GLN:HG3	1.87	0.56
1:B:375:LEU:HD12	1:B:398:SER:CB	2.35	0.56
1:A:171:SER:OG	1:A:198:SER:HB2	2.08	0.54
1:A:94:ASN:HD21	1:A:120:ARG:HD3	1.75	0.51
1:B:280:VAL:HG22	1:B:307:MET:CE	2.39	0.51
1:B:375:LEU:HD12	1:B:398:SER:HB3	1.92	0.50
1:B:272:GLN:O	1:B:275:LYS:HG2	2.14	0.48
1:B:100:ARG:HD2	1:B:172:GLN:HB3	1.96	0.47
1:B:129:LEU:HB2	1:B:173:PHE:HD1	1.78	0.47
1:A:243:ASP:OD2	1:A:245:SER:HB2	2.16	0.46
1:A:341:LYS:HG3	1:A:345:ARG:HD2	1.98	0.45
1:B:100:ARG:HB2	1:B:101:PRO:HD3	2.00	0.44
1:B:181:ALA:N	1:B:182:PRO:CD	2.80	0.44
1:B:233:GLN:HG2	2:B:595:HOH:O	2.18	0.43
1:A:165:ILE:HD13	1:A:165:ILE:HA	1.86	0.42
1:B:92:LEU:HD11	1:B:102:LEU:HD12	2.00	0.42
1:B:268:LYS:HG3	1:B:292:ASP:HA	2.01	0.42
1:B:18:LEU:HD11	1:B:247:ALA:HB2	2.00	0.42
1:B:31:ILE:HG13	1:B:179:MET:HE3	2.01	0.42
1:A:55:ASN:ND2	1:B:307:MET:HE3	2.33	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:31:ILE:CD1	1:A:179:MET:HE3	2.49	0.41
1:A:288:ILE:HG23	1:A:295:ILE:HD11	2.02	0.41
1:B:116:THR:HB	1:B:150:PRO:HB3	2.02	0.41
1:A:323:ALA:O	1:A:371:PRO:HG3	2.21	0.41
1:B:416:ASP:OD1	1:B:416:ASP:N	2.54	0.40
1:B:172:GLN:O	1:B:172:GLN:OE1	2.40	0.40
1:B:306:ASP:C	1:B:306:ASP:OD1	2.65	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:233:GLN:OE1	1:B:36:ASN:ND2[2_546]	1.93	0.27

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	424/446 (95%)	412 (97%)	12 (3%)	0	100	100
1	B	424/446 (95%)	409 (96%)	13 (3%)	2 (0%)	24	27
All	All	848/892 (95%)	821 (97%)	25 (3%)	2 (0%)	43	51

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	171	SER
1	B	412	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	355/372 (95%)	342 (96%)	13 (4%)	30	41
1	B	355/372 (95%)	346 (98%)	9 (2%)	42	56
All	All	710/744 (95%)	688 (97%)	22 (3%)	35	48

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	SER
1	A	120	ARG
1	A	124	ARG
1	A	165	ILE
1	A	172	GLN
1	A	175	THR
1	A	215	SER
1	A	231	SER
1	A	283	LYS
1	A	295	ILE
1	A	342	GLU
1	A	412	LYS
1	A	426	CYS
1	B	68	ASP
1	B	69	CYS
1	B	99	MET
1	B	126	ILE
1	B	163	VAL
1	B	214	VAL
1	B	268	LYS
1	B	416	ASP
1	B	426	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	55	ASN
1	A	276	HIS
1	A	310	ASN
1	A	337	ASN
1	B	3	GLN
1	B	13	ASN
1	B	83	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	426/446 (95%)	-0.47	2 (0%) 87 85	17, 28, 52, 85	0
1	B	426/446 (95%)	-0.47	5 (1%) 76 74	19, 28, 49, 81	0
All	All	852/892 (95%)	-0.47	7 (0%) 82 80	17, 28, 51, 85	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	173	PHE	3.5
1	A	169	ILE	2.9
1	B	170	SER	2.6
1	A	1	MET	2.4
1	B	171	SER	2.4
1	B	169	ILE	2.3
1	B	172	GLN	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.