



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 7, 2026 – 01:30 AM UTC

PDB ID : 5XWP / pdb_00005xwp
Title : Crystal structure of LbuCas13a-crRNA-target RNA ternary complex
Authors : Liu, L.; Li, X.; Li, Z.; Wang, Y.
Deposited on : 2017-06-30
Resolution : 3.09 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2010 (3.10-3.06)
Clashscore	190562	2102 (3.10-3.06)
Ramachandran outliers	187476	1982 (3.10-3.06)
Sidechain outliers	187428	1981 (3.10-3.06)
RSRZ outliers	180081	2010 (3.10-3.06)
RNA backbone	3983	1054 (3.30-2.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1160	43% 41% 11% . .
1	B	1160	42% 42% 12% . .
2	C	59	20% 39% 31% 8% .
2	E	59	25% 41% 24% 8% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	D	30	23%53%23%
3	F	30	33%33%30%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 21838 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uncharacterized protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	1125	Total	C	N	O	S	Se	0	0	0
			9160	5877	1539	1720	3	21			
1	B	1114	Total	C	N	O	S	Se	0	0	0
			8940	5718	1508	1690	3	21			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP C7NBY4
A	1048	ALA	ARG	engineered mutation	UNP C7NBY4
A	1053	ALA	HIS	engineered mutation	UNP C7NBY4
B	0	SER	-	expression tag	UNP C7NBY4
B	1048	ALA	ARG	engineered mutation	UNP C7NBY4
B	1053	ALA	HIS	engineered mutation	UNP C7NBY4

- Molecule 2 is a RNA chain called RNA (59-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	58	Total	C	N	O	P	0	0	0
			1229	553	224	394	58			
2	E	58	Total	C	N	O	P	0	0	0
			1229	553	224	394	58			

- Molecule 3 is a RNA chain called RNA (30-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	30	Total	C	N	O	P	0	0	0
			644	288	116	210	30			
3	F	29	Total	C	N	O	P	0	0	0
			624	279	113	203	29			


- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	10	Total 10	O 10	0	0
4	B	2	Total 2	O 2	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Uncharacterized protein






• Molecule 1: Uncharacterized protein






• Molecule 2: RNA (59-MER)

Chain C: 20% 39% 31% 8% .



• Molecule 2: RNA (59-MER)

Chain E: 25% 41% 24% 8% .




• Molecule 3: RNA (30-MER)

Chain D: 23% 53% 23%



• Molecule 3: RNA (30-MER)

Chain F: 33% 33% 30%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	95.58Å 132.87Å 139.96Å 90.00° 90.80° 90.00°	Depositor
Resolution (Å)	48.18 – 3.09 48.18 – 3.09	Depositor EDS
% Data completeness (in resolution range)	95.1 (48.18-3.09) 91.3 (48.18-3.09)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.18 (at 3.07Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155)	Depositor
R, R_{free}	0.218 , 0.244 0.219 , 0.242	Depositor DCC
R_{free} test set	2881 reflections (4.47%)	wwPDB-VP
Wilson B-factor (Å ²)	58.5	Xtriage
Anisotropy	0.074	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.25 , 28.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.36$, $\langle L^2 \rangle = 0.18$	Xtriage
Estimated twinning fraction	0.186 for h,-k,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	21838	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.98	1/9290 (0.0%)	1.08	60/12431 (0.5%)
1	B	0.94	2/9063 (0.0%)	1.15	66/12146 (0.5%)
2	C	0.73	1/1375 (0.1%)	0.95	4/2137 (0.2%)
2	E	0.67	1/1375 (0.1%)	0.89	6/2137 (0.3%)
3	D	0.63	0/721	0.79	0/1122
3	F	0.66	0/699	0.77	1/1088 (0.1%)
All	All	0.91	5/22523 (0.0%)	1.07	137/31061 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
All	All	0	4

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	2	A	O3'-P	-7.93	1.49	1.61
2	C	2	A	O3'-P	-7.34	1.50	1.61
1	A	494	ALA	C-N	5.31	1.40	1.33
1	B	406	MSE	CA-C	-5.25	1.45	1.52
1	B	679	ASP	CA-C	-5.20	1.45	1.52

The worst 5 of 137 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	403	THR	N-CA-C	-24.97	70.75	110.20
1	B	570	LEU	N-CA-C	20.72	144.44	114.39
1	A	163	ILE	N-CA-C	14.63	125.50	110.72

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	404	GLY	CA-C-N	-12.61	104.98	122.77
1	B	404	GLY	C-N-CA	-12.61	104.98	122.77

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1025	VAL	Peptide
1	A	705	GLY	Peptide
1	B	421	VAL	Peptide
1	B	695	ASN	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	9160	0	9025	697	0
1	B	8940	0	8665	699	0
2	C	1229	0	627	87	0
2	E	1229	0	627	95	0
3	D	644	0	321	40	0
3	F	624	0	310	36	0
4	A	10	0	0	0	0
4	B	2	0	0	0	0
All	All	21838	0	19575	1580	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 38.

The worst 5 of 1580 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:566:LEU:HD12	1:A:686:LEU:CD1	1.58	1.30
1:A:406:MSE:CE	1:A:461:PHE:HB3	1.62	1.30
1:B:920:PHE:O	1:B:921:THR:CG2	1.82	1.24

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:265:SER:HB3	1:A:268:LYS:HG2	1.24	1.19
1:B:846:ILE:HD11	1:B:887:LEU:HD21	1.21	1.15

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1115/1160 (96%)	1079 (97%)	34 (3%)	2 (0%)	43	71
1	B	1104/1160 (95%)	1064 (96%)	38 (3%)	2 (0%)	43	71
All	All	2219/2320 (96%)	2143 (97%)	72 (3%)	4 (0%)	43	71

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	77	ASP
1	B	921	THR
1	B	1024	GLU
1	A	914	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	967/1061 (91%)	778 (80%)	189 (20%)	1 6
1	B	925/1061 (87%)	747 (81%)	178 (19%)	1 6
All	All	1892/2122 (89%)	1525 (81%)	367 (19%)	1 6

5 of 367 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	231	ILE
1	B	562	ARG
1	B	286	LYS
1	B	439	GLU
1	B	684	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 57 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	57	ASN
1	B	997	ASN
1	B	271	GLN
1	B	987	GLN
1	B	856	HIS

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	C	58/59 (98%)	27 (46%)	5 (8%)
2	E	57/59 (96%)	22 (38%)	4 (7%)
3	D	29/30 (96%)	10 (34%)	0
3	F	28/30 (93%)	11 (39%)	2 (7%)
All	All	172/178 (96%)	70 (40%)	11 (6%)

5 of 70 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	C	2	A
2	C	3	C
2	C	4	C

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	C	5	A
2	C	7	C

5 of 11 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	E	31	A
2	E	32	C
3	F	4	A
3	F	2	G
2	C	30	C

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1104/1160 (95%)	-1.30	0 100 100	17, 43, 82, 138	0
1	B	1093/1160 (94%)	-1.19	2 (0%) 91 83	20, 51, 113, 149	0
2	C	58/59 (98%)	-1.84	0 100 100	22, 36, 68, 81	0
2	E	58/59 (98%)	-1.77	0 100 100	29, 43, 82, 92	0
3	D	30/30 (100%)	-1.95	0 100 100	26, 36, 75, 105	0
3	F	29/30 (96%)	-1.96	0 100 100	30, 40, 82, 122	0
All	All	2372/2498 (94%)	-1.29	2 (0%) 92 86	17, 46, 98, 149	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	575	LYS	2.9
1	B	574	TRP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.