



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 06:49 AM UTC

PDB ID : 6XWS / pdb_00006xws
Title : Crystal Structure of Drosophila CAL1 1-160 bound to CENP-A/H4
Authors : Jeyaprakash, A.A.; Medina-Pritchard, B.; Lazou, V.; Zou, J.; Byron, O.; Abad, M.A.; Rappsilber, J.; Heun, P.
Deposited on : 2020-01-24
Resolution : 4.36 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 4.36 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1078 (4.82-3.90)
Clashscore	190562	1125 (4.82-3.90)
Ramachandran outliers	187476	1020 (4.80-3.90)
Sidechain outliers	187428	1004 (4.80-3.90)
RSRZ outliers	180081	1074 (4.82-3.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	225	
2	B	103	
3	C	108	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3386 atoms, of which 1671 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone H3-like centromeric protein cid.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	69	Total	C	H	N	O	S	0	0	0
			1131	353	573	98	101	6			

- Molecule 2 is a protein called Histone H4.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	68	Total	C	H	N	O	S	0	0	0
			1123	341	583	105	93	1			

- Molecule 3 is a protein called Chromosome alignment defect 1,Chromosome alignment defect 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	105	Total	C	H	N	O	S	0	0	0
			1132	378	515	111	125	3			

4 Data and refinement statistics

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, α , β , γ	199.70Å 199.70Å 76.75Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	172.95 – 4.36 172.94 – 4.36	Depositor EDS
% Data completeness (in resolution range)	89.0 (172.95-4.36) 34.2 (172.94-4.36)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.22 (at 4.30Å)	Xtriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.306 , 0.323 0.305 , 0.331	Depositor DCC
R_{free} test set	117 reflections (1.86%)	wwPDB-VP
Wilson B-factor (Å ²)	180.0	Xtriage
Anisotropy	0.298	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 322.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.81	EDS
Total number of atoms	3386	wwPDB-VP
Average B, all atoms (Å ²)	193.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.27	0/565	0.76	0/759
2	B	0.26	0/545	0.92	1/731 (0.1%)
3	C	0.18	0/315	0.46	0/425
All	All	0.25	0/1425	0.77	1/1915 (0.1%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	77	ALA	N-CA-C	6.25	124.11	110.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	558	573	572	31	0
2	B	540	583	582	29	1
3	C	617	515	340	9	1
All	All	1715	1671	1494	49	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:72:THR:O	2:B:76:HIS:ND1	2.15	0.79
2:B:39:ALA:HB1	2:B:44:VAL:HB	1.67	0.74
2:B:66:VAL:HA	2:B:90:ALA:HB1	1.72	0.71
1:A:70:LEU:HD12	2:B:71:VAL:HG13	1.76	0.68
2:B:74:THR:O	2:B:79:ARG:O	2.12	0.68
1:A:86:MET:HE2	3:C:22:TRP:CD2	2.30	0.67
3:C:29:PHE:O	3:C:31:SER:N	2.28	0.66
2:B:33:PRO:O	2:B:37:ARG:HG3	1.97	0.64
1:A:70:LEU:CD1	2:B:71:VAL:HG13	2.28	0.63
2:B:35:ILE:HD13	2:B:52:TYR:HA	1.80	0.62
1:A:76:ALA:HB1	2:B:82:VAL:HG12	1.84	0.60
1:A:86:MET:HE2	3:C:22:TRP:CE2	2.39	0.57
1:A:72:VAL:HG22	2:B:82:VAL:HB	1.86	0.57
1:A:98:LEU:HD22	1:A:114:LEU:HD23	1.87	0.57
2:B:72:THR:HG23	2:B:76:HIS:CE1	2.40	0.56
1:A:88:LEU:HD11	2:B:59:LEU:HD22	1.88	0.56
3:C:5:VAL:O	3:C:7:ASP:N	2.39	0.55
3:C:31:SER:OG	3:C:32:LEU:N	2.38	0.55
3:C:11:LEU:O	3:C:15:VAL:HG23	2.08	0.53
2:B:65:ASN:OD1	2:B:68:ARG:NH2	2.42	0.53
1:A:84:CYS:HB3	2:B:63:LEU:HD21	1.91	0.52
1:A:84:CYS:CB	2:B:63:LEU:HD21	2.39	0.52
2:B:57:GLY:O	2:B:60:LYS:HG2	2.10	0.52
1:A:71:ARG:O	1:A:72:VAL:HB	2.10	0.51
1:A:56:LEU:HD21	1:A:60:PHE:CE2	2.46	0.51
3:C:34:ASP:O	3:C:35:GLY:C	2.54	0.51
1:A:80:MET:HE1	2:B:67:ILE:HG13	1.94	0.49
1:A:83:SER:HB2	2:B:91:LEU:HD21	1.96	0.48
1:A:71:ARG:O	1:A:72:VAL:CB	2.62	0.47
2:B:37:ARG:O	2:B:41:ARG:HG3	2.15	0.46
1:A:80:MET:HE1	2:B:67:ILE:CD1	2.46	0.46
1:A:88:LEU:HD22	2:B:55:THR:HG23	1.97	0.45
1:A:107:LEU:O	2:B:51:ILE:HD11	2.17	0.44
1:A:80:MET:HE1	2:B:67:ILE:HD11	2.00	0.44
1:A:99:THR:HG22	1:A:104:ARG:O	2.18	0.43
1:A:60:PHE:CE2	2:B:64:GLU:HG3	2.53	0.43
1:A:86:MET:HE2	3:C:22:TRP:CG	2.54	0.43
1:A:105:VAL:O	1:A:105:VAL:HG12	2.18	0.43
1:A:81:GLN:HA	1:A:84:CYS:SG	2.59	0.43
1:A:84:CYS:HB2	2:B:59:LEU:HD11	2.01	0.42
1:A:71:ARG:C	1:A:72:VAL:HG23	2.44	0.42
1:A:94:ASP:OD1	3:C:14:MET:HE2	2.18	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:72:VAL:HG12	1:A:73:THR:N	2.35	0.42
2:B:95:GLY:C	2:B:97:THR:H	2.27	0.42
1:A:94:ASP:O	1:A:97:MET:HB3	2.20	0.42
2:B:95:GLY:HA2	2:B:98:LEU:HG	2.01	0.41
2:B:74:THR:HG21	2:B:81:THR:C	2.46	0.41
1:A:109:VAL:HG12	2:B:54:GLU:HG3	2.03	0.40
1:A:95:SER:O	1:A:111:ASP:HB3	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:89:TYR:OH	3:C:120:UNK:O[11_654]	1.86	0.34

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	67/225 (30%)	62 (92%)	4 (6%)	1 (2%)	8	38
2	B	66/103 (64%)	62 (94%)	4 (6%)	0	100	100
3	C	42/108 (39%)	31 (74%)	8 (19%)	3 (7%)	1	11
All	All	175/436 (40%)	155 (89%)	16 (9%)	4 (2%)	5	28

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	72	VAL
3	C	30	ALA
3	C	6	VAL
3	C	35	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	61/198 (31%)	61 (100%)	0	100	100
2	B	56/79 (71%)	56 (100%)	0	100	100
3	C	27/39 (69%)	27 (100%)	0	100	100
All	All	144/316 (46%)	144 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

The following chains have linkage breaks:

Mol	Chain	Number of breaks
3	C	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	C	47:PHE	C	62:UNK	N	36.83
1	C	72:UNK	C	99:UNK	N	17.84

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	69/225 (30%)	1.91	25 (36%) 1 3	153, 182, 247, 253	0
2	B	68/103 (66%)	1.77	20 (29%) 1 4	129, 184, 231, 241	0
3	C	44/108 (40%)	2.40	19 (43%) 0 2	168, 215, 328, 332	0
All	All	181/436 (41%)	1.98	64 (35%) 1 3	129, 188, 252, 332	0

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	50	LYS	9.1
1	A	71	ARG	8.5
3	C	4	ALA	7.9
3	C	41	ALA	7.0
3	C	8	GLU	7.0
2	B	45	LYS	6.8
1	A	100	LYS	6.7
3	C	17	GLU	6.7
2	B	47	ILE	6.5
2	B	81	THR	6.3
1	A	115	MET	5.9
2	B	76	HIS	5.8
1	A	91	ARG	5.2
1	A	110	ARG	5.0
2	B	41	ARG	4.9
3	C	15	VAL	4.9
1	A	52	PRO	4.8
2	B	37	ARG	4.7
1	A	51	LEU	4.6
2	B	48	SER	4.2
2	B	44	VAL	4.1
3	C	40	VAL	4.0
3	C	6	VAL	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	C	39	ASP	3.8
3	C	44	ASP	3.8
3	C	42	GLU	3.8
1	A	108	GLU	3.7
2	B	93	ARG	3.6
1	A	95	SER	3.5
1	A	88	LEU	3.3
1	A	64	TYR	3.3
1	A	68	GLU	3.2
3	C	11	LEU	3.1
2	B	80	LYS	3.0
2	B	52	TYR	2.9
1	A	92	LEU	2.9
1	A	75	GLY	2.8
1	A	69	PRO	2.8
3	C	46	LEU	2.7
1	A	114	LEU	2.7
3	C	25	LYS	2.7
1	A	82	GLU	2.6
2	B	38	LEU	2.5
2	B	82	VAL	2.5
1	A	67	ASP	2.5
1	A	63	LYS	2.5
2	B	40	ARG	2.4
3	C	26	MET	2.4
1	A	96	TYR	2.4
2	B	50	LEU	2.4
3	C	31	SER	2.3
1	A	105	VAL	2.3
3	C	5	VAL	2.3
3	C	45	ASN	2.3
1	A	87	TYR	2.2
2	B	79	ARG	2.2
3	C	19	SER	2.2
1	A	61	ILE	2.2
2	B	32	LYS	2.2
1	A	53	PHE	2.1
3	C	33	GLU	2.1
2	B	39	ALA	2.1
2	B	36	ARG	2.0
2	B	91	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.